



MATH
QA
371
H72
1858a

CORNELL
UNIVERSITY
LIBRARIES



Mathematics
Library
White Hall



Cornell University
Library

The original of this book is in
the Cornell University Library.

There are no known copyright restrictions in
the United States on the use of the text.

<http://www.archive.org/details/cu31924059412902>

Production Note

Cornell University Library produced this volume to replace the irreparably deteriorated original. It was scanned using Xerox software and equipment at 600 dots per inch resolution and compressed prior to storage using CCITT Group 4 compression. The digital data were used to create Cornell's replacement volume on paper that meets the ANSI Standard Z39.48-1984. The production of this volume was supported in part by the Commission on Preservation and Access and the Xerox Corporation. Digital file copyright by Cornell University Library 1991.



CORNELL
UNIVERSITY
LIBRARY



GIFT OF

J. E. Oliver
through
Lucien Augustus Wait

MATHEMATICS

A TREATISE

ON

DIFFERENTIAL EQUATIONS,

AND ON THE

CALCULUS OF FINITE DIFFERENCES.

By J. HYMERS, D.D.

LATE FELLOW AND TUTOR OF ST JOHN'S COLLEGE, CAMBRIDGE.

SECOND EDITION, ENLARGED.

LONDON :
LONGMAN, BROWN, GREEN, LONGMANS, AND ROBERTS.
1858.

CAMBRIDGE:
PRINTED BY C. J. OLAY, M.A.
AT THE UNIVERSITY PRESS.

the gift of
J. E. Oliver
through
Lucien Augustus Wait

CONTENTS.

DIFFERENTIAL EQUATIONS.

SECTION I.

DIFFERENTIAL EQUATIONS OF THE FIRST ORDER AND DEGREE.

| ART. | PAGE |
|---|------|
| 1— 5. NATURE and Formation of Differential Equations | 1 |
| 6—10. Exact Differential Equations of the First Order | 4 |
| 11, 12. Homogeneous Equations | 8 |
| 13—15. Linear Equations of the First Order | 11 |
| 16—18. Riccati's Equation, and other Equations, in which the Variables are separable by particular substitutions | 14 |
| 19, 20. Euler's Equation | 19 |
| 21—27. Factors which render integrable a Differential Equation of the First Order | 22 |
| 28. Geometrical Problems producing Differential Equations of the First Order | 30 |

SECTION II.

DIFFERENTIAL EQUATIONS OF THE FIRST ORDER, BUT NOT OF THE FIRST DEGREE.

| | |
|---|----|
| 29. Case in which the Equation can be resolved into its Simple Factors | 35 |
| 30—36. Cases in which this resolution is impossible. Clairaut's Form... | 37 |

SECTION III.

SINGULAR SOLUTIONS OF DIFFERENTIAL EQUATIONS OF THE FIRST ORDER.

| | |
|---|----|
| 37—40. Nature of a Singular Solution; mode of deducing it from the complete Integral; its geometrical signification..... | 47 |
| 41—44. Test of a Given Solution being a Singular Solution. Mode of obtaining the Singular Solutions directly from the Differential Equation | 52 |

SECTION IV.

DIFFERENTIAL EQUATIONS OF THE SECOND ORDER, AND OF HIGHER ORDERS.

| ART. | PAGE |
|---|------|
| 45—48. Cases in which only three of the four quantities $x, y, \frac{dy}{dx}, \frac{d^2y}{dx^2}$ are involved | 59 |
| 49—55. Linear Equations of the Second and Higher Orders | 69 |
| 56—66. Integration of Linear Equations with Constant Coefficients by separation of Symbols | 76 |
| 67. Method of Parameters | 91 |
| 68—76. Method by changing the Independent Variable | 92 |
| 77—82. Simultaneous Equations. Jacobi's Equation | 107 |
| 83. Solutions expressed by Definite Integrals | 115 |
| 84—89. Approximate Solutions of Differential Equations | 117 |

SECTION V.

DIFFERENTIAL EQUATIONS INVOLVING TWO OR MORE INDEPENDENT VARIABLES.

| | |
|---|-----|
| 90—92. Explanation of the Notation used | 126 |
| 93—96. Integration of Differential Functions of two or more Variables... | 128 |
| 97—102. Total Differential Equations | 132 |
| 103—111. Partial Differential Equations of the First Order | 138 |
| 112—117. Partial Differential Equations when linear solved by separation of symbols | 147 |
| 118—120. Change of the independent variables | 155 |
| 121. Partial Differential Equations of a higher degree than the first... | 162 |
| 122—129. Partial Differential Equations of the Second and Higher Orders, not being Linear | 164 |
| 130, 131. Integration by a Series. Simultaneous Equations. | 174 |
| 132, 133. Singular Solutions. Geometrical Problems | 177 |

FINITE DIFFERENCES.

SECTION I.

DIRECT METHOD OF DIFFERENCES.

| ART. | PAGE |
|---|------|
| 1— 6. DEFINITIONS and Principles..... | 1 |
| 7—16. Differences of Explicit Functions | 4 |
| 17—25. Relations between the successive Values and the Differences of a Function | 8 |
| 26—32. Generating Functions | 14 |
| 33—39. Separation of the Symbols of Operation from those of Quantity | 19 |

SECTION II.

INVERSE METHOD OF DIFFERENCES.

| | |
|--|----|
| 40—63. Integration of Explicit Functions | 26 |
| 64—71. Numbers of Bernoulli | 41 |

SECTION III.

EQUATIONS OF DIFFERENCES.

| | |
|---|----|
| 72, 73. Origin and Nature of Equations of Differences | 50 |
| 74. Linear Equations of Differences of the First Order | 51 |
| 75. Indirect Integrals of Equations of Differences..... | 53 |
| 76—85. Linear Equations of Differences of all Orders..... | 55 |
| 86—89. Certain Equations reducible to Linear Equations. Simultaneous Equations | 71 |
| 90—95. Equations of Partial Differences, and Mixed Differences | 80 |
| 96—99. Problems on continued Fractions. Functional Equations | 89 |
| 100. Geometrical Problems involving Finite Differences | 97 |

SECTION IV.

SUMMATION OF SERIES.

| ART. | | PAGE |
|-----------|---|------|
| 101, 102. | By Integrating the General Term | 100 |
| 103—108. | Recurring Series..... | 107 |
| 109—122. | Application of the Integral Calculus to the Summation of Series | 111 |
| 123—133. | Convergency and Divergency of Series..... | 124 |
| 134—142. | Interpolation of Series..... | 132 |

ERRATA.

DIFFERENTIAL EQUATIONS.

| | | | |
|------|--------------|-----------------------------------|-------------------------------------|
| Page | 37, line 16, | for $\int dx \frac{dp}{dz}$ | read $\int dx x \frac{dp}{dz}$. |
| 38, | 5, | $\frac{d}{(1+p^2)^{\frac{1}{2}}}$ | $\frac{a}{(1+p^2)^{\frac{1}{2}}}$. |
| 69, | 2, | Ex. 5 | Ex. 6. |
| 80, | 16, | nn^2 | $11n^2$. |
| 96, | 6, | $\pm p_{n-1}y$ | $\pm p_{n-1} \frac{dy}{dt}$. |
| 96, | 11, | $\pm p_n$ | $\pm p_{n-1} \frac{d}{dt}$. |
| 115, | 14, | a_1^n | a_1^n . |
| — | 17, | a_2^n | a_2^n . |
| 160, | 9, | $m = -1$ | $m = 1$. |

FINITE DIFFERENCES.

| | | | |
|------|-------------|---------------|-----------------|
| Page | 49, line 8, | for Art. 72 | read Art. 67. |
| 98, | 20, | \cos_θ | $\cos \theta$. |

INTEGRATION

OF

DIFFERENTIAL EQUATIONS

BETWEEN TWO OR MORE VARIABLES.

SECTION I.

DIFFERENTIAL EQUATIONS OF THE FIRST ORDER AND DEGREE.

1. IN that part of the Integral Calculus which relates to the integration of explicit functions of one variable, we have to determine the relation between y and x from the equation

$$\frac{dy}{dx} = f(x);$$

in the present portion, we have to determine it from the equation

$$f\left(\frac{d^n y}{dx^n}, \frac{d^{n-1} y}{dx^{n-1}}, \dots, \frac{dy}{dx}, y, x\right) = 0;$$

or to assign the relation between x, y, z (where z is a function of the independent variables x and y), or between a greater number of variables and their functions, from the equation

$$f\left(\frac{dz}{dx}, \frac{dz}{dy}, z, x, y\right) = 0,$$

or from other equations in which a greater number of variables and differential coefficients of higher orders are involved.

2. A differential equation is said to be of the n^{th} order, when the differential coefficient of the highest order which it involves is the n^{th} .

A differential equation of any order is said, moreover, to be of the first, second, &c., degree, when the differential coefficient

which marks its order, is raised to the first, second, &c., power: or when it involves a product at most of m dimensions in differential coefficients and their powers, it is said to be of the m^{th} degree.

To integrate a differential equation of any order, is to pass to the primitive equation between the variables and the constants, from which the proposed may have been derived by the process of differentiation.

3. We shall begin with the simplest case, viz. that of differential equations of the first order and degree, which will be of the form

$$M + N \frac{dy}{dx} = 0,$$

M and N being functions of x and y .

Every differential equation of the first order and degree is either the direct derived equation of a primitive; or it results from the combination of the derived equation with its primitive, so as to eliminate a constant which enters in each only to the first power; the former sort are called exact, the latter inexact.

4. First, let $u = f(x, y) = 0$ be an equation between x and y , by virtue of which y is a function of x ; then, as is proved in the Differential Calculus, $\frac{dy}{dx}$ is given by the equation

$$\frac{du}{dx} + \frac{du}{dy} \cdot \frac{dy}{dx} = 0,$$

the partial differential coefficients $\frac{du}{dx}$, $\frac{du}{dy}$, being formed as if the variables x and y were independent of one another; or, since $\frac{du}{dx}$, $\frac{du}{dy}$, are functions of x and y which we may represent by M and N ,

$$M + N \frac{dy}{dx} = 0,$$

a differential equation of the first order and degree, of which $f(x, y) = 0$ is the primitive or integral. Now if $f(x, y)$,

besides other constants which are affected with x and y , contain a term $+C$ independent of x and y , this will not enter into M and N , having disappeared in differentiating; and if there be no such term, we may add it, and $f(x, y) + C = 0$ is still a relation between x and y which satisfies the equation

$$M + N \frac{dy}{dx} = 0;$$

under this form it is called the complete integral; and the constant C , which does not appear in the differential equation, is called the arbitrary constant; if the integral did not contain such a term as $+C$, it would not be sufficiently general, and would be only a particular case of the complete integral.

We shall presently give the test which every equation of this sort must satisfy, and the mode of integrating it. It is evident that no equation of the first order which is not of the first degree can be exact.

5. Next, let C_1 be another constant which enters to the first power in the equation

$$f(x, y) + C = 0,$$

then C_1 will be affected with x and y , and will consequently appear to the first power in

$$M + N \frac{dy}{dx} = 0;$$

and if a value of C_1 be obtained from either of these equations and substituted in the other, the result will be

$$M_1 + N_1 \frac{dy}{dx} = 0,$$

an equation of the first order and degree, involving all the constants which enter into $f(x, y) + C = 0$, except C_1 . Hence whilst the direct derived equation of

$$f(x, y) + C = 0, \text{ viz. } M + N \frac{dy}{dx} = 0,$$

does not involve the term C which is independent of x and y , there will be as many other differential equations of the first

order and degree that have $f(x, y) + C = 0$ for their primitive, as it has independent constants entering only in the first power; if any constant enter in a dimension above the first, the differential equation obtained by eliminating it, will evidently not be of the first degree.

There are two principal methods of integrating equations of this sort, which consist either in separating the variables, by substitution, or some algebraical process; or in restoring the factor which makes them exact.

Exact Differential Equations of the First Order.

6. Let y be a function of x determined by the equation

$$u = f(x, y) = 0;$$

then the equation which gives the value of $\frac{dy}{dx}$ is

$$\frac{du}{dx} + \frac{du}{dy} \cdot \frac{dy}{dx} = 0, \quad \text{or} \quad M + N \frac{dy}{dx} = 0;$$

the differential coefficients $\frac{du}{dx}$, $\frac{du}{dy}$ being formed on the hypothesis that the variables x and y are independent of one another; then, as proved in the Differential Calculus,

$$\frac{dM}{dy} = \frac{dN}{dx}.$$

Conversely, an equation of the form $M + N \frac{dy}{dx} = 0$ being proposed in which M and N are functions of x and y , if the condition

$$\frac{dM}{dy} = \frac{dN}{dx}$$

(which is called the criterion of integrability) be satisfied, the equation results from the immediate differentiation of an equation of the form $f(x, y) = 0$; and to find its integral amounts to finding a function of two variables $f(x, y)$ whose differential shall be $Mdx + Ndy$, and then to put $f(x, y)$ equal to a constant; if the above condition be not satisfied, there exists no

equation by the simple differentiation of which, the given equation can be produced.

7. To integrate the exact differential equation

$$M + N \frac{dy}{dx} = 0.$$

Let the equation from which it is derived be

$$u = f(x, y) = 0;$$

$$\text{then } \frac{du}{dx} = M, \quad \frac{du}{dy} = N, \quad \text{and } \frac{dM}{dy} = \frac{dN}{dx};$$

$$\therefore u = \int dx M + Y,$$

denoting by Y a function of y which may have disappeared, since M is the differential coefficient of u relative to x , on the hypothesis that x and y are independent;

$$\therefore \frac{du}{dy} = \frac{d}{dy}(\int dx M) + \frac{dY}{dy} = N,$$

$$\therefore \frac{dY}{dy} = N - \frac{d}{dy}(\int dx M), \quad \text{and } Y = \int dy \{N - \frac{d}{dy}(\int dx M)\},$$

$$\therefore u = \int dx M + \int dy \{N - \frac{d}{dy}(\int dx M)\} + C = 0,$$

the complete integral involving one arbitrary constant.

8. OBS. The equation $Y = \int dy \{N - \frac{d}{dy}(\int dx M)\}$ will be absurd, unless the expression $N - \frac{d}{dy}(\int dx M)$ be independent of x : therefore its differential coefficient with respect to x must vanish;

$$\therefore \frac{dN}{dx} - \frac{d}{dx} \frac{d}{dy}(\int dx M) = \frac{dN}{dx} - \frac{d}{dy} \frac{d}{dx}(\int dx M) = \frac{dN}{dx} - \frac{dM}{dy}$$

must equal zero; which it does, since the criterion of integrability is supposed to be satisfied. Hence it will be necessary only to integrate those terms in N which involve y only; and if N can be reduced to such a form as to contain no such terms, the solution will be $u = \int dx M + C = 0$.

9. As the simplest case of exact equations, we may first notice those in which the variables are separated; they will be of the form

$$X + Y \frac{dy}{dx} = 0,$$

where X denotes a function of x only, and Y a function of y only; here the criterion of integrability is manifestly satisfied, for

$$\frac{dX}{dy} = \frac{dY}{dx} = 0;$$

and the complete integral is

$$\int dx X + \int dy Y \frac{dy}{dx} = C, \quad \text{or} \quad \int dx X + \int dy Y = C.$$

To this case may likewise be reduced the equation

$$XY_1 + YX_1 \frac{dy}{dx} = 0,$$

which becomes, when divided by $X_1 Y_1$,

$$\frac{X}{X_1} + \frac{Y}{Y_1} \frac{dy}{dx} = 0.$$

$$\text{Ex. 1.} \quad \frac{1}{\sqrt{a^2 - x^2}} + \frac{1}{\sqrt{a^2 - y^2}} \frac{dy}{dx} = 0;$$

$$\therefore \sin^{-1} \frac{x}{a} + \sin^{-1} \frac{y}{a} = \sin^{-1} \frac{C}{a},$$

$$\text{or} \quad \sin^{-1} \left(\frac{x}{a} \sqrt{1 - \frac{y^2}{a^2}} + \frac{y}{a} \sqrt{1 - \frac{x^2}{a^2}} \right) = \sin^{-1} \frac{C}{a},$$

$$\text{or} \quad x \sqrt{a^2 - y^2} + y \sqrt{a^2 - x^2} = aC;$$

at which we may also arrive, by multiplying the proposed equation by xy , and integrating by parts, which gives

$$-y \sqrt{a^2 - x^2} + \int dx \sqrt{a^2 - x^2} \frac{dy}{dx} - x \sqrt{a^2 - y^2} + \int dx \sqrt{a^2 - y^2} + C = 0,$$

or, since the part affected by the sign $\int dx$ viz.

$$\sqrt{a^2 - y^2} + \sqrt{a^2 - x^2} \frac{dy}{dx},$$

is equal to zero by the proposed,

$$y \sqrt{a^2 - x^2} + x \sqrt{a^2 - y^2} = C.$$

$$\text{Ex. 2. } 1 + y + y^2 + (1 + x + x^2) \frac{dy}{dx} = 0,$$

$$C(x + y + 1) = 2xy + x + y - 1.$$

10. The following are instances of the integration of **exact** differential equations by the method of Art. 7.

$$\text{Ex. 1. } ax + by + c + (bx + my + n) \frac{dy}{dx} = 0.$$

$$\therefore \frac{dM}{dy} = b = \frac{dN}{dx},$$

$$\frac{du}{dx} = ax + by + c; \quad \therefore u = \frac{ax^2}{2} + (by + c)x + Y,$$

$$\frac{du}{dy} = bx + \frac{dY}{dy} = bx + my + n;$$

$$\therefore \frac{dY}{dy} = my + n; \quad \therefore Y = \frac{1}{2} my^2 + ny + C;$$

$$\therefore \frac{1}{2} (ax^2 + my^2) + (by + c)x + ny + C = 0.$$

$$\text{Ex. 2. } \frac{1}{\sqrt{x^2 + y^2}} + \left(\frac{1}{y} - \frac{x}{y \sqrt{x^2 + y^2}} \right) \frac{dy}{dx} + c = 0,$$

$$\log (x + \sqrt{x^2 + y^2}) + cx + C = 0.$$

$$\text{Ex. 3. } \frac{2y - 2x \frac{dy}{dx}}{y \sqrt{x^2 - y^2}} + c = 0, \quad \log \left(\frac{x + \sqrt{x^2 - y^2}}{x - \sqrt{x^2 - y^2}} \right) + cx + C = 0.$$

$$\text{Ex. 4. } \frac{y^2 + ax + (1 - xy) \frac{dy}{dx}}{y^3 + 3ayx - a + a^2 x^2} = 0. \quad \text{Shew this to be exact.}$$

Ex. 5. $(x^2 + y^2 - a^2) \frac{dy}{dx} + (x^2 + 2xy + a^2) = 0,$

$$a^2x + x^2y + \frac{1}{3}x^3 + \frac{1}{3}y^3 - a^2y = C.$$

Ex. 6. $\frac{x^4 \frac{dy}{dx} + x^4 y^2 - a}{x^4(1-xy)^2 - ax^2} = 0$, here $\int dx M + C = 0$,

$$\text{or } \log \frac{\sqrt{a} + x(1-xy)}{\sqrt{a} - x(1-xy)} - \frac{2\sqrt{a}}{x} + C = 0.$$

Homogeneous Equations.

11. We come next to the case of inexact equations, in which the variables are separable by substitution; of these the most important class is homogeneous equations.

Let $M + N \frac{dy}{dx} = 0$ be a homogeneous equation, that is, one in which each of the functions M and N is or can be expressed by series of the form

$$M = ay^m x^{-m} + by^n x^{-n} + cy^p x^{-p} + \&c.,$$

$$N = \alpha y^\mu x^{-\mu} + \beta y^\nu x^{-\nu} + \gamma y^r x^{-r} + \&c.,$$

the sum of the dimensions of x and y in each term of M and N being equal to r .

Let $y = xz$ where z denotes a new function of x , then

$$M = x^r \left(a \frac{y^m}{x^m} + b \frac{y^n}{x^n} + \&c. \right) = x^r f(z),$$

$$N = x^r \left(\alpha \frac{y^\mu}{x^\mu} + \beta \frac{y^\nu}{x^\nu} + \&c. \right) = x^r \phi(z),$$

$$\frac{dy}{dx} = z + x \frac{dz}{dx}.$$

Hence, making these substitutions in the given equation, and dividing by x^r ,

$$f(z) + \phi(z) \left(z + x \frac{dz}{dx} \right) = 0, \quad \text{or } \frac{1}{x} + \frac{1}{\frac{f(z)}{\phi(z)} + z} \cdot \frac{dz}{dx} = 0,$$

in which the variables are separated. Similarly, the variables may be separated by making $x = yz$; and the latter substitution

will be more convenient when N is a more complicated expression than M .

Hence it is easy to effect the separation of the variables in equations which are either homogeneous, or can be made homogeneous; besides these, the number of equations in which that separation is possible, is very limited.

Ex. 1. $3y^2x + 2x^3 + y^3 \frac{dy}{dx} = 0$.

Here $\frac{M}{N} = \frac{3y^2x + 2x^3}{y^3} = 3 \frac{x}{y} + 2 \frac{x^3}{y^3} = \frac{3}{z} + \frac{2}{z^3}$;

$$\therefore \frac{1}{x} + \frac{1}{\frac{3}{z} + \frac{2}{z^3} + z} \cdot \frac{dz}{dx} = 0, \quad \text{or} \quad \frac{1}{x} + \frac{z^3}{z^4 + 3z^2 + 2} \frac{dz}{dx} = 0,$$

$$\text{or} \quad \frac{1}{x} + \left(\frac{2z}{z^2 + 2} - \frac{z}{z^2 + 1} \right) \frac{dz}{dx} = 0;$$

$$\therefore \log x + \log(z^2 + 2) - \frac{1}{2} \log(z^2 + 1) = \log C;$$

$$\therefore \frac{x(z^2 + 2)}{\sqrt{z^2 + 1}} = C; \quad \text{or} \quad y^2 + 2x^2 = C\sqrt{x^2 + y^2}.$$

2. $y^2 + (xy + x^2) \frac{dy}{dx} = 0, \quad y = C \sqrt{1 + 2 \frac{y}{x}}.$

3. $(x - y) \frac{dy}{dx} = x + y, \quad \tan^{-1} \frac{y}{x} = \log \sqrt{\frac{x^2 + y^2}{C^2}}.$

4. $(x^3 - y^2) \frac{dy}{dx} - 2xy = 0, \quad x^2 + y^2 = Cy.$

5. $x \frac{dy}{dx} - y = \sqrt{x^2 - y^2}, \quad \sin^{-1} \frac{y}{x} = \log \frac{x}{C}.$

6. $x \frac{dx}{dy} + y = \sqrt{x^2 + y^2}, \quad 2Cy + C^2 = x^2.$

7. $\sqrt{y} + (\sqrt{y} - \sqrt{x}) \frac{dy}{dx} = 0,$

$$\log(y - \sqrt{xy} + x) + \frac{2}{\sqrt{3}} \tan^{-1} \left(\frac{2\sqrt{x} - \sqrt{y}}{\sqrt{3y}} \right) = C.$$

12. In the following instances, the equations are not homogeneous, but are made so by simple substitutions.

Ex. 1. $ax + by + c + (mx + ny + p) \frac{dy}{dx} = 0.$

Let $ax + by + c = z$, $mx + ny + p = v$,

z and v denoting functions of x ; then $\frac{dy}{dx} = -\frac{z}{v}$, and

$$a + b \frac{dy}{dx} = \frac{dz}{dx}, \quad m + n \frac{dy}{dx} = \frac{dv}{dx};$$

$$\therefore \frac{mv - nz}{av - bz} = \frac{dv}{dz} \text{ or } mv - nz + (bz - av) \frac{dv}{dz} = 0;$$

this being homogeneous, assume $v = zw$, w being a function of z ,

$$\therefore \frac{1}{z} + \frac{(aw - b)}{n - (m + b)w + aw^2} \frac{dw}{dz} = 0,$$

in which the variables are separated.

Ex. 2. $x^m \left(ay + bx \frac{dy}{dx} \right) = y^n \left(ay + \beta x \frac{dy}{dx} \right),$

$$\text{or } \frac{1}{y} (bx^m - \beta y^n) \frac{dy}{dx} = (ay^n - ax^m) \frac{1}{x}.$$

Let $x^m = z$, $y^n = v$, z and v being functions of x ,

$$\therefore \frac{m}{x} = \frac{1}{z} \frac{dz}{dx}, \quad \frac{n}{y} \frac{dy}{dx} = \frac{1}{v} \frac{dv}{dx}, \quad \therefore \frac{nx}{my} \frac{dy}{dx} = \frac{z}{v} \frac{dv}{dz};$$

$$\therefore (bz - \beta v) \frac{mv}{nv} \frac{dv}{dz} = av - az,$$

which is homogeneous.

Ex. 3. $\frac{dy}{dx} + ay^nx^p + by^qx^m = 0$ will become homogeneous by

making $y = z^{\frac{p+1}{1-n}}$, the equation of condition between

$$m, n, p, q, \text{ being } (p+1)(1-q) = (m+1)(1-n).$$

Ex. 4. $\frac{dy}{dx} = \frac{xy^3}{a^2 + xy}$.

This may be written $-\frac{d}{dx} \left(\frac{1}{y} \right) = \frac{x}{\frac{a^2}{y} + x}$,

and therefore becomes homogeneous when z is written for $\frac{1}{y}$.

Linear Equations of the First Order.

13. The next important class of inexact equations of the first order which admit of being integrated, are linear equations, the general form of which is

$$\frac{dy}{dx} + Py = Q,$$

P and Q being functions of x ; they are called linear because they involve no power of y above the first.

Assume $y = vz$, v and z being functions of x ,

$$\therefore v \frac{dz}{dx} + z \frac{dv}{dx} + P vz = Q.$$

Now z being an indeterminate quantity, may be assumed so that the equation last written down may resolve itself into two others, each of which admits of the separation of its variables; to this end let

$$v \frac{dz}{dx} + P zv = 0,$$

or, dividing by v , $\frac{dz}{dx} + Pz = 0$, or $\frac{1}{z} \frac{dz}{dx} + P = 0$;

$$\therefore \log z = -\int dx P, \text{ or } z = e^{-\int dx P}.$$

The remaining part of the equation gives

$$z \frac{dv}{dx} = Q, \text{ or, substituting for } z, \frac{dv}{dx} = Q e^{\int dx P};$$

$$\therefore v = \int dx Q e^{\int dx P} + C,$$

$$\text{and } y = e^{-\int dx P} \{ \int dx Q e^{\int dx P} + C \},$$

the complete primitive involving one arbitrary constant.

OBS. It is unnecessary to add a constant after performing the integration indicated in the equation $z = e^{\int dx P}$; for let

$$z = e^{\int dx P + C} = C_1 e^{\int dx P};$$

$$\text{then } y = zv = C_1 e^{\int dx P} \left\{ \frac{1}{C_1} \int dx Q e^{\int dx P} + C \right\} = e^{\int dx P} (\int dx Q e^{\int dx P} + C C_1),$$

which is the same result as before, since $C C_1$ is equivalent only to a single constant.

14. If we differentiate the result

$$y e^{\int dx P} = \int dx Q e^{\int dx P} + C,$$

$$\text{we get } e^{\int dx P} \left(\frac{dy}{dx} + P y \right) = e^{\int dx P} Q,$$

which shews that if we multiply the proposed equation by $e^{\int dx P}$, each member is separately integrable; and this is the most convenient practical mode of integrating it. When it is once known that the factor which makes the equation integrable is a function of x only, its value may be immediately found; for let it be denoted by X ; then

$$X \frac{dy}{dx} + (Py - Q) X = 0 \text{ is exact,}$$

$$\therefore \frac{dX}{dx} = \frac{d}{dy} (Py - Q) X = PX, \text{ or } \frac{1}{X} \frac{dX}{dx} = P,$$

$$\therefore \log X = \int dx P, \text{ or } X = e^{\int dx P}.$$

15. It must be observed that if the second member of the equation $\frac{dy}{dx} + Py = Q$ be multiplied by any power of y , it is still reducible to the standard form of a linear equation of the first order. For suppose the equation to be

$$\frac{dy}{dx} + Py = Q y^n,$$

then dividing both sides by y^n , and multiplying by $-(n-1)$, we get

$$\frac{d}{dx} \left(\frac{1}{y^{n-1}} \right) - \frac{1}{y^{n-1}} (n-1) P = -Q (n-1).$$

Hence the factor which makes both sides integrable is $e^{-(n-1)\int dx P}$, and the result is

$$\frac{1}{y^{n-1}} = e^{(n-1)\int dx P} \{ - (n-1) \int dx Q e^{-(n-1)\int dx P} + C \}.$$

Ex. 1. $\frac{dy}{dx} - \frac{x}{1+x^2} y = \frac{a}{1+x^2},$

$$P = -\frac{x}{1+x^2}, \quad \int dx P = -\frac{1}{2} \log(1+x^2) = \log \frac{1}{\sqrt{1+x^2}},$$

$$\therefore e^{\int dx P} = \frac{1}{\sqrt{1+x^2}},$$

$$\therefore \frac{y}{\sqrt{1+x^2}} = \int dx \frac{a}{(1+x^2)^{\frac{3}{2}}} + C = \frac{ax}{\sqrt{1+x^2}} + C;$$

$$\therefore y = ax + C\sqrt{1+x^2}.$$

Ex. 2. $\frac{dy}{dx} + y = xy^3,$

$$\text{or, } \frac{d}{dx} \left(\frac{1}{y^3} \right) - \frac{2}{y^3} = -2x,$$

$$P = -2, \quad \int dx P = -2x, \quad e^{\int dx P} = e^{-2x};$$

$$\therefore \frac{e^{-2x}}{y^3} = \int dx e^{-2x} (-2x) = e^{-2x} x - \int dx e^{-2x} = e^{-2x} x + \frac{1}{2} e^{-2x} + C;$$

$$\therefore \frac{1}{y^3} = x + \frac{1}{2} + C e^{2x}.$$

3. $\frac{dy}{dx} - xy = x, \quad y = C e^{\frac{1}{2}x^2} - 1.$

4. $\frac{dy}{dx} + \frac{xy}{1-x^2} = x\sqrt{y}, \quad \sqrt{y} = C(1-x^2)^{\frac{1}{2}} - \frac{1}{2}(1-x^2).$

5. $\frac{dy}{dx} + \frac{1-2x}{x^2} y = 1, \quad y = x^2(1 + C e^{\frac{1}{x}}).$

6. $\frac{dy}{dx} = x^3 y^3 - xy, \quad \frac{1}{y^2} = x^2 + 1 + C e^{x^2}.$

7. $\frac{dy}{dx} = a \sin x + by, \quad y = -a \cdot \frac{b \sin x + \cos x}{1+b^2} + C e^{bx}.$

Riccati's Equation.

16. There are certain cases of the equation

$$\frac{dy}{dx} + by^2 = ax^m$$

(called Riccati's Equation, after the Mathematician who first considered it) in which the variables are separable.

First, let $m = 0$, then $\frac{dy}{dx} = a - by^2$, or $\frac{\frac{dy}{dx}}{a - by^2} = 1$, where the variables are separated.

Secondly, let $m = -2$, and assume $y = \frac{u}{x}$, u being a function of x :

$$\therefore \frac{1}{x} \frac{du}{dx} - \frac{u}{x^2} + \frac{bu^2}{x^2} = \frac{a}{x^3};$$

$$\therefore x \frac{du}{dx} = a + u - bu^2,$$

where the variables are separated.

Thirdly, let $m = -4$, and assume $y = \frac{1}{bx} + \frac{u}{x^2}$,

$$\text{then } -\frac{1}{bx^2} - \frac{2u}{x^3} + \frac{1}{x^3} \frac{du}{dx} + \frac{1}{bx^3} + \frac{2u}{x^3} + \frac{bu^2}{x^4} = \frac{a}{x^4};$$

$$\therefore x^2 \frac{du}{dx} + bu^2 = a,$$

where the variables are separated.

17. Besides the above, the variables are likewise separable in the cases when $m = \frac{-4i}{2i \pm 1}$, i being any integer from 0 to infinity; all which values of m evidently lie between 0 and -2 .

First, let $m = \frac{-4i}{2i-1}$; assume $y = \frac{1}{bx} + \frac{1}{x^2}u$;

$$\therefore by^2 - ax^m = \frac{1}{bx^2} + \frac{2}{x^3u} + \frac{b}{x^4u^2} - ax^m,$$

$$\frac{dy}{dx} = -\frac{1}{bx^2} - \frac{2}{x^3u} - \frac{1}{x^2u^2} \frac{du}{dx};$$

therefore adding these together,

$$\frac{b}{x^4u^2} - ax^m - \frac{1}{x^2u^2} \frac{du}{dx} = 0,$$

$$\text{or } x^2 \frac{du}{dx} + au^2 x^{m+4} - b = 0.$$

Now, let $x = z^{\frac{1}{m+3}}$, then

$$\frac{du}{dx} = \frac{du}{dz} \cdot \frac{dz}{dx} = (m+3) z^{\frac{m+2}{m+3}} \frac{du}{dz};$$

$$\therefore (m+3) z^{\frac{m+4}{m+3}} \frac{du}{dz} + au^2 z^{\frac{m+4}{m+3}} - b = 0;$$

$$\therefore \frac{du}{dz} + \frac{a}{m+3} u^2 = \frac{b}{m+3} z^{-\frac{m+4}{m+3}}, \text{ or } \frac{du}{dz} + b_1 u^2 = a_1 z^{n_1}.$$

$$\text{But } m = \frac{-4i}{2i-1}, \therefore -\frac{m+4}{m+3} = -\frac{4i-4}{2i-3} = -\frac{4(i-1)}{2(i-1)-1} = m_1.$$

Hence by these substitutions the equation is transformed into another of exactly the same form, with $i-1$ instead of i in the index of the variable in the second member.

Similarly, by substituting $\frac{1}{b_1 z} + \frac{1}{u_1 z^2}$ for u , and $z_1^{\frac{1}{m_1+3}}$ for z , we shall transform the equation into another of the same form where $m_2 = \frac{-4(i-2)}{2(i-2)-1}$; and consequently, after i substitutions the index of the variable in the second member will become zero, and the variables will be separated.

Secondly, let $m = \frac{-4i}{2i+1}$; assume $y = \frac{1}{u}$, then

$$-\frac{1}{u^2} \frac{du}{dx} + \frac{b}{u^3} = ax^m, \text{ or } -\frac{du}{dx} + b = ax^m u^2,$$

$$\text{let } x = z^{\frac{1}{m+1}}, \text{ then } \frac{du}{dx} = \frac{du}{dz} \cdot \frac{dz}{dx} = (m+1) z^{\frac{m}{m+1}} \frac{du}{dz};$$

$$\therefore -(m+1) z^{\frac{m}{m+1}} \frac{du}{dz} + b = a z^{\frac{m}{m+1}} u^2,$$

$$\text{or } \frac{du}{dz} + \frac{a}{m+1} u^2 = \frac{b}{m+1} z^{-\frac{m}{m+1}},$$

$$\text{but } m = \frac{-4i}{2i+1}, \quad \therefore -\frac{m}{m+1} = -\frac{-4i}{-2i+1} = \frac{-4i}{2i-1}.$$

Hence by these substitutions this case is reduced to the former; and therefore when $m = \frac{-4i}{2i+1}$, the variables in the equation $\frac{dy}{dx} + by^2 = ax^m$ can be separated. It may be observed that the more general equation, $\frac{dy}{dx} + by^2 x^{q-1} = ax^p$, is reducible to this form by putting $x^q = z$.

18. We shall now give some other instances of equations in which the variables are separable by particular substitutions.

$$\text{Ex. 1. } a \left(x \frac{dy}{dx} - y \right) = \left(x + y \frac{dy}{dx} \right) \sqrt{x^2 + y^2 - a^2}.$$

When an equation contains the expressions

$$y \frac{dy}{dx} + x, \quad x \frac{dy}{dx} - y, \quad \sqrt{x^2 + y^2},$$

the introduction of polar co-ordinates will sometimes effect the separation of the variables; that is, to assume

$$x = \rho \cos \theta, \quad y = \rho \sin \theta,$$

ρ being supposed a function of θ , for then

$$y \frac{dy}{d\theta} + x \frac{dx}{d\theta} = \rho \frac{d\rho}{d\theta}, \quad x \frac{dy}{d\theta} - y \frac{dx}{d\theta} = \rho^2.$$

Hence the proposed equation, which considering x and y as functions of θ , may be written

$$a \left(x \frac{dy}{d\theta} - y \frac{dx}{d\theta} \right) = \left(x \frac{dx}{d\theta} + y \frac{dy}{d\theta} \right) \sqrt{x^2 + y^2 - a^2},$$

becomes $a\rho^2 = \rho \frac{d\rho}{d\theta} \sqrt{\rho^2 - a^2}$, or $a = \frac{1}{\rho} \sqrt{\rho^2 - a^2} \frac{d\rho}{d\theta}$;

$$\therefore a\theta = \sqrt{\rho^2 - a^2} - a \sec^{-1} \frac{\rho}{a} + C,$$

$$\text{or } a \tan^{-1} \frac{y}{x} = \sqrt{x^2 + y^2 - a^2} - a \sec^{-1} \frac{\sqrt{x^2 + y^2}}{a} + C.$$

Ex. 2. $y - x \frac{dy}{dx} + \frac{dy}{dx} \sqrt{(ax + by) \frac{x}{y}} = 0,$

$$\text{or } y \frac{dx}{dy} - x + \sqrt{(ax + by) \frac{x}{y}} = 0;$$

assume $\frac{x}{y} = z$, z being a function of y ,

$$\text{then } y \frac{dx}{dy} - x = y^2 \frac{dz}{dy}; \quad \therefore y^2 \frac{dz}{dy} + \sqrt{(az + b) zy} = 0,$$

where the variables are separated.

Ex. 3. $\frac{dy}{dx} = f(mx + ny)$. Let $mx + ny = z$, then

$$\frac{dz}{dx} = m + n \frac{dy}{dx} = m + nf(z), \text{ where the variables are separated.}$$

Ex. 4. $\frac{dy}{dx} + p + qy + ry^2 = 0$, p , q , and r being functions of x . Having given $y = u$ a particular integral, to find the complete integral.

Assume $y = u + z^{-1}$, then

$$\frac{dz}{dx} - (q + 2ur)z - r = 0, \text{ a linear equation;}$$

$$\therefore \frac{1}{y - u} = z = e^{\int dx(q + 2ur)} \times \int dx re^{-\int dx(q + 2ur)}.$$

Thus for the equation $\frac{dy}{dx} = 1 + (x - y)(ax - by)$,

$$\text{or } \frac{dy}{dx} = 1 + ax^2 - (a + b)xy + by^2,$$

$q = (a+b)x$, $r = -b$, and evidently $u = x$;

$$\therefore \int dx (q + 2ur) = \frac{1}{2} (a-b) x^2;$$

$$\therefore \frac{1}{y-x} = -be^{\frac{1}{2}(a-b)x^2} \times \int dx e^{-\frac{1}{2}(a-b)x^2}.$$

Ex. 5. $(1-xy) \frac{dy}{dx} + y^2 + ax = 0.$

Assume $y = \frac{z-ax^2}{1+xz}$, so that $z = \frac{y+ax^2}{1-xy}$;

$$\text{then } \frac{1}{z^3-a} \cdot \frac{dz}{dx} + \frac{x}{1+ax^3} = 0,$$

where the variables are separated.

Ex. 6. $(y-x) \frac{dy}{dx} = \frac{n(1+y^2)^{\frac{1}{2}}}{\sqrt{1+x^2}}$; assume $y = \frac{x-z}{1+xz}$,

$$\therefore \frac{z}{(1+z^2)(z+n\sqrt{1+z^2})} \frac{dz}{dx} - \frac{1}{1+x^2} = 0.$$

Ex. 7. $\frac{dy}{dx} = \frac{y(n+cx)}{y+a+bx+cx^2}$. Let $z = \frac{y(n+cx)}{y+a+bx+cx^2}$;

$$\therefore \frac{1}{(n-z)^2-b(n-z)+ac} \cdot \frac{1}{z} \frac{dz}{dx} = \frac{1}{(a+bx+cx^2)(n+cx)}.$$

Ex. 8. $\frac{dy}{dx} + p + qy + ry^2 = 0$, where $q = \frac{d}{dx} \log \sqrt{\frac{r}{p}}$.

$$\therefore \left(\frac{dy}{dx} + p + ry^2 \right) \sqrt{\frac{r}{p}} + y \frac{d}{dx} \sqrt{\frac{r}{p}} = 0,$$

$$\text{or } \frac{d}{dx} \left(y \sqrt{\frac{r}{p}} \right) + \left(1 + \frac{y^2 r}{p} \right) \sqrt{\frac{r}{p}} = 0;$$

$$\therefore \tan^{-1} \left(y \sqrt{\frac{r}{p}} \right) + \int dx \sqrt{rp} = C,$$

$$\text{or } y = \sqrt{\frac{p}{r}} \tan (C - \int dx \sqrt{rp}).$$

Thus if we take $p = x^m$, $r = x^n$, then $q = \frac{n-m}{2x}$, and the solution of

$$\frac{dy}{dx} + x^m + \frac{1}{2x} (n-m) y + x^n y^2 = 0, \text{ is}$$

$$y = x^{\frac{m-n}{2}} \tan \left\{ C - \frac{2x^{\frac{1}{2}(m+n+2)}}{m+n+2} \right\}.$$

Euler's Equation.

19. To integrate the equation

$$\frac{dy}{dx} \sqrt{a+bx+cx^2+ex^3+fx^4} + \sqrt{a+by+cy^2+ey^3+fy^4} = 0;$$

or, considering x and y as functions of a new variable t ,

$$\frac{dy}{dt} \sqrt{X} + \frac{dx}{dt} \sqrt{Y} = 0.$$

Let the function of t which expresses x be determined by the equation $\frac{dx}{dt} = \sqrt{X}$, and therefore that which expresses y by the equation $\frac{dy}{dt} = -\sqrt{Y}$; also let $x+y=p$, $x-y=q$, p and q being functions of t .

Then since $\left(\frac{dx}{dt}\right)^2 = X$;

$$\therefore 2 \frac{dx}{dt} \frac{d^2x}{dt^2} = \frac{dX}{dt}, \text{ or } \frac{d^2x}{dt^2} = \frac{1}{2} \frac{dX}{dx};$$

$$\text{similarly } \frac{d^2y}{dt^2} = \frac{1}{2} \frac{dY}{dy}.$$

$$\begin{aligned} \therefore \frac{d^2p}{dt^2} &= \frac{1}{2} \left(\frac{dX}{dx} + \frac{dY}{dy} \right) \\ &= \frac{1}{2} \{ 2b + 2c(x+y) + 3e(x^2+y^2) + 4f(x^3+y^3) \} \\ &= b + c(x+y) + \frac{3e}{4} \{ (x+y)^2 + (x-y)^2 \} \end{aligned}$$

$$+ 2f(x+y) \frac{(x+y)^2 + 3(x-y)^2}{4}$$

$$= b + cp + \frac{3e}{4}(p^2 + q^2) + \frac{1}{2}fp(p^2 + 3q^2),$$

$$\text{and } \frac{dp}{dt} \cdot \frac{dq}{dt} = X - Y = b(x-y) + c(x^2 - y^2) + e(x^3 - y^3) + f(x^4 - y^4)$$

$$= bq + cpq + \frac{e}{4}q(3p^2 + q^2) + \frac{1}{2}fpq(p^2 + q^2);$$

$$\therefore \frac{d^2p}{dt^2} - \frac{1}{q} \frac{dp}{dt} \cdot \frac{dq}{dt} = \frac{e}{2}q^2 + fpq^2,$$

$$\text{or } \frac{d}{dt} \left(\frac{1}{q} \frac{dp}{dt} \right)^2 = e \frac{dp}{dt} + 2fp \frac{dp}{dt};$$

$$\therefore \left(\frac{1}{q} \frac{dp}{dt} \right)^2 = C + ep + fp^2;$$

$$\therefore \frac{dp}{dt} = q \sqrt{C + ep + fp^2},$$

$$\text{or } \sqrt{X} - \sqrt{Y} = (x-y) \sqrt{C + e(x+y) + f(x+y)^2},$$

the integral required. The discovery of this integral, which is due to Euler, was of great importance, as being the first step towards the foundation of the Theory of Elliptic Functions.

20. To integrate the equation

$$\sqrt{1 - c^2 \sin^2 \psi} + \sqrt{1 - c^2 \sin^2 \phi} \frac{d\psi}{d\phi} = 0,$$

or considering ϕ and ψ as functions of another variable t ,

$$\sqrt{1 - c^2 \sin^2 \psi} \frac{d\phi}{dt} + \sqrt{1 - c^2 \sin^2 \phi} \frac{d\psi}{dt} = 0.$$

$$\text{Let } \frac{d\phi}{dt} = \sqrt{1 - c^2 \sin^2 \phi}, \text{ and } \therefore \frac{d\psi}{dt} = -\sqrt{1 - c^2 \sin^2 \psi};$$

$$\therefore \left(\frac{d\phi}{dt} \right)^2 = 1 - c^2 \sin^2 \phi, \quad \left(\frac{d\psi}{dt} \right)^2 = 1 - c^2 \sin^2 \psi;$$

$$\therefore \frac{d^2\phi}{dt^2} + \frac{d^2\psi}{dt^2} = -\frac{1}{2}c^2 (\sin 2\phi + \sin 2\psi),$$

$$\frac{d^2\phi}{dt^2} - \frac{d^2\psi}{dt^2} = -\frac{1}{2}c^2 (\sin 2\phi - \sin 2\psi).$$

Let $p = \phi + \psi, \quad q = \phi - \psi,$

$$\therefore \frac{d^2p}{dt^2} = -c^2 \sin p \cos q,$$

$$\frac{d^2q}{dt^2} = -c^2 \cos p \sin q,$$

and $\frac{dp}{dt} \cdot \frac{dq}{dt} = \left(\frac{d\phi}{dt}\right)^2 - \left(\frac{d\psi}{dt}\right)^2 = -c^2 (\sin^2 \phi - \sin^2 \psi)$

$$= \frac{c^2}{2} (\cos 2\phi - \cos 2\psi) = -c^2 \sin p \sin q;$$

$$\therefore \left(\frac{dp}{dt}\right)^{-1} \cdot \frac{d^2p}{dt^2} = \frac{\cos q}{\sin q} \frac{dq}{dt}; \quad \therefore \log \left(\frac{dp}{dt}\right) = \log (C \sin q);$$

$$\therefore \frac{dp}{dt} = C \sin q; \quad \text{similarly } \frac{dq}{dt} = C \sin p;$$

$$\therefore \sqrt{1 - c^2 \sin^2 \phi} \mp \sqrt{1 - c^2 \sin^2 \psi} = C \sin (\phi \mp \psi)$$

is the integral of the proposed equation; which is only Euler's equation under a different form.

The equations $\sqrt{1 + y^2} + \sqrt{1 + x^2} \frac{dy}{dx} = 0,$

$$\sqrt{y - y^3} + \sqrt{x - x^3} \frac{dy}{dx} = 0,$$

are immediately reducible to the above form, viz.

$$\sqrt{1 - \frac{1}{2} \sin^2 \psi} + \sqrt{1 - \frac{1}{2} \sin^2 \phi} \frac{d\psi}{d\phi} = 0;$$

the former by making $x = \tan \frac{1}{2} \phi, \quad y = \tan \frac{1}{2} \psi;$ the latter by making $\sqrt{x} = \cos \phi, \quad \sqrt{y} = \cos \psi.$

On the Factors which render integrable a Differential Equation of the First Order.

21. The most natural way of obtaining the complete integral of a differential equation of the first order, is to prepare it so that its first member may become an exact differential coefficient; for then we shall have only to integrate and add a constant. This preparation is always possible by means of a factor, when the equation is reduced to the form $\frac{dy}{dx} + K = 0$. For let an equation $f(x, y, C) = 0$ be resolved with respect to C , so that

$$C = \phi(x, y);$$

$$\therefore \text{ by differentiation, } 0 = P + Q \frac{dy}{dx}, \text{ or } \frac{dy}{dx} + \frac{P}{Q} = 0.$$

Now the equation $M + N \frac{dy}{dx} = 0$ may be put under the form $\frac{dy}{dx} + K = 0$, which agrees with the preceding, and may consequently be supposed to have arisen from the elimination of a constant between the primitive $f(x, y, C) = 0$, and its immediately derived equation. On this supposition, therefore, $\frac{dy}{dx} + K = 0$ is identical with $\frac{dy}{dx} + \frac{P}{Q} = 0$;

$$\therefore \frac{dy}{dx} + K = \frac{1}{Q} \left(P + Q \frac{dy}{dx} \right),$$

$$\text{or } \frac{d}{dx} \phi(x, y) = Q \left(\frac{dy}{dx} + K \right), \text{ identically.}$$

The second member therefore is an exact differential coefficient, which proves that there always exists a factor proper to render the expression $\frac{dy}{dx} + K$ integrable.

22. But although the existence of the factor in every case is thus established, the investigation of it is usually attended with greater difficulties than the solution of the original equation.

For let $P + Q \frac{dy}{dx} = 0$ be an exact differential equation; and let z , a function of x and y , be a common factor of P and Q so that $P = Mz$, $Q = Nz$, by the removal of which, the equation is reduced to the inexact state

$$M + N \frac{dy}{dx} = 0;$$

then because $P + Q \frac{dy}{dx} = 0$ is exact,

$$\frac{dP}{dy} = \frac{dQ}{dx}, \quad \text{or} \quad \frac{d(Mz)}{dy} = \frac{d(Nz)}{dx},$$

$$\text{or } z \frac{dM}{dy} + M \frac{dz}{dy} = z \frac{dN}{dx} + N \frac{dz}{dx},$$

$$\text{or } N \frac{dz}{dx} - M \frac{dz}{dy} = z \left(\frac{dM}{dy} - \frac{dN}{dx} \right);$$

an equation between x , y , z , and the partial differential coefficients of z , for determining the factor z . The consideration of this equation in its general state must be reserved till we come to treat of partial differential equations of the first order; but the following particular cases may be noticed.

23. First, suppose that the factor is a function of only one of the variables x , then $\frac{dz}{dy} = 0$, and the equation becomes

$$\frac{1}{z} \frac{dz}{dx} = \frac{1}{N} \left(\frac{dM}{dy} - \frac{dN}{dx} \right),$$

which, being integrated, gives z ; for the hypothesis requires that the second member should be independent of y .

Similarly, if the factor be a function of y only, it will result from the integration of

$$\frac{1}{z} \frac{dz}{dy} = \frac{1}{M} \left(\frac{dN}{dx} - \frac{dM}{dy} \right),$$

of which the second member is independent of x .

Hence, if in any equation $M + N \frac{dy}{dx} = 0$ we find

$$\frac{1}{N} \left(\frac{dM}{dy} - \frac{dN}{dx} \right) = X$$

a function of x only, or

$$\frac{1}{M} \left(\frac{dN}{dx} - \frac{dM}{dy} \right) = Y$$

a function of y only; the factors which make it integrable are respectively $e^{\int dx X}$, $e^{\int dy Y}$.

Ex. 1. $\frac{dy}{dx} + (Py - Q) = 0$, the linear equation of the first order.

This compared with $M + N \frac{dy}{dx} = 0$, gives

$$M = Py - Q, \quad N = 1;$$

$$\therefore \frac{dM}{dy} - \frac{dN}{dx} = P, \quad \text{and} \quad \frac{1}{N} \left(\frac{dM}{dy} - \frac{dN}{dx} \right) = P,$$

a function of x only; therefore the factor is $e^{\int dx P}$.

$$\text{Ex. 2.} \quad y^2 + (1 - xy) \frac{dy}{dx} = 0,$$

$$M = y^2, \quad N = 1 - xy,$$

$$\frac{dN}{dx} - \frac{dM}{dy} = -3y;$$

$$\therefore \int dy \frac{1}{M} \left(\frac{dN}{dx} - \frac{dM}{dy} \right) = - \int dy \frac{3}{y} = \log \frac{1}{y^3};$$

therefore the factor is $\frac{1}{y^3}$.

$$\text{Ex. 3.} \quad (y - x) \frac{dy}{dx} - \frac{1}{2cx} = 0.$$

The factor by which this is made integrable is a function of both the variables $\frac{1}{x} e^{cy^2}$, as may be shewn by introducing it, and applying the criterion of integrability to the equation.

Ex. 4. $\frac{dy}{dx} + \frac{1}{q} \frac{dp}{dx} - \frac{1}{p} \frac{dq}{dx} y^2 = 0,$

where p and q are any functions of x , is made integrable by the factor $\frac{1}{(p + qy)^2}$, and its complete integral is

$$\frac{1}{q(p + qy)} + \int dx \frac{1}{pq^2} \frac{dq}{dx} = C.$$

24. In the case of homogeneous equations, a factor proper to render them integrable, is readily discovered by means of the property that if u be a homogeneous function of n dimensions of the independent quantities t and z , then

$$nu = t \frac{du}{dt} + z \frac{du}{dz}.$$

For suppose V , a homogeneous function of x and y of m dimensions, to be a factor which makes $M + N \frac{dy}{dx}$ an exact differential coefficient, M and N being homogeneous functions of x and y of r dimensions; then if U denote the primitive function, it will be homogeneous and of $m + r + 1$ dimensions, and we shall have

$$VM + VN \frac{dy}{dx} = \frac{dU}{dx};$$

hence since VM and VN are the partial differential coefficients of U with respect to x and y respectively,

$$xVM + yVN = (m + r + 1) U;$$

$$\therefore \frac{M + N \frac{dy}{dx}}{Mx + Ny} = \frac{1}{m + r + 1} \cdot \frac{1}{U} \frac{dU}{dx},$$

and as the second member is an exact differential coefficient, .. it follows that the first is so likewise, and consequently, that $M + N \frac{dy}{dx}$ is made exact by means of the multiplier

$$\frac{1}{Mx + Ny}.$$

25. The property of homogeneous functions assumed above is easily proved. Let u be a homogeneous function of the independent quantities t and z of n dimensions; then if we change t into $t(1+h)$ and z into $z(1+h)$, u will become

$$u(1+h)^n = u + nuh + \&c.$$

But by Taylor's theorem, u will also become

$$u + \frac{du}{dt} \cdot ht + \frac{du}{dz} \cdot hz + \&c.,$$

therefore, equating the coefficients of h ,

$$nu = t \frac{du}{dt} + z \frac{du}{dz}.$$

And, generally, if u be a homogeneous function of n dimensions of any number of independent quantities $t, z, w, \&c.$, and we change them into $t(1+h), z(1+h), w(1+h), \&c.$, the new value of u will be equally expressed by $u(1+h)^n$ or by $e^{(t \frac{d}{dt} + z \frac{d}{dz} + \dots) \log(1+h)} u$; and equating the coefficients of h^r in these two identical expressions, we get, separating as above the symbols of operation from those of quantity,

$$n(n-1) \dots (n-r+1) u = \left(t \frac{d}{dt} + z \frac{d}{dz} + w \frac{d}{dw} + \dots \right)^r u.$$

Ex. $xy + y^2 + (xy - x^2) \frac{dy}{dx} = 0.$

The factor is

$$\frac{1}{(xy + y^2)x + (xy - x^2)y} = \frac{1}{2y^2x};$$

$$\therefore \frac{xy + y^2}{2y^2x} + \frac{xy - x^2}{2y^2x} \frac{dy}{dx} = 0,$$

is an exact differential coefficient, and gives the primitive by Art. 7.

$$\frac{x}{2y} + \frac{1}{2} \log(xy) + C = 0.$$

26. Whenever the variables can be separated in an equation, a factor which makes it integrable can also be found.

For suppose that $M + N \frac{dy}{dx} = 0$, by the introduction of two other variables u and z , is transformed into $R + S \frac{du}{dz} = 0$, so that

$$M + N \frac{dy}{dx} = R + S \frac{du}{dz};$$

and suppose V to be a function of u and z , such that if we divide $R + S \frac{du}{dz}$ by it, the variables are separated, i.e. $\frac{R}{V}$ contains z only, and $\frac{S}{V}$ contains u only;

$$\therefore \frac{1}{V} \left(M + N \frac{dy}{dx} \right) = \frac{R}{V} + \frac{S}{V} \frac{du}{dz}$$

is an exact differential coefficient; and consequently $\frac{1}{V}$, which, upon restoring the values of u and z , becomes a function of x and y , is a factor which makes $M + N \frac{dy}{dx} = 0$ integrable.

Ex. 1. $a + bx^2y^2 + x^2 \frac{dy}{dx} = 0$.

Assuming $y = \frac{u}{x}$, we find

$$a + bx^2y^2 + x^2 \frac{dy}{dx} = a + bu^2 + x \frac{du}{dx} - u,$$

and dividing by $x(a - u + bu^2)$, we get

$$\frac{a + bx^2y^2 + x^2 \frac{dy}{dx}}{x(a - u + bu^2)} = \frac{1}{x} + \frac{\frac{du}{dx}}{a - u + bu^2};$$

$$\therefore \frac{1}{x(a - u + bu^2)} = \frac{1}{ax - x^2y + bx^3y^2}$$

is a factor which makes the proposed equation integrable.

Ex. 2. $\frac{dy}{dx} + y^2 - \frac{a}{x^4} = 0.$

The integrating factor here is $\frac{x^2}{x^2(1-xy)^2 - a}$; as results from the substitution by which (Art. 16) the variables are separated.

Ex. 3. $y^2 + ax + (1-xy)\frac{dy}{dx} = 0.$

By assuming $y = \frac{z - ax^2}{1 + xz}$, it may be shewn that a factor which makes the proposed integrable is

$$\frac{1}{y^3 + 3axyx - a + a^2x^3}.$$

Ex. 4. $M + N\frac{dy}{dx} = 0$, a homogeneous equation.

In this case we know, that making $y = xz$, we have

$$M = x^r f(z), \quad N = x^r \phi(z),$$

$$\text{and } M + N\frac{dy}{dx} = x^r f(z) + x^r \phi(z) \left(z + x \frac{dz}{dx} \right);$$

consequently, dividing by

$$x^{r+1} \{f(z) + z\phi(z)\} = Mx + Ny,$$

we get

$$\frac{M + N\frac{dy}{dx}}{Mx + Ny} = \frac{1}{x} + \frac{\phi(z)\frac{dz}{dx}}{f(z) + z\phi(z)};$$

$$\therefore \frac{1}{Mx + Ny}$$

is a factor which makes the proposed equation integrable.

OBS. That $\frac{M + N\frac{dy}{dx}}{Mx + Ny}$ is an exact differential coefficient, provided M and N be homogeneous functions of x and y of the same dimensions, admits of an easy proof as follows.

We must shew that

$$\frac{d}{dy} \left(\frac{M}{Mx + Ny} \right) = \frac{d}{dx} \left(\frac{N}{Mx + Ny} \right).$$

Now putting $\frac{y}{x} = z$, we have

$$\begin{aligned} \frac{M}{Mx + Ny} &= \frac{1}{x} \cdot \frac{1}{1 + \frac{Ny}{Mx}} = \frac{1}{x} \cdot \frac{1}{1 + F(z)}, \\ \frac{N}{Mx + Ny} &= \frac{1}{y} \left(1 - \frac{Mx}{Mx + Ny} \right) = \frac{1}{y} - \frac{1}{y} \cdot \frac{1}{1 + F(z)}; \\ \therefore \frac{d}{dy} \left(\frac{M}{Mx + Ny} \right) &= -\frac{1}{x} \cdot \frac{\frac{dF(z)}{dz} \cdot \frac{1}{x}}{\{1 + F(z)\}^2}, \\ \frac{d}{dx} \left(\frac{N}{Mx + Ny} \right) &= -\frac{1}{y} \cdot \frac{\frac{dF(z)}{dz} \cdot \frac{y}{x^2}}{\{1 + F(z)\}^2}, \end{aligned}$$

which expressions are evidently equal to one another.

27. There always exists an infinite number of factors which render an equation of the first order and degree integrable.

For let z be the factor by means of which the equation

$$M + N \frac{dy}{dx} = 0$$

is made integrable, and $u=0$ its complete primitive, so that

$$z \left(M + N \frac{dy}{dx} \right) = \frac{du}{dx};$$

multiply both sides by $F(u)$, where $F(u)$ denotes any function of u , and we have

$$zF(u) \left(M + N \frac{dy}{dx} \right) = F(u) \frac{du}{dx};$$

and as the second member is an exact differential coefficient, it follows that the first is so likewise; therefore $zF(u)$ is a factor which makes the proposed equation integrable, whatever form be assigned to $F(u)$.

Geometrical Problems producing Equations of the first order and degree.

28. The following geometrical problems are added to illustrate this part of the subject.

I. To determine the trajectory of a given family of curves.

Let AA' , BB' (fig. 1) be two of a family of curves resulting from the equation $f(X, Y, c) = 0$, by giving particular values to the constant c ; and let AB be a curve which cuts AA' , BB' , and all the curves resulting from the equation by giving all possible values to c , at the same angle; then AB is called the trajectory of this family of curves. Let x, y , be the co-ordinates of the point A in AB , between which we are required to find a relation; AT' , AT , tangents to the curve and trajectory at A , $\tan TAT' = a$; and let a value $\psi(X, Y)$ of $\frac{dY}{dX}$ be obtained from the equation $f(X, Y, c) = 0$, not involving c ; then at the point A , $\tan AT'N = \psi(x, y)$, and $\tan ATN = \frac{dy}{dx}$,

$$\therefore a = \frac{\frac{dy}{dx} - \psi(x, y)}{1 + \frac{dy}{dx} \cdot \psi(x, y)},$$

the differential equation to the curve AB ; and as it does not involve c , AB will cut every curve in the series at an angle whose tangent $= a$. The equation when integrated will involve an arbitrary constant, and consequently will represent a system of curves, every one of which cuts the former system at the same angle; the constant may be determined, if a point through which the trajectory is to pass, be given. If the angle TAT' be a right angle, or a be infinite, the differential equation to the trajectory, which is then called orthogonal, is

$$1 + \frac{dy}{dx} \cdot \psi(x, y) = 0.$$

Ex. 1. To find the orthogonal trajectory to all curves resulting from the equation $y^2(c - x) = x^3$, by giving all possible values to c .

$$\text{Here } c - x = \frac{x^3}{y^3}, \quad \therefore -1 = \frac{3x^2}{y^3} - \frac{2x^3}{y^3} \frac{dy}{dx};$$

$$\therefore \frac{dy}{dx} = \psi(x, y) = \frac{y^3 + 3x^2y}{2x^3},$$

therefore, substituting for $\psi(x, y)$ its value, the differential equation to the trajectory is

$$2x^3 + (y^3 + 3x^2y) \frac{dy}{dx} = 0;$$

a homogeneous equation whose integral is (Art. 11)

$$x^2 + y^2 = C\sqrt{2x^2 + y^2}.$$

Similarly, let $f(\rho, \theta, c) = 0$ be the equation to the curve AA' referred to polar co-ordinates, and let it give for $\rho \frac{d\theta}{d\rho}$ the value $\psi(\rho, \theta)$ independent of c ; then considering ρ and θ as co-ordinates of the point A in the curve AB ,

$$\tan SAT' = \psi(\rho, \theta), \quad \tan SAT = \rho \frac{d\theta}{d\rho},$$

$$\therefore a = \frac{\psi(\rho, \theta) - \rho \frac{d\theta}{d\rho}}{1 + \psi(\rho, \theta) \rho \frac{d\theta}{d\rho}},$$

which is the differential equation to the trajectory; or if it be orthogonal,

$$1 + \psi(\rho, \theta) \rho \frac{d\theta}{d\rho} = 0.$$

EX. 2. Let the curves be a system of circles touching a straight line in the same point, then taking that point as the origin and measuring θ from the line, their equation is

$$\rho = c \sin \theta,$$

$$\therefore \frac{1}{\rho} = \frac{\cos \theta}{\sin \theta} \frac{d\theta}{d\rho}, \quad \text{or } \rho \frac{d\theta}{d\rho} = \frac{\sin \theta}{\cos \theta} = \psi(\rho, \theta);$$

$$\therefore 1 + \frac{\sin \theta}{\cos \theta} \rho \frac{d\theta}{d\rho} = 0, \quad \text{or } \cos \theta + \sin \theta \rho \frac{d\theta}{d\rho} = 0,$$

$$\text{or } \frac{d}{d\rho} \left(\frac{\rho}{\cos \theta} \right) = 0; \quad \therefore \rho = C \cos \theta,$$

the equation to the orthogonal trajectory, which represents a system of circles passing through the given point and having the given line for their diameter.

We may generalize this problem, by finding the orthogonal trajectory of all circles described through two given points.

II. To determine a curve such, that the locus of the extremity of its polar subtangent shall be a straight line.

The polar subtangent is a line drawn from the origin perpendicular to the radius vector to meet the tangent.

Let ρ , θ , be the polar co-ordinates of any point P in the curve sought (fig. 2); then those of the extremity T of its polar subtangent will be $\rho^2 \frac{d\theta}{d\rho}$ and $\theta - \frac{\pi}{2}$, which must satisfy the equation to a straight line, viz.

$$\rho' = c \sec (\theta' - \alpha);$$

$$\therefore \rho^2 \frac{d\theta}{d\rho} = c \sec \left(\theta - \alpha - \frac{\pi}{2} \right) = \frac{c}{\sin (\theta - \alpha)}.$$

$$\therefore \frac{c}{\rho^2} = \sin (\theta - \alpha) \frac{d\theta}{d\rho}, \quad \text{and } \frac{c}{\rho} = \cos (\theta - \alpha) + C;$$

$$\therefore \rho = \frac{c}{C + \cos (\theta - \alpha)},$$

the equation to curves of the second order, having the pole for one of their foci.

III. To find a curve in which SG varies as SP , PG being a normal at P , and SG a fixed line through S , (fig. 2).

Taking SG for the axis of x , the equation to the normal at P is

$$(Y - y) \frac{dy}{dx} + X - x = 0;$$

$$\text{therefore making } Y = 0, \quad X = SG = x + y \frac{dy}{dx},$$

$$\text{and } SP = \sqrt{x^2 + y^2}, \quad \therefore x + y \frac{dy}{dx} = e \sqrt{x^2 + y^2},$$

$$\text{or } \sqrt{x^2 + y^2} = ex + C,$$

the equation to a curve of the second order.

IV. To find a curve which is always cut by its radius vector at an angle proportional to the corresponding angle of revolution; that is, $\angle SPT \propto \angle ASP$, (fig. 2).

Let ρ, θ , be the co-ordinates of any point in the curve, then the angle at which the radius vector cuts the curve, has for its tangent $\rho \frac{d\theta}{d\rho}$;

$$\therefore \rho \frac{d\theta}{d\rho} = \tan n\theta, \quad \text{or } \frac{1}{\rho} \frac{d\rho}{d\theta} = \frac{\cos n\theta}{\sin n\theta}, \quad \therefore \left(\frac{\rho}{C}\right)^n = \sin n\theta.$$

V. To find the locus of the centre of an ellipse rolling along a straight line.

Let ON be the line along which the ellipse rolls, and which touches the ellipse at P ; $ON = x$, $NC = y$, the co-ordinates of its centre; then CP is a normal to the locus of C (fig. 7), and therefore

$$= y \sqrt{1 + \left(\frac{dy}{dx}\right)^2},$$

$$\text{but } a^2 + b^2 - CP^2 = \frac{a^2 b^2}{CN^2},$$

$$\therefore a^2 + b^2 - y^2 - y^2 \left(\frac{dy}{dx}\right)^2 = \frac{a^2 b^2}{y^2}, \quad \text{or } \frac{dx}{dy} = \frac{y^2}{\sqrt{(a^2 - y^2)(y^2 - b^2)}}$$

the differential equation to the required curve, in which the variables are separated. Generally, if $p = f(\rho)$ be the relation between the radius vector and perpendicular on the tangent in any curve, then the locus of the pole, when the curve rolls along a straight line, will have for its equation

$$y = f\left\{y \sqrt{1 + \left(\frac{dy}{dx}\right)^2}\right\}.$$

Thus if the curve be a parabola, to find the locus of its focus we have $p = \sqrt{ap}$;

$$\therefore y = \sqrt{ay \left\{ 1 + \left(\frac{dy}{dx} \right)^2 \right\}^{\frac{1}{2}}}, \quad \text{or } y = a \sqrt{1 + \left(\frac{dy}{dx} \right)^2},$$

the equation to the common catenary.

VI. To find a curve such that the intersection of the tangent at any point and a line drawn from the pole inclined at a constant angle to the radius vector of that point, shall trace out a given curve.

Let $\angle ASP = \theta$, $SP = \rho$ (fig. 2) be the polar co-ordinates of any point P in the required curve, YQ a tangent at P intersected by a perpendicular upon it from S in Y , and by a line through S inclined at a constant angle $PSQ = \alpha$ to SP in Q . Also let $\angle ASQ = \theta'$, $SQ = \rho'$ be the co-ordinates of Q , which is supposed to lie in a given curve whose equation is $\frac{1}{\rho'} = f(\theta')$; then $\theta' = \theta + \alpha$; and, calling $\angle PSY = \phi$,

$$\frac{1}{\rho'} = \frac{\cos(\alpha + \phi)}{\rho \cos \phi} = \frac{\cos \alpha}{\rho} - \frac{\sin \alpha}{\rho^2} \frac{d\rho}{d\theta}, \quad \text{since } \cot \phi = \frac{1}{\rho} \frac{d\rho}{d\theta}.$$

Hence the equation to the locus of P is

$$\frac{d}{d\theta} \left(\frac{1}{\rho} \right) + \frac{\cot \alpha}{\rho} = \frac{1}{\sin \alpha} f(\theta + \alpha),$$

$$\text{or } \frac{1}{\rho} e^{\theta \cot \alpha} + C = \frac{1}{\sin \alpha} \int d\theta e^{\theta \cot \alpha} f(\theta + \alpha).$$

Suppose $\frac{a}{\rho'} = \frac{\cos \beta + c \cos \theta'}{a}$, a curve of the second order;

then $\frac{a}{\rho} + C e^{-\theta \cot \alpha} = \frac{\cos \beta}{\cos \alpha} + c \cos \theta$, another curve of the second order when $C = 0$. The result seems to fail when $\alpha = \frac{1}{2} \pi$; but in that case, first changing the constant into $C' + \frac{\cos \beta}{\cos \alpha}$, we find

$$\frac{a}{\rho} + C' - \theta \cos \beta = c \cos \theta.$$

SECTION II.

DIFFERENTIAL EQUATIONS OF THE FIRST ORDER BUT NOT OF THE FIRST DEGREE.

29. WHEN a differential equation of the first order is of a higher degree than the first, we know that it is not obtained by the direct differentiation of its primitive, but results from eliminating a constant, (which enters into the primitive in a dimension above the first,) between the primitive and its derived equation; the degree of the differential equation and the dimension of the constant eliminated above the lowest dimension in which it appears, being always the same. The general form of such equations free from radicals, is

$$\left(\frac{dy}{dx}\right)^n + p_1 \left(\frac{dy}{dx}\right)^{n-1} + p_2 \left(\frac{dy}{dx}\right)^{n-2} + \dots + p_{n-1} \frac{dy}{dx} + p_n = 0,$$

the coefficients being functions of x and y .

If this can be resolved with respect to $\frac{dy}{dx}$ into its n simple factors, it will assume the form

$$\left(\frac{dy}{dx} + q_1\right) \left(\frac{dy}{dx} + q_2\right) \dots \left(\frac{dy}{dx} + q_n\right) = 0;$$

then each of these factors put equal to zero, will be an equation of the first order and degree, whose integral may be found by the methods of the preceding Section; and any one of these integrals, as well as the continued product of any number of them, will evidently satisfy the proposed equation. If, therefore, we integrate the n equations,

$$\frac{dy}{dx} + q_1 = 0, \quad \frac{dy}{dx} + q_2 = 0, \quad \dots \quad \frac{dy}{dx} + q_n = 0,$$

and complete them all with the same constant C , as the proposed equation is of the first order, we shall obtain the required primitive involving C in the n^{th} power, by equating their continued product to zero.

$$\text{Ex. 1.} \quad \left(\frac{dy}{dx}\right)^2 + \frac{2x}{y} \frac{dy}{dx} - 1 = 0;$$

$$\therefore \frac{dy}{dx} + \frac{x}{y} = \pm \sqrt{1 + \frac{x^2}{y^2}}, \quad \text{or} \quad \frac{y \frac{dy}{dx} + x}{\sqrt{y^2 + x^2}} = \pm 1;$$

$$\therefore +\sqrt{y^2 + x^2} = x + C, \quad \text{and} \quad -\sqrt{y^2 + x^2} = x + C;$$

$$\therefore (\sqrt{y^2 + x^2} - C - x)(\sqrt{y^2 + x^2} + C + x) = 0,$$

$$\therefore y^2 + x^2 - (C + x)^2 = 0, \quad \text{or} \quad y^2 = 2Cx + C^2.$$

$$\text{Ex. 2.} \quad x^2 \left(\frac{dy}{dx}\right)^2 - 2xy \frac{dy}{dx} + y^2 - x^2 y^2 - x^4 = 0.$$

$$\therefore x \frac{dy}{dx} - y = \pm x \sqrt{x^2 + y^2}, \quad \text{or} \quad \frac{\frac{d}{dx} \left(\frac{y}{x}\right)}{\pm \sqrt{1 + \frac{y^2}{x^2}}} = 1.$$

$$\therefore \log \left(\frac{y}{x} + \sqrt{1 + \frac{y^2}{x^2}}\right) = x + C,$$

$$\text{and} \quad -\log \left(\frac{y}{x} + \sqrt{1 + \frac{y^2}{x^2}}\right) = \log \left(\sqrt{1 + \frac{y^2}{x^2}} - \frac{y}{x}\right) = x + C;$$

$$\therefore \sqrt{x^2 + y^2} + y = cxe^x, \quad \text{changing the constant,}$$

$$\text{and} \quad \sqrt{x^2 + y^2} - y = cxe^x;$$

$$\therefore (\sqrt{x^2 + y^2} - cxe^x + y)(\sqrt{x^2 + y^2} - cxe^x - y) = 0,$$

$$\text{or} \quad y = \frac{x}{2} \left(\frac{1}{ce^x} - ce^x\right).$$

$$\text{Ex. 3.} \quad y^2 + \left(y \frac{dy}{dx}\right)^2 = 2nx + 2ny \frac{dy}{dx}.$$

$$y^2 + (x - c)^2 = n^2 + 2nx,$$

the equation to a circle.

This is the solution of the Problem to find a curve in which the square of the normal is always proportional to the sum of the abscissa and subnormal.

30. When the resolution of the proposed equation into its simple factors is impossible, there are still various forms for which the complete primitive can be determined, or its determination made to depend on elimination; this is done by means of substitution, or differentiation, or other analytical artifices, of which we shall now give some instances.

OBS. The arbitrary constant in what follows is often reserved under sign of integration.

31. When the equation contains only one of the variables, x suppose, and can be solved with respect to that variable, so that $x = f\left(\frac{dy}{dx}\right)$; let $\frac{dy}{dx}$ be denoted by p , then $x = f(p)$; and integrating the equation $\frac{dy}{dx} = p$ by parts, we get

$$y = xp - \int dx \frac{dp}{dx} = pf(p) - \int p f'(p) dp;$$

between which and the equation $x = f(p)$, eliminating p , we shall obtain the required integral.

Similarly, if we have $y = f(p)$, since

$$\frac{dx}{dp} = \frac{dx}{dy} \cdot \frac{dy}{dp} = \frac{1}{p} \frac{df(p)}{dp},$$

we shall have to eliminate p between $y = f(p)$,

$$\text{and } x = \int dp \frac{1}{p} \frac{df(p)}{dp}.$$

Ex. 1. $x + x \left(\frac{dy}{dx}\right)^2 = 1$, or $x = \frac{1}{1+p^2}$;

$$\therefore y = \frac{p}{1+p^2} - \tan^{-1} p + C;$$

$$\therefore y = \sqrt{x(1-x)} - \tan^{-1} \sqrt{\frac{1-x}{x}} + C.$$

Ex. 2. $y = a \sqrt{1+p^2}$, $x + C = a \log (\sqrt{y^2 - a^2} + y)$.

This is the solution of the problem in which it is required to find a curve such that the perpendicular on the tangent from the foot of the ordinate shall be constant.

Ex. 3. $y \sqrt{1+p^2} = ap$. Here $\frac{dy}{dp} = \frac{d}{(1+p^2)^{\frac{3}{2}}}$;

$$\therefore x + C = \int dp \frac{a}{p(1+p^2)^{\frac{3}{2}}} = \frac{a}{\sqrt{1+p^2}} + a \log \left(\frac{p}{\sqrt{1+p^2} + 1} \right),$$

$$\text{or } x + C = \sqrt{a^2 - y^2} + a \log \left(\frac{y}{\sqrt{a^2 - y^2} + a} \right).$$

This is the solution of the problem to find a curve such that the tangent terminated at the axis of x shall be of a constant length.

32. An equation not coming immediately under this case, may sometimes be reduced to it by putting $p = xz$, or $p = yz$.

Ex. $\left(\frac{dy}{dx}\right)^3 + ax \frac{dy}{dx} + x^3 = 0$. Let $p = xz$,

$$\text{then } x(z^3 + 1) + az = 0, \text{ or } x = -\frac{az}{1+z^3};$$

$$\therefore \frac{dy}{dz} = \frac{dy}{dx} \cdot \frac{dx}{dz} = -\frac{a^2 z^2 (2z^3 - 1)}{(z^3 + 1)^3},$$

and z must be eliminated between the integral of this, and the equation $x = -\frac{az}{1+z^3}$.

33. When the equation contains both the variables x and y , provided it be homogeneous with respect to them, we may assume $\frac{y}{x} = z$; then the equation will take the form, (which is not solvable with respect to p by supposition,)

$$f(z, p) = 0.$$

Suppose this capable of being solved with respect to z , and let it give $z = \phi(p)$; now $y = xz$ gives $p = z + x \frac{dz}{dx}$,

$$\text{or } \frac{1}{x} = \frac{1}{p - z} \frac{dz}{dx},$$

substitute the above value of z , and integrate this equation; then p must be eliminated between the result which will be of the form $\log x = F(p)$, and $y = x\phi(p)$.

$$\text{Ex. } y - x \frac{dy}{dx} = nx \sqrt{1 + \left(\frac{dy}{dx}\right)^2};$$

$$\therefore \frac{y}{x} = p + n \sqrt{1 + p^2};$$

$$\therefore \frac{1}{x} = \frac{\frac{d}{dx}(p + n \sqrt{1 + p^2})}{-n \sqrt{1 + p^2}} = -\frac{1}{n} \left(\frac{\frac{dp}{dx}}{\sqrt{1 + p^2}} + n \frac{\frac{d}{dx} \sqrt{1 + p^2}}{\sqrt{1 + p^2}} \right);$$

$$\therefore \log x = -\frac{1}{n} \{ \log(p + \sqrt{1 + p^2}) + n \log \sqrt{1 + p^2} \} + \log C;$$

$$\therefore x = \frac{C(p + \sqrt{1 + p^2})^{-\frac{1}{n}}}{\sqrt{1 + p^2}}, \quad \frac{y}{x} = p + n \sqrt{1 + p^2},$$

between which equations p must be eliminated.

This is the solution of the problem, to find a curve such that the perpendicular upon the tangent from the origin shall vary as the abscissa to the point of contact.

$$\text{Let } n = 1. \quad \frac{C}{x} = 1 + p^2 + p \sqrt{1 + p^2}, \quad \frac{y}{x} = p + \sqrt{1 + p^2};$$

$$\therefore \frac{y^2 + (C - x)^2}{x^2} = (1 + p^2)(p + \sqrt{1 + p^2})^2 = \frac{C^2}{x^2},$$

or $y^2 = 2Cx - x^2$, the equation to a circle.

34. Another integrable form is $y = x \frac{dy}{dx} + f\left(\frac{dy}{dx}\right)$, which

is called Clairaut's form, after the Mathematician who first considered it. Substituting p for $\frac{dy}{dx}$, and differentiating, we get successively,

$$y = xp + f(p),$$

$$\frac{dy}{dx} = p = p + x \frac{dp}{dx} + \frac{d}{dp} f(p) \cdot \frac{dp}{dx};$$

$$\therefore \{x + \frac{d}{dp} f(p)\} \frac{dp}{dx} = 0,$$

which resolves itself into the two

$$x + \frac{d}{dp} f(p) = 0, \quad \frac{dp}{dx} = 0.$$

The first of these gives $p = \phi(x)$ suppose; this value substituted for p in the proposed equation, furnishes a relation between x and y which satisfies the proposed equation, but which involves no arbitrary constant, and cannot therefore be the complete primitive. The other equation must therefore lead to the complete primitive; but this gives $p = C$, and by substituting this value of p in the proposed we find

$$y = Cx + f(C).$$

Hence Clairaut's form has the property, that the complete primitive is obtained by substituting the arbitrary constant C for p , in that form. If we integrate $p = C$, we find $y = Cx + C'$; but the condition of the proposed equation being satisfied gives $C' = f(C)$, the same result as before.

We shall afterwards return to the consideration of the other solution, which is called the singular solution, and is not derivable from the complete integral. It is evident that Clairaut's form may be put into the rather more general shape

$$y = (x + c)p + f(p).$$

Ex. 1. $y = xp + \frac{\alpha(1+p^2)}{p}.$

Differentiating, we get $p = p + x \frac{dp}{dx} + a \left(1 - \frac{1}{p^2}\right) \frac{dp}{dx}$,

$$\text{or } \left(x + a - \frac{a}{p^2}\right) \frac{dp}{dx} = 0;$$

$$\therefore \frac{dp}{dx} = 0 \text{ gives } p = C, \text{ and } y = Cx + \frac{a(1+C^2)}{C},$$

the complete integral; and

$$x + a - \frac{a}{p^2} = 0 \text{ gives } p = \pm \sqrt{\frac{a}{x+a}},$$

which, substituted in $py = a + p^2(x+a)$, gives the singular solution

$$\pm y \sqrt{\frac{a}{x+a}} = 2a, \text{ or } y^2 = 4a(x+a).$$

$$\text{Ex. 2. } y = x(p-c) + \sqrt{a^2 - c^2 + p^2 a^2},$$

$$y = x(C-c) + \sqrt{a^2 - c^2 + C^2 a^2}, \text{ the complete integral;}$$

$$\frac{y^2}{a^2 - c^2} + \frac{(x-c)^2}{a^2} = 1, \text{ the singular solution.}$$

The two foregoing examples are the solutions of the problems to find a curve for which the locus of the intersection of the tangent and perpendicular upon it from the origin shall be a straight line, and a circle, respectively. For the co-ordinates of the said point of intersection are

$$X = -\frac{p(y - px)}{1 + p^2}, \quad Y = \frac{y - px}{1 + p^2};$$

and these substituted in the equations to a straight line and circle, viz. $X + a = 0$, $Y^2 + (X - c)^2 = a^2$, lead to the preceding equations.

Ex. 3. To find a curve such that, α, β, γ being the angles which the sides of a triangle on a given base and with its vertex a point in the curve, form with the tangent at that point, we may always have $\tan^2 \beta = \tan \alpha \tan \gamma$.

Let the given base $SC = c$, $SN = x$, $NP = y$, the co-ordinates of a point P in the curve, PT a tangent, PG a normal at that point, AP parallel to SC ; $\angle APT = \alpha$, $SPT = \beta$, $CPT = \gamma$ (fig. 8); then $\tan^2 \beta = \tan \alpha \tan \gamma$ gives

$$1 - \frac{\cos^2 \beta}{\cos^2 \alpha} = \tan \alpha \tan (\gamma - \alpha);$$

$$\therefore 1 - \left(\frac{SG}{SP}\right)^2 = \frac{GN}{PN} \cdot \frac{PN}{CN} = \frac{GN}{CN};$$

$$\text{or } 1 - \frac{\left(x + y \frac{dy}{dx}\right)^2}{x^2 + y^2} = -\frac{y}{x - c} \frac{dy}{dx};$$

$$\text{or } \frac{x + y \frac{dy}{dx} - c}{x - c} = \frac{\left(x + y \frac{dy}{dx}\right)^2}{x^2 + y^2};$$

$$\text{or } \frac{\rho \frac{d\rho}{dx} - c}{x - c} = \left(\frac{d\rho}{dx}\right)^2; \text{ or, putting } \frac{d\rho}{dx} = p,$$

$$\rho = (x - c)p + \frac{c}{p};$$

$$\therefore p = p + (x - c) \frac{dp}{dx} - \frac{c}{p^2} \frac{dp}{dx}; \therefore \frac{dp}{dx} = 0, \text{ or } p = C;$$

$$\therefore \sqrt{x^2 + y^2} = (x - c) C + \frac{c}{C}, \text{ the complete integral,}$$

representing a conic Section, the two fixed points being its focus and center, and its eccentricity $= C$; and $(x - 2c)^2 + y^2 = 0$, is the singular solution, representing a point, viz. the other focus.

Ex. 4. $y = xp + \sqrt{b^2 + a^2 p^2},$

$$y = Cx + \sqrt{b^2 + C^2 a^2}, \text{ the complete integral,}$$

$$a^2 y^2 + b^2 x^2 = a^2 b^2, \text{ the singular solution.}$$

This is the solution of the problem to find a curve, such that the product of the perpendiculars dropped from two given

points upon the tangent may be invariable; for taking the line joining the two given points (whose distance suppose $= 2c$) for the axis of x , and their middle point for origin, and x, y the co-ordinates of any point in the curve, the equation to the tangent at that point will be

$$Y - y = \frac{dy}{dx} (X - x), \quad \text{or} \quad Y = \frac{dy}{dx} X + \left(y - x \frac{dy}{dx} \right),$$

and the lengths of the perpendiculars dropped upon this line from the points $(c, 0)$, $(-c, 0)$ will be

$$\frac{-pc - (y - xp)}{\sqrt{1 + p^2}}, \quad \frac{pc - (y - xp)}{\sqrt{1 + p^2}};$$

and the product of these $= \frac{(y - xp)^2 - c^2 p^2}{1 + p^2} = b^2$, suppose;

$$\therefore y = xp + \sqrt{b^2 + a^2 p^2}, \quad \text{putting } a^2 = b^2 + c^2.$$

35. A still more general case is the equation

$$y = xf(p) + \phi(p),$$

which by differentiation is reduced to a linear equation of the first order in x ; for we get

$$p = f(p) + x \frac{d}{dp} f(p) \frac{dp}{dx} + \frac{d}{dp} \phi(p) \frac{dp}{dx};$$

$$\therefore \{p - f(p)\} \frac{dx}{dp} = x \frac{d}{dp} f(p) + \frac{d}{dp} \phi(p);$$

$$\therefore \frac{dx}{dp} + x \frac{\frac{d}{dp} f(p)}{f(p) - p} = - \frac{\frac{d}{dp} \phi(p)}{f(p) - p},$$

which gives $x = F(p)$; then p must be eliminated between this and the proposed equation.

Ex. 1. $y = xp^2 + 2p$; $\therefore p = p^2 + 2xp \frac{dp}{dx} + 2 \frac{dp}{dx},$

$$\text{or } \frac{dx}{dp} + \frac{2x}{p - 1} = - \frac{2}{p^2 - p},$$

which is made integrable by the factor $(p-1)^2$;

$$\therefore x(p-1)^2 = - \int dp \frac{2p-2}{p} = -2p + \log p^2 + C.$$

But $p = -\frac{1}{x} \pm \sqrt{\frac{y}{x} + \frac{1}{x^2}}$; therefore substituting this in the preceding, we obtain the complete primitive between x and y .

Ex. 2. $y = xmp + n\sqrt[3]{1+p^3}$;

$$xp^{\frac{m}{m-1}} = -\frac{n}{m-1} \int dp \frac{p^{\frac{2m-1}{m-1}}}{(1+p^3)^{\frac{1}{3}}}.$$

Ex. 3. $y + p(a-x) = n \int dx \sqrt{1+p^2}$;

$$y = \frac{(a-x)^{n+1}}{2c^n(n+1)} + \frac{c^n}{2(n-1)(a-x)^{n-1}} + C'.$$

This is the solution of the problem of finding the path of a point P which moves uniformly towards another point Q also moving uniformly in a straight line.

For taking A (fig. 3) for the origin, and AB , which is perpendicular to By the line in which Q moves, for the axis of x , we have, supposing P and Q to start together from A and B , $BQ = nAP$, or if $AN = x$, $NP = y$, $AB = a$,

$$y + (a-x)p = n \int dx \sqrt{1+p^2}.$$

36. In the following examples the method of substitution succeeds.

Ex. 1. $(1-p^2)xy = p(x^2 - y^2 - c^2)$,

which expresses that the normal bisects the angle between the focal distances; $2c$ being the distance of the foci, the origin at the middle point between them, and the line joining them the axis of x .

$$\text{Let } p = \frac{xz}{y}, \quad \therefore y^2 = x^2z - c^2 \frac{z}{1+z};$$

therefore, differentiating,

$$\left\{ x^2 - \frac{c^2}{(1+z)^2} \right\} \frac{dz}{dx} = 0;$$

this resolves itself into

$$x = \pm \frac{c}{1+z}, \text{ which gives } y^2 + (x-c)^2 = 0,$$

the singular solution;

and $\frac{dz}{dx} = 0$, or $z = C$, which gives $y^2 = C \left(x^2 - \frac{c^2}{1+C} \right)$, the complete integral.

If we integrate $p = \frac{Cx}{y}$, we get $y^2 = Cx^2 + C'$, where C' must be determined by the condition of the proposed equation being satisfied; and by this condition, in general whenever the method of solution raises the order of the equation, must the number of constants be reduced.

By the same substitution may be solved the more general form

$$axy p^2 + p(x^2 - ay^2 - b) - xy = 0.$$

$$\text{Ex. 2. } x \frac{dy}{dx} - y = X \sqrt{\left(\frac{dy}{dx}\right)^2 - \frac{2y}{x} \frac{dy}{dx} + 1},$$

$$\text{or } \frac{dy}{dx} - \frac{y}{x} = \frac{X}{x} \sqrt{\left(\frac{dy}{dx} - \frac{y}{x}\right)^2 + 1 - \frac{y^2}{x^2}}.$$

$$\text{Let } y = xz, \text{ then } \frac{dy}{dx} = z + x \frac{dz}{dx} = \frac{y}{x} + x \frac{dz}{dx};$$

$$\therefore x^2 \left(\frac{dz}{dx}\right)^2 = \frac{X^2}{x^2} \left\{ x^2 \left(\frac{dz}{dx}\right)^2 + 1 - z^2 \right\};$$

$$\therefore \frac{1}{\sqrt{1-z^2}} \frac{dz}{dx} = \frac{X}{x \sqrt{x^2 - X^2}}.$$

$$\text{If } X = 1, \text{ we have } \sin^{-1} z = \sec^{-1} x + C,$$

$$\text{or } \sin^{-1} \frac{y}{x} = \sec^{-1} x + C.$$

Ex. 3 and 4.

$$\frac{y - xp}{\sqrt{1+p^2}} = f(\sqrt{x^2 + y^2}),$$

$$\frac{y - xp}{\sqrt{x^2 + y^2} \sqrt{1+p^2}} = f\left(\frac{x}{\sqrt{x^2 + y^2}}\right).$$

Introducing polar co-ordinates, we get for the first,

$$\frac{\rho^2}{\sqrt{\left(\frac{d\rho}{d\theta}\right)^2 + \rho^2}} = f(\rho), \quad \text{or} \quad \frac{d\theta}{d\rho} = \frac{f(\rho)}{\rho \sqrt{\rho^2 - \{f(\rho)\}^2}}.$$

The second gives

$$\frac{\rho}{\sqrt{\left(\frac{d\rho}{d\theta}\right)^2 + \rho^2}} = f(\cos \theta), \quad \text{or} \quad \frac{1}{\rho} \frac{d\rho}{d\theta} = \frac{\sqrt{1 - \{f(\cos \theta)\}^2}}{f(\cos \theta)}.$$

The former expresses that the perpendicular on the tangent from the origin is a given function of the radius vector; the latter that the sine of the angle at which the radius vector cuts the curve is a given function of the cosine of the angle at which it is inclined to the axis of x .

SECTION III.

ON THE SINGULAR SOLUTIONS OF DIFFERENTIAL EQUATIONS OF THE FIRST ORDER.

37. FROM the complete integral of a differential equation we can deduce as many particular integrals as we please, by giving to the arbitrary constant particular values. But some differential equations are satisfied by a relation between x and y not containing an arbitrary constant, and not deducible from the complete integral; such a relation is called, as has been said, a singular solution of the differential equation. The existence of such solutions depends upon the fact, that when a solution of a differential equation has been obtained in any manner, it will still be a solution after a quantity of any kind has been introduced in any way, provided the same derived equation result. This is merely an extension of the principle on which the arbitrary constant is added.

38. Before entering upon the general theory, it may be useful to consider the following particular instance.

Let the equation

$$y = x \frac{dy}{dx} + a + a \left(\frac{dy}{dx} \right)^2 \dots\dots\dots(1)$$

be proposed, which, since it falls under Clairaut's form, has for its complete integral

$$y = Cx + a + aC^2; \dots\dots\dots(2)$$

C being the arbitrary constant. If we now regard C , not as a constant, but as a function of x , and differentiate, we get

$$\frac{dy}{dx} = C + (x + 2aC) \frac{dC}{dx};$$

and if we eliminate C between this and $y = Cx + a + aC^2$, we shall obtain a differential equation, but not the proposed one,

for that arises by eliminating C by means of the equation $\frac{dy}{dx} = C$. But if C be so determined as to make the coefficient of $\frac{dC}{dx}$ vanish, that is, if $C = -\frac{x}{2a}$, then the derived equation will be $\frac{dy}{dx} = C$, and the result of the elimination of C will be the proposed equation.

Substituting for C its value, we get

$$y = -\frac{x^2}{2a} + a + \frac{x^2}{4a} = -\frac{x^2}{4a} + a,$$

a relation which manifestly satisfies the proposed equation; for it gives $\frac{dy}{dx} = -\frac{x}{2a}$; and these values of y and $\frac{dy}{dx}$, being substituted in the proposed equation, make it identical. But this solution contains no arbitrary constant, and being the equation to a parabola it cannot, either by making $C=0$, or any other constant quantity, arise from the complete integral which is the equation to a straight line; it is consequently a singular solution, and arises from the complete integral by changing C into a function of x so determined as to make the term involving $\frac{dC}{dx}$ disappear from the value of $\frac{dy}{dx}$.

Thus we see how the singular solution arises from the complete integral; next, let us consider its geometrical signification. The proposed differential equation expresses that the curves to which it belongs have the property, that the tangent at any point is intersected by a perpendicular upon it from a given point, in a given straight line.

Take the given point S (fig. 4) for the origin, and TS , AS respectively parallel and perpendicular to the given line AC , for the axes of x and y . Let TC be the tangent at a point whose co-ordinates are x and y , then its equation is

$$Y - y = \frac{dy}{dx}(X - x),$$

and the equation to the perpendicular upon it from S is

$$Y \frac{dy}{dx} = -X,$$

and for their point of intersection

$$Y \left\{ 1 + \left(\frac{dy}{dx} \right)^2 \right\} - y = -x \frac{dy}{dx},$$

but this point is always in AC for which $Y = AS = a$;

$$\therefore y - x \frac{dy}{dx} = a + a \left(\frac{dy}{dx} \right)^2,$$

the same as the proposed equation.

The complete integral $y = Cx + a(1 + C^2)$, which represents a series of straight lines, evidently satisfies the problem for all values of C ; for let Tb be any one of these lines, then $y = -\frac{x}{C}$ is the equation to a line through S perpendicular to it; and combining the equations to get the co-ordinates of their point of intersection, we have

$$y = -C^2y + a(1 + C^2), \quad \text{or } y = a;$$

the intersection consequently falls in AC . And as a straight line is its own tangent at every point, the equation

$$y = Cx + a(1 + C^2), \text{ for all values of } C,$$

represents a line such that the intersection of the tangent at any point, and a perpendicular upon it from S , falls in the given line AC . Now the curve which is generated by the perpetual intersections of these lines will also satisfy the problem; for each of the lines will be a tangent to it, and therefore perpendiculars from S upon its tangents will intersect them in the various points of AC . To get the equation to this curve we must, according to the usual method, differentiate with respect to the parameter C , which gives

$$0 = x + 2Ca,$$

and eliminate C between this, and the equation

$$y = Cx + a(1 + C^2),$$

which gives

$$y = -\frac{x^2}{4a} + a, \quad \text{or} \quad 4a(a - y) = x^2,$$

the equation to a parabola, vertex A , focus S , of which curve it is a well-known property that the perpendicular from the focus intersects the tangent at any point, in the line touching the parabola at its vertex.

This result being obtained by exactly the same process as the singular solution was obtained, of course coincides with it; hence it appears that the singular solution belongs to the curve which touches the family of curves resulting from the complete integral by making the arbitrary constant assume all possible values. The conclusions arrived at in this particular instance, we shall now shew to hold generally.

39. Having given the complete integral of a differential equation, to find its singular solution.

Let $f\left(x, y, \frac{dy}{dx}\right) = 0$ be a proposed differential equation, and suppose it to result from the elimination of the arbitrary constant c , between the equation $F(x, y, c) = 0$, and its immediately derived equation

$$M + N \frac{dy}{dx} = 0.$$

Now change c into c' any function of x and y , then our equation becomes $F(x, y, c') = 0$; and its immediately derived equation is

$$M' + N' \frac{dy}{dx} + C' \frac{dc'}{dx} = 0 \dots\dots\dots (1),$$

c' entering into M' and N' just as c did into M and N , and $\frac{dc'}{dx}$ being formed in the usual way for a function of two variables x and y , the latter of which is dependent on the former. Now C' is the differential coefficient of $F(x, y, c')$ with respect to c' , regarding x and y as constant, and will therefore usually involve x, y , and c' ; and if put equal to zero, will give such a value for c' as makes the last term of equation (1) disappear;

and then the elimination of c' must evidently produce the proposed equation $f\left(x, y, \frac{dy}{dx}\right) = 0$. Let this value of c' be substituted in $F(x, y, c) = 0$; then this equation is changed into $\phi(x, y) = 0$, and furnishes a relation between x and y which satisfies the equation $f\left(x, y, \frac{dy}{dx}\right) = 0$, but contains no arbitrary constant, and is not deducible from the complete integral by giving a particular value to the constant; since it results from the complete integral by substituting for c a variable value deduced from the equation $\frac{d}{dc}F(x, y, c) = 0$. Consequently, the relation $\phi(x, y) = 0$ is the singular solution required.

40. To explain the geometrical signification of the singular solution of a differential equation.

Let $F(x, y, c) = 0$ (1) be the complete integral of a differential equation between two variables; if we differentiate it with regard to c , we have $\frac{d}{dc}F(x, y, c) = 0$ (2); and if between these equations we eliminate c , we get $\phi(x, y) = 0$ (3) where c does not appear, and which is a singular solution of the differential equation of which (1) is the complete primitive. Suppose equation (1) to be the equation to a system of curves, in which the position and dimensions of any particular curve is defined by a particular value of the parameter c ; also let equation (3) be the equation to a curve referred to the same co-ordinate axes. Now equations (1) and (2), when c receives a certain value, are satisfied by the same values of x and y . Hence from the manner of its formation, equation (3) is satisfied by the same values; or the curves which are represented by (3) and (1), with a particular value of c , have a common point. But equations (3) and (1), being each a solution of the same differential equation, furnish the same value of $\frac{dy}{dx}$ for the same values of x and y ; consequently the curves touch one another at their common point. The same thing happens for every one of the system of curves which equation (1) represents. Therefore the curve represented by the singular solution touches in a point every curve represented

by the complete primitive. This is the geometrical interpretation of the singular solution of a differential equation of the first order.

41. Having given a solution of a differential equation, to find whether it is included in the complete integral or not.

Let $\frac{dy}{dx} = f(x, y)$ be the proposed differential equation, and $y = F(x, c)$ its complete integral, c being the arbitrary constant; and when $c = c'$, let this become $y = u$, u containing no arbitrary constant; then $y = u$ is a particular integral of the proposed.

Hence since $F(x, c) - u$ becomes zero when $c = c'$, we have

$$F(x, c) - u = (c - c')^m \cdot z = az \text{ suppose,}$$

z being a function of x and c , which is neither infinite nor zero when $c = c'$, or when $a = 0$; and m expressing the highest power of $c - c'$ which enters into every term of $F(x, c) - u$. Consequently the complete integral becomes

$$y = u + az,$$

which being substituted in the proposed equation,

$$\frac{dy}{dx} = f(x, y),$$

$$\text{gives } \frac{dy}{dx} + a \frac{dz}{dx} = f(x, u + az) \quad (1).$$

Now since z is neither infinite nor zero when $a = 0$, we may expand it in a series of ascending powers of a , in the form

$$z = K + Aa^\alpha + Ba^\beta + \&c.,$$

α, β , &c. being increasing and positive, and K, A, B , &c. functions of x ;

$$\therefore \frac{du}{dx} + a \frac{dz}{dx} = \frac{du}{dx} + \frac{dK}{dx} a + \frac{dA}{dx} a^{\alpha+1} + \&c.$$

Again, the development of $f(x, u + az)$ will be of the form

$$f(x, u + az) = f(x, u) + M(az)^m + N(az)^n + \&c.,$$

m, n , &c. being increasing and positive. Hence, by substitution in equation (1), observing that $\frac{du}{dx} = f(x, u)$, we get

$$\frac{dK}{dx} \cdot a + \frac{dA}{dx} \cdot a^{a+1} + \&c.$$

$$= Ma^m (K + Aa^a + \&c.)^m + Na^n (K + Aa^a + \&c.)^n + \&c.$$

Now unless this equation is identical, $y = u$ cannot result from $y = F(x, c)$ by changing c into c' ; and therefore $y = u$ cannot be a particular integral of the proposed; and consequently if it satisfies the proposed, it must be a singular solution. Now the indices $m, n, \&c.$, are known, for they result from writing $u + az$ for y in $f(x, y)$ and expanding according to powers of az ; and we must endeavour to determine $\alpha, \beta, \&c.$ so that the two members of the equation may be identical. If m be > 1 , this can easily be effected; for we must have $\frac{dK}{dx} = 0$, or $K = \text{a constant}$; $a + 1 = m$, and $\frac{dA}{dx} = MK^m$; and so on for the other terms. Consequently it will be possible to make the two members identical, and $y = u$ will be a particular integral. In the same way the identity may be established if $m = 1$. But if $m < 1$, there is no term on the first side corresponding to $MK^m a^m$; and since K cannot be equal to zero, it is impossible to satisfy the identity; and therefore $y = u$ is a singular solution. Hence to discover whether a given solution, $y = u$, of a differential equation

$$\frac{dy}{dx} = f(x, y),$$

is a singular solution or not; we must write $u + h$ for y in the value of $\frac{dy}{dx}$, and if the expansion in ascending powers of h involve a power of h , whose index is < 1 , the solution in question is a singular solution; otherwise it is a particular integral.

42. To deduce the singular solutions from the differential equation, without knowing its complete primitive.

Let $y = u$ be a singular solution of the equation

$$\frac{dy}{dx} = f(x, y);$$

then by the preceding article, substituting $u + h$ for y , we get

$$f(x, u + h) = f(x, u) + Mh^m + Nh^n + \&c.$$

where m , n , &c. are proper fractions ;

$$\therefore \frac{d}{du}f(x, u+h) = \frac{d}{dh}f(x, u+h) = mMh^{m-1} + nNh^{n-1} + \&c.;$$

consequently, when $h=0$, $\frac{d}{du}f(x, u) = \infty$.

But $\frac{d}{du}f(x, u)$ is what $\frac{d}{dy}f(x, y)$ becomes when $y=u$; and therefore, conversely, every value u of y , which satisfies $\frac{dy}{dx}=f(x, y)$, and makes $\frac{d}{dy}f(x, y) = \infty$, is a singular solution of the equation

$$\frac{dy}{dx} = f(x, y).$$

43. It is not essential to give the equation the explicit form $\frac{dy}{dx}=f(x, y)$. For let $\frac{dy}{dx}=p$, and let $V=0$ be the given relation between x , y and p ; then we may regard p as a function of x and y determined by the equation $V=0$; hence forming the differential coefficients $\frac{dV}{dy}$, $\frac{dV}{dp}$ as if y and p were independent of one another, we get

$$\frac{dV}{dy} + \frac{dV}{dp} \frac{dp}{dy} = 0, \quad \text{or} \quad \frac{dp}{dy} = -\frac{dV}{dy} \div \frac{dV}{dp}.$$

Hence the condition $\frac{d}{dy}f(x, y) = \infty$ is equivalent to $\frac{dV}{dp} = 0$, provided $\frac{dV}{dy}$ remains finite; and therefore the singular solutions of the equation $V=F(x, y, p)=0$, are determined by eliminating p between $V=0$ and $\frac{dV}{dp}=0$, provided always that these solutions satisfy the proposed equation and do not make $\frac{dV}{dy}=0$. And conversely if the solution of an equation be given, and we deduce from it the value of $\frac{dy}{dx}=p$, then if this value make $\frac{dV}{dp}$ vanish without making at the same time $\frac{dV}{dy}=0$, it is

a singular solution; otherwise it is a particular integral. It is evident that if we consider y as the independent variable, and put the equation under the form

$$V = F\left(x, y, \frac{dx}{dy}\right) = 0,$$

the same reasonings would shew that singular solutions may be obtained by eliminating $p' = \frac{dx}{dy}$, between $V = 0$ and $\frac{dV}{dp'} = 0$, provided that these solutions do not at the same time make $\frac{dV}{dx} = 0$.

Ex. 1. To find the singular solution of

$$y - x \frac{dy}{dx} + x - \frac{y}{\frac{dy}{dx}} = a.$$

Here $\frac{dV}{dp} = -x + \frac{y}{p^2} = 0$; $\therefore p^2 = \frac{y}{x}$, and the proposed becomes

$$(x + y - a)p = y + xp^2, \text{ or } (x + y - a)p = 2y, \\ \text{or } (x + y - a)^2 = 4xy,$$

which may be reduced to the form $\sqrt{x} + \sqrt{y} = \sqrt{a}$.

$$2. \quad \left(y - x \frac{dy}{dx}\right) \left(x - \frac{y}{\frac{dy}{dx}}\right) = a^2. \quad 4xy = a^2.$$

$$3. \quad \left(y - x \frac{dy}{dx}\right)^2 + \left(x - \frac{y}{\frac{dy}{dx}}\right)^2 = a^2. \quad x^{\frac{2}{3}} + y^{\frac{2}{3}} = a^{\frac{2}{3}}.$$

$$4. \quad \left(y - x \frac{dy}{dx}\right) \left\{y - (x - 2a) \frac{dy}{dx}\right\} = b^2, \quad y^2 = \frac{b^2}{a^2} (2ax - x^2).$$

The three former examples determine respectively the curves which have the properties that $OT + OT'$ is constant, that the area of the triangle TOT' is constant, and that TT' is constant, TT' being the tangent at any point meeting the axes Ox , Oy in T and T' , (fig. 5); the last determines a curve such that the

product of the portions of two fixed parallel straight lines intercepted between the tangent at any point and the axis of x , shall be invariable.

$$5. \quad y + \left\{ x - \tan^{-1} \left(\frac{dy}{dx} \right) \right\} \frac{dy}{dx} - 1 = 0.$$

$$\text{Here } \frac{dV}{dp} = x - \tan^{-1} p - \frac{p}{1+p^2} = 0;$$

$$\therefore x = \tan^{-1} p + \frac{p}{1+p^2};$$

$$y = \frac{1}{1+p^2}, \quad \therefore x = \cos^{-1} \sqrt{y} + \sqrt{y-y^2},$$

which represents a cycloid whose base coincides with the axis of x , the origin being in the centre of the base. This is the solution of the problem to find a curve always touched by the same diameter of a circle rolling along a straight line.

$$6. \quad \frac{dy}{dx} = \frac{x}{\sqrt{x^2 + y^2 - a^2} - y};$$

$$\therefore \frac{dp}{dy} = \frac{-x}{(\sqrt{x^2 + y^2 - a^2} - y)^2} \left(\frac{y}{\sqrt{x^2 + y^2 - a^2}} - 1 \right);$$

\therefore the relation $x^2 + y^2 - a^2 = 0$ makes $\frac{dp}{dy}$ infinite, and satisfies the proposed equation; it is consequently a singular solution of the proposed. But if we suppose the solution given, we may find whether it is comprised in the complete integral or not, by Art. 41. For we have $y = \sqrt{a^2 - x^2}$; therefore substituting $\sqrt{a^2 - x^2} + h$ for y in the value of $\frac{dy}{dx}$, we get

$$\frac{x}{\sqrt{2h}\sqrt{a^2 - x^2} + h - \sqrt{a^2 - x^2} - h} = \frac{-x}{\sqrt{a^2 - x^2}} \left\{ 1 + \frac{\sqrt{2h}}{(a^2 - x^2)^{\frac{1}{2}}} - \&c. \right\},$$

when developed according to powers of h ; and as the index of h is a proper fraction, $x^2 + y^2 - a^2 = 0$ cannot be comprised in the complete integral, and is therefore a singular solution.

7. $(x-a) \left(\frac{dy}{dx}\right)^2 - y \frac{dy}{dx} + a = 0$, to find whether $y^2 = 4a(x-a)$, $y = x$, are particular integrals or singular solutions. We get

$$\frac{dV}{dp} = 2(x-a)p - y, \quad \frac{dV}{dy} = -p.$$

Hence the solution $y = x$ which gives $p = 1$, does not make $\frac{dV}{dp}$ vanish, and is therefore a particular integral. In fact the proposed equation being

$$y = (x-a) \frac{dy}{dx} + \frac{a}{\frac{dy}{dx}},$$

its complete integral is (Art. 34)

$$y = (x-a) C + \frac{a}{C};$$

and this becomes $y = x$, when $C = 1$. But the solution

$$y^2 = 4a(x-a) \text{ gives } yp = 2a;$$

and these values of y and p reduce $\frac{dV}{dp}$ to zero without making $\frac{dV}{dy}$ vanish; consequently $y^2 = 4a(x-a)$ is a singular solution.

$$8. \quad y^2 \left(\frac{dy}{dx}\right)^2 + 2xy \frac{dy}{dx} + ax + by = 0.$$

$$\frac{dV}{dp} = 2y(py+x), \quad \frac{dV}{dy} = 2p(py+x).$$

The value of p which makes $\frac{dV}{dp}$ vanish is $p = -\frac{x}{y}$, and the result of the elimination of p is $ax + by - x^2 = 0$; but as $\frac{dV}{dy}$ vanishes for the same value of p , this cannot be a singular solution. In fact, it does not satisfy the proposed equation, and is not a solution of any sort.

$$9. \left(\frac{dy}{dx}\right)^2 - 2ay \frac{dy}{dx} + bx = 0, \quad \frac{dV}{dp} = 2p - 2ay, \quad \frac{dV}{dy} = -2ap;$$

the value of p which makes $\frac{dV}{dp}$ vanish, is $p = ay$, and this value does not make $\frac{dV}{dy}$ vanish; but the result of the elimination of p , $a^2y^2 = bx$, as it does not satisfy the given equation, is not a singular solution.

44. Every factor proper to make a proposed differential equation integrable, is made infinite by the singular solution.

Let $\frac{dy}{dx} + f(x, y) = 0$ be the proposed equation, $F(x, y) = c$ its complete integral, and z the factor which makes it integrable, so that

$$z \left\{ \frac{dy}{dx} + f(x, y) \right\} = \frac{d}{dx} F(x, y);$$

also, let $y = u$ be the singular solution; then this is not deducible from the complete integral, and therefore if u be written for y in $F(x, y)$, the result will not be constant; if therefore we substitute u for y , in the preceding equation, since the second member will have a finite value, and the factor $\frac{dy}{dx} + f(x, y)$ of the first member will be zero, the value of the other factor z corresponding to this substitution must be infinite.

This property will sometimes lead to the discovery of the factor which makes an equation integrable; as in the example

$$(a^2 - x^2) \frac{dy}{dx} + xy = a \sqrt{x^2 + y^2 - a^2},$$

a singular solution of which is $x^2 + y^2 - a^2 = 0$; if we try a factor of the form

$$(x^2 - a^2)^m (y^2 + x^2 - a^2)^n,$$

we arrive at $m = -1$, $n = -\frac{1}{2}$; and the factor which makes the proposed integrable is

$$(x^2 - a^2)^{-1} (x^2 + y^2 - a^2)^{-\frac{1}{2}}.$$

SECTION IV.

DIFFERENTIAL EQUATIONS OF THE SECOND ORDER, AND OF HIGHER ORDERS.

45. EVERY differential equation of the n^{th} order admits of a primitive with n arbitrary constants.

Let $f(x, y, c_1, c_2, \dots c_n) = 0$ be an equation between the variables x and y , containing n constants $c_1, c_2, \dots c_n$. Let the first n derived equations be

$$f_1(x, y, \frac{dy}{dx}, c_1, \dots c_n) = 0,$$

$$f_2(x, y, \frac{dy}{dx}, \frac{d^2y}{dx^2}, c_1, \dots c_n) = 0,$$

.....

$$f_n(x, y, \frac{dy}{dx}, \frac{d^2y}{dx^2}, \dots \frac{d^ny}{dx^n}, c_1, \dots c_n) = 0.$$

Between these n equations and the original, the n constants may be eliminated, and the result will be

$$F\left(x, y, \frac{dy}{dx}, \frac{d^2y}{dx^2}, \dots \frac{d^ny}{dx^n}\right) = 0, \dots\dots\dots (1)$$

a differential equation in which none of the constants enter.

Conversely, a differential equation of the n^{th} order being proposed, it must admit of a primitive containing n arbitrary constants, because that number of constants, and no more, can be eliminated in its formation. Hence every differential equation of the n^{th} order admits of a primitive containing n arbitrary constants.

46. Again, between the original equation and its first $n - 1$ derived equations, $n - 1$ of the constants may be eliminated,

and a differential equation of the $(n-1)^{\text{th}}$ order with one constant will result.

Every such differential equation, having the same primitive with equation (1), is a first integral of that equation; hence a differential equation of the n^{th} order has n first integrals, each a differential equation of the $(n-1)^{\text{th}}$ order, and containing one constant.

Also, between the original equation and the first r of its derived equations, r of the constants may be eliminated, and a differential equation of the r^{th} order containing $n-r$ constants will result, which is an integral of equation (1). Now r constants can be eliminated in a number of ways equal to the number of combinations of n things taken r together, or

$$\frac{n(n-1) \dots (n-r+1)}{1 \cdot 2 \cdot 3 \dots r}.$$

This then is the number of integrals which equation (1) has of the $(n-r)^{\text{th}}$ order, each a differential equation of the r^{th} order and containing $n-r$ constants.

47. Of the general equation of the second order

$$F\left(\frac{d^2y}{dx^2}, \frac{dy}{dx}, y, x\right) = 0,$$

we shall first of all consider the following particular cases, in which $\frac{d^2y}{dx^2}$ is involved with only one, or two, of the other quantities $x, y, \frac{dy}{dx}$; and which admit of integration, or rather of reduction to forms of the first order.

I. $F\left(\frac{d^2y}{dx^2}, x\right) = 0$. Let this by resolution give

$$\frac{d^2y}{dx^2} = f(x), \quad \therefore \frac{dy}{dx} = \int dx f(x) + C;$$

and integrating again and adding another constant, we obtain the complete integral. The same process applies to $\frac{d^2y}{dx^2} = f(x)$.

Also, if we have $F\left(\frac{d^{n+1}y}{dx^{n+1}}, \frac{d^ny}{dx^n}\right) = 0$, and put $\frac{d^ny}{dx^n} = u$, we get $F\left(\frac{du}{dx}, u\right) = 0$; and if this can be integrated, and gives $u = f(x)$, it is reduced to the case just noticed.

$$\text{Ex. 1. } x^2 \frac{d^2y}{dx^2} = a, \quad \therefore \frac{d^2y}{dx^2} = \frac{a}{x^2}, \quad \frac{dy}{dx} = -\frac{a}{x} + C;$$

$$\therefore y = a \log \frac{1}{x} + Cx + C'.$$

$$\text{Ex. 2. } \frac{d^2y}{dx^2} \cdot \frac{d^3y}{dx^3} + x = 0, \quad \therefore \frac{d^2y}{dx^2} = \sqrt{c^2 - x^2};$$

$$\therefore y = \frac{1}{6} (2c^2 + x^2) \sqrt{c^2 - x^2} + \frac{1}{2} c^2 x \sin^{-1} \frac{x}{c} + Cx + C'.$$

$$\text{II. } F\left(\frac{d^2y}{dx^2}, \frac{dy}{dx}\right) = 0. \text{ Let this by resolution give}$$

$$\frac{d^2y}{dx^2} = f\left(\frac{dy}{dx}\right), \text{ or } \frac{dp}{dx} = f(p), \text{ putting } \frac{dy}{dx} = p;$$

$$\therefore \frac{dx}{dp} = \frac{1}{f(p)}, \quad x = \int dp \frac{1}{f(p)};$$

$$\text{and } \frac{dy}{dp} = \frac{dy}{dx} \cdot \frac{dx}{dp} = \frac{p}{f(p)}, \quad \therefore y = \int dp \frac{p}{f(p)};$$

p must be eliminated between these two equations.

$$\text{Ex. } a \frac{d^2y}{dx^2} = \frac{dy}{dx}; \quad \therefore \frac{dp}{dx} = \frac{p}{a}; \quad \therefore \log p = \frac{x}{a} + C,$$

$$\text{and } \frac{dy}{dp} = \frac{dy}{dx} \cdot \frac{dx}{dp} = p \cdot \frac{a}{p} = a; \quad \therefore y = ap + C';$$

$$\therefore \frac{x}{a} + C = \log \frac{y - C'}{a}.$$

$$\text{III. } F\left(\frac{d^2y}{dx^2}, y\right) = 0. \text{ Let this give } \frac{d^2y}{dx^2} = f(y);$$

$$\therefore 2 \frac{dy}{dx} \frac{d^2y}{dx^2} = 2f(y) \frac{dy}{dx}, \text{ and } \left(\frac{dy}{dx}\right)^2 = C + 2 \int dy f(y);$$

$$\therefore \frac{dy}{dx} = \sqrt{C + 2 \int dy f(y)}, \text{ and } x = \int dy \frac{1}{\sqrt{C + 2 \int dy f(y)}}.$$

Also $F\left(\frac{d^{n+2}y}{dx^{n+2}}, \frac{d^n y}{dx^n}\right) = 0$, putting $\frac{d^n y}{dx^n} = u$, becomes

$$F\left(\frac{d^2u}{dx^2}, u\right) = 0;$$

and this, treated as above, gives $x = \phi(u)$; and if $x = \phi(u)$ can be solved with respect to u , it is brought under Case I.

Ex. 1. $a^2 \frac{d^2y}{dx^2} + y = 0; \therefore 2a^2 \frac{dy}{dx} \frac{d^2y}{dx^2} + 2y \frac{dy}{dx} = 0;$

$$\therefore a^2 \left(\frac{dy}{dx}\right)^2 + y^2 = C; \therefore \frac{a}{\sqrt{C - y^2}} \cdot \frac{dy}{dx} = 1,$$

$$\therefore a \sin^{-1} \frac{y}{\sqrt{C}} = x + C'.$$

Ex. 2. $ny^3 \frac{d^2y}{dx^2} - 1 = 0; \frac{1}{n} (x - c')^2 = \frac{1}{c} y^2 - \frac{1}{c^2}.$

This is the solution of the problem, in which a curve concave towards the axis of x is sought whose radius of curvature shall vary as the cube of its normal: for this requires that

$$(1 + p^2)^{\frac{3}{2}} \div \frac{dp}{dx} = ny^3 (1 + p^2)^{\frac{3}{2}}.$$

Ex. 3. $\sqrt{ay} \frac{d^2y}{dx^2} = 1.$

$$\frac{x}{\sqrt{a}} = \frac{2}{3} (\sqrt{y} + C)^3 - 2C \sqrt{y} + C'.$$

IV. $F\left(\frac{d^2y}{dx^2}, \frac{dy}{dx}, x\right) = 0.$

This becomes of the first order in p and x , by putting p for $\frac{dy}{dx}$ and $\frac{dp}{dx}$ for $\frac{d^2y}{dx^2}$; let its integral be $\phi(x, p, C) = 0.$

If this by resolution give p or $\frac{dy}{dx} = f(x)$, then $y = \int dx f(x)$; among other cases, this will happen when the proposed is of the form $\frac{d^2y}{dx^2} + P \frac{dy}{dx} = Q$; for the latter can be solved as a linear equation of the first order.

If it gives $x = f(p)$, then

$$y = \int dx p = xp - \int dx x \frac{dp}{dx} = xp - \int p f(p) dp;$$

and p must be eliminated between these equations.

$$\text{Ex. 1. } \frac{d^2y}{dx^2} + \frac{1}{x} \frac{dy}{dx} = 0; \quad \therefore x \frac{d^2y}{dx^2} + \frac{dy}{dx} = \frac{d}{dx} \left(x \frac{dy}{dx} \right) = 0;$$

$$\therefore x \frac{dy}{dx} = C; \quad \therefore y = C \log x + C'.$$

$$\text{Ex. 2. } (1+x^2) \frac{d^2y}{dx^2} + 1 + \left(\frac{dy}{dx} \right)^2 = 0;$$

$$\therefore \frac{1}{1+p^2} \frac{dp}{dx} + \frac{1}{1+x^2} = 0, \quad \tan^{-1} p + \tan^{-1} x = \tan^{-1} c;$$

$$\therefore p = \frac{c-x}{1+cx} = \frac{1}{c} \left(\frac{1+c^2}{1+cx} - 1 \right);$$

$$\therefore y = \frac{1+c^2}{c^2} \log(1+cx) - \frac{x}{c} + C'.$$

$$\text{Ex. 3. } x(a+bx) \frac{d^2y}{dx^2} + (c+ex) \frac{dy}{dx} = 0;$$

$$\therefore \frac{1}{p} \frac{dp}{dx} = - \frac{c+ex}{x(a+bx)} = - \frac{cx^{-2}}{ax^{-1}+b} - \frac{e}{a+bx};$$

$$\therefore p = c' x^{-\frac{c}{a}} (a+bx)^{\frac{c}{a}-\frac{e}{b}} \text{ and } y = C + c' \int dx x^{-\frac{c}{a}} (a+bx)^{\frac{c}{a}-\frac{e}{b}}.$$

$$\text{Ex. 4. } \frac{\left\{ 1 + \left(\frac{dy}{dx} \right)^2 \right\}^{\frac{3}{2}}}{\frac{d^2y}{dx^2}} = f(x); \quad \therefore \frac{1}{(1+p^2)^{\frac{3}{2}}} \frac{dp}{dx} = \frac{1}{f(x)};$$

$$\therefore \frac{p}{\sqrt{1+p^2}} = \int dx \frac{1}{f(x)} = X, \text{ suppose ;}$$

$$\therefore p = \frac{X}{\sqrt{1-X^2}}, \text{ and } y = \int dx \frac{X}{\sqrt{1-X^2}}.$$

This is the solution of the inverse problem of the radius of curvature, in which it is required to find a curve whose radius of curvature shall be a given function of the abscissa.

$$\text{Ex. 5. } \frac{d^2y}{dx^2} + (e^x - 1) \frac{dy}{dx} = e^{2x}, \quad y = e^x + Ce^{-x} + C'.$$

$$\text{V. } F\left(\frac{d^2y}{dx^2}, \frac{dy}{dx}, y\right) = 0.$$

Putting $\frac{dy}{dx} = p$, we get

$$\frac{d^2y}{dx^2} = \frac{dp}{dy} \cdot \frac{dy}{dx} = p \frac{dp}{dy};$$

and the substitution of $p \frac{dp}{dy}$ for $\frac{d^2y}{dx^2}$, and of p for $\frac{dy}{dx}$, will make the proposed of the first order in p and y ; let its integral be

$$\phi(p, y, C) = 0.$$

If this by resolution give $p = \frac{dy}{dx} = f(y)$, then

$$x = \int dy \frac{1}{f(y)}.$$

If it gives $y = f(p)$, then

$$x = \int dy \frac{1}{p} = \frac{y}{p} + \int dy \frac{y}{p^2} \frac{dp}{dy} = \frac{y}{p} + \int dp \frac{f(p)}{p^2},$$

and p must be eliminated between these equations.

$$\text{Ex. 1. } y \frac{d^2y}{dx^2} + \left(\frac{dy}{dx}\right)^2 = 0, \text{ or } \frac{d}{dx} \left(y \frac{dy}{dx}\right) = 0;$$

$$\therefore y \frac{dy}{dx} = C; \quad \therefore y^2 = 2Cx + C'.$$

$$\text{Ex. 2. } y \frac{d^2y}{dx^2} + n \left(\frac{dy}{dx} \right)^2 + n = 0;$$

$$\therefore yp \frac{dp}{dy} + np^2 + n = 0;$$

$$\therefore \frac{1}{2} \log (1 + p^2) + n \log y = \frac{1}{2} \log C;$$

$$\therefore (1 + p^2) y^{2n} = C;$$

$$\therefore p = \sqrt{Cy^{-2n} - 1}, \text{ and } x = \int dy \frac{y^n}{\sqrt{C - y^{2n}}}.$$

This is the solution of the problem in which it is required to find a curve whose radius of curvature shall vary as its normal; for this condition gives

$$y \sqrt{1 + p^2} = \frac{n(1 + p^2)^{\frac{1}{2}}}{\pm \frac{dp}{dx}}, \text{ or } y \frac{dp}{dx} + (\mp n)(1 + p^2) = 0,$$

— or + according as the curve is convex or concave to the axis of x . If $n = 1$ the curve is a circle, if $n = 2$ a cycloid, if $n = -1$ a common catenary.

$$\text{Ex. 3. } 1 + \left(\frac{dy}{dx} \right)^2 - 2y \frac{d^2y}{dx^2} = 3ay \left\{ 1 + \left(\frac{dy}{dx} \right)^2 \right\}^{\frac{1}{2}}.$$

$$\text{This gives } \frac{dx}{dy} = \frac{ay^{\frac{1}{2}} + C}{\sqrt{y - (ay^{\frac{1}{2}} + C)^2}}.$$

$$\text{Ex. 4. } y^2 \frac{d^2y}{dx^2} + \sqrt{1 + \left(\frac{dy}{dx} \right)^2} = 0, \text{ gives}$$

$$\frac{dx}{dy} = \frac{y}{\sqrt{(Cy + 1)^2 - y^2}}.$$

$$\text{VI. } F\left(\frac{d^2y}{dx^2}, y, x\right) = 0.$$

As there is no substitution by which this can be generally reduced to an equation of the first order between two variables, the artifice to be employed in any case will depend upon the

nature of the example proposed. Among other substitutions for $\frac{d^2y}{dx^2}$, the two following may be noticed,

$$x \frac{d^2y}{dx^2} = \frac{d}{dx} \left\{ x^2 \frac{d}{dx} \left(\frac{y}{x} \right) \right\}, \text{ and } x \frac{d^2y}{dx^2} = \frac{d^2(xy)}{dx^2} - 2 \frac{dy}{dx}.$$

Ex. 1. $x^2 \frac{d^2y}{dx^2} = 2y;$

$$\therefore \frac{d^2(xy)}{dx^2} - 2 \frac{dy}{dx} = \frac{2y}{x}, \text{ or } \frac{d^2(xy)}{dx^2} = \frac{2}{x} \frac{d(xy)}{dx};$$

$$\therefore \frac{d}{dx} (xy) = Cx^2; \text{ and } y = \frac{Cx^2}{3} + \frac{C'}{x}.$$

Ex. 2. $(x^2 + y^2)^2 \frac{d^2y}{dx^2} + a^2 y = 0,$ let $\frac{y}{x} = z$, then

$$\frac{d}{dx} \left(x^2 \frac{dz}{dx} \right) + \frac{a^2}{x^2} \frac{z}{(1+z^2)^2} = 0;$$

$$\therefore x^2 \frac{dz}{dx} \cdot \frac{d}{dx} \left(x^2 \frac{dz}{dx} \right) + \frac{a^2 z}{(1+z^2)^2} \frac{dz}{dx} = 0;$$

$$\therefore \left(x^2 \frac{dz}{dx} \right)^2 - \frac{a^2}{1+z^2} = C,$$

$$\text{or } x^2 \frac{dz}{dx} = \sqrt{C + \frac{a^2}{1+z^2}},$$

$$\text{or } -\frac{1}{x} = \int dz \frac{\sqrt{1+z^2}}{\sqrt{a^2 + C(1+z^2)}}.$$

Ex. 3. $\frac{d^2y}{dx^2} = ax + by;$

$$\text{or } \frac{d^2}{dx^2} (ax + by) = b(ax + by),$$

which becomes $\frac{d^2z}{dx^2} = bz$, putting $ax + by = z$, and so falls under Case III.

48. Of the general equation of the second order involving $x, y, \frac{dy}{dx}, \frac{d^2y}{dx^2}$, the following particular cases may be noticed, in which P denotes a function of x .

$$1. \quad P \frac{d^2y}{dx^2} + \frac{1}{2} \frac{dP}{dx} \cdot \frac{dy}{dx} + cy = 0;$$

this when multiplied by $2 \frac{dy}{dx}$ may be written

$$\frac{d}{dx} \left\{ P \left(\frac{dy}{dx} \right)^2 \right\} + c \frac{d}{dx} y^2 = 0;$$

$$\therefore P \left(\frac{dy}{dx} \right)^2 + cy^2 = C, \quad \text{or} \quad \frac{1}{\sqrt{C - cy^2}} \frac{dy}{dx} = \frac{1}{\sqrt{P}}.$$

$$\text{Thus if } (1 - x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} + q^2 y = 0;$$

$$\therefore \frac{1}{\sqrt{C - q^2 y^2}} \frac{dy}{dx} = \frac{1}{\sqrt{1 - x^2}}, \quad \text{or} \quad \frac{1}{q} \sin^{-1} \frac{qy}{\sqrt{C}} = \sin^{-1} x + C',$$

which may be put under the form

$$y = C_1 \cos (q \sin^{-1} x) + C_2 \sin (q \sin^{-1} x).$$

$$\text{Similarly, } (ax + bx^2) \frac{d^2y}{dx^2} + \left(\frac{1}{2} a + bx \right) \frac{dy}{dx} + cy = 0,$$

$$\text{leads to } \frac{1}{\sqrt{C - cy^2}} \frac{dy}{dx} = \frac{1}{\sqrt{ax + bx^2}}.$$

$$2. \quad \frac{d^2y}{dx^2} + P \left(\frac{dy}{dx} - \frac{y}{x} \right) = 0;$$

$$\therefore \frac{d}{dx} \left(x \frac{dy}{dx} - y \right) + P \left(x \frac{dy}{dx} - y \right) = 0;$$

$$\therefore x \frac{dy}{dx} - y = C e^{-\int dx P} \quad \text{or} \quad \frac{d}{dx} \left(\frac{y}{x} \right) = \frac{C}{x^2} e^{-\int dx P}.$$

$$\therefore y = C x \int dx \cdot \frac{1}{x^2} e^{-\int dx P}.$$

48*. When the equation

$$F\left(x, y, \frac{dy}{dx}, \frac{d^2y}{dx^2}\right) = 0, \quad \text{or} \quad F\left(x, y, p, \frac{dp}{dx}\right) = 0,$$

is homogeneous, reckoning the dimensions of p and $\frac{dp}{dx}$ to be 0 and -1 respectively; it may be reduced to an equation of the first order by putting

$$y = xz, \quad \text{and} \quad \frac{dp}{dx} = \frac{q}{x}.$$

For each term, if r denote its dimensions, will consist of some function of p multiplied by a factor of the form

$$\left(\frac{y}{x}\right)^m x^{n+r} \left(\frac{dp}{dx}\right)^n,$$

m and n being any numbers from 0 to ∞ ; therefore upon making the substitutions stated above, every term will be divisible by x^r , and the equation will assume the form

$$F(p, q, z) = 0;$$

and if this can be solved relative to q , we shall have

$$q = \phi(z, p); \quad \text{but} \quad q = x \frac{dp}{dx} = \frac{(p-z)}{\frac{dz}{dx}} \frac{dp}{dx} = (p-z) \frac{dp}{dz};$$

$$\therefore \phi(z, p) + (z-p) \frac{dp}{dz} = 0;$$

let this give

$$p = \psi(z), \quad \text{then} \quad \psi(z) = z + x \frac{dz}{dx},$$

which will give the required integral $z = \frac{y}{x} = f(x)$.

$$\text{Ex.} \quad n \left\{ 1 + \left(\frac{dy}{dx} \right)^2 \right\}^{\frac{n}{2}} = \frac{d^2y}{dx^2} \sqrt{x^2 + y^2};$$

putting $\frac{dy}{dx} = p$, $y = xz$, $\frac{dp}{dx} = \frac{q}{x}$, we find

$$n(1+p^2)^{\frac{n}{2}} = q \sqrt{1+z^2};$$

$$\therefore q = \frac{n(1+p^2)^{\frac{1}{2}}}{\sqrt{1+z^2}} = (p-z) \frac{dp}{dz},$$

which is the same as Ex. 5, Art. 18.

Linear Equations of the Second and Higher Orders.

49. The linear equation of the n^{th} order is

$$\frac{d^n y}{dx^n} + p_1 \frac{d^{n-1} y}{dx^{n-1}} + p_2 \frac{d^{n-2} y}{dx^{n-2}} + \dots + p_{n-1} \frac{dy}{dx} + p_n y = X,$$

all the coefficients being functions of x , and each term of the first member involving either y , or one of its differential coefficients, in the first power. The first step towards the integration of this equation is the establishment of the following theorem.

If there be n particular values $u_1, u_2, u_3 \dots u_n$, functions of x , which, when substituted for y , satisfy the equation

$$\frac{d^n y}{dx^n} + p_1 \frac{d^{n-1} y}{dx^{n-1}} + p_2 \frac{d^{n-2} y}{dx^{n-2}} + \dots + p_n y = 0,$$

its complete integral is

$$y = a_1 u_1 + a_2 u_2 + a_3 u_3 + \dots + a_n u_n,$$

$a_1, a_2, \dots a_n$ being arbitrary constants.

For let this value of y be substituted in the expression

$$\frac{d^n y}{dx^n} + p_1 \frac{d^{n-1} y}{dx^{n-1}} + \dots + p_n y,$$

and it becomes

$$\begin{aligned} & \frac{d^n}{dx^n} (a_1 u_1 + a_2 u_2 + \dots + a_n u_n) \\ & + p_1 \frac{d^{n-1}}{dx^{n-1}} (a_1 u_1 + a_2 u_2 + \dots + a_n u_n) + \dots + p_n (a_1 u_1 + a_2 u_2 + \dots + a_n u_n), \end{aligned}$$

or, collecting the terms multiplied by the factors $a_1, a_2, \dots a_n$,

$$\begin{aligned} & a_1 \left(\frac{d^n u_1}{dx^n} + p_1 \frac{d^{n-1} u_1}{dx^{n-1}} + \dots + p_n u_1 \right) \\ & + a_2 \left(\frac{d^n u_2}{dx^n} + p_1 \frac{d^{n-1} u_2}{dx^{n-1}} + \dots + p_n u_2 \right) + \dots \\ & + a_n \left(\frac{d^n u_n}{dx^n} + p_1 \frac{d^{n-1} u_n}{dx^{n-1}} + \dots + p_n u_n \right); \end{aligned}$$

Now, since $u_1, u_2, \dots u_n$ satisfy the equation, each of the quantities within brackets is equal to zero, and therefore the whole is identically zero; and therefore the assumed value of y satisfies the equation; and it contains n arbitrary constants, consequently it is the complete integral of the equation.

Thus the equation $\frac{d^2 y}{dx^2} - n^2 y = 0$ is satisfied by $y = e^{nx}$, and since it is not altered by changing the sign of n , it is also satisfied by $y = e^{-nx}$; $\therefore y = a_1 e^{nx} + a_2 e^{-nx}$ is the complete integral.

Linear Equations with constant Coefficients.

50. To integrate the equation

$$\frac{d^n y}{dx^n} + p_1 \frac{d^{n-1} y}{dx^{n-1}} + p_2 \frac{d^{n-2} y}{dx^{n-2}} + \dots + p_n y = q,$$

all the coefficients and q being constant.

First write $y + \frac{q}{p_n}$ instead of y ; and the equation becomes

$$\frac{d^n y}{dx^n} + p_1 \frac{d^{n-1} y}{dx^{n-1}} + \dots + p_n y = 0,$$

which shews that we may treat the proposed equation as if it had no term q independent of y , provided we divide that term by the coefficient of y and add the quotient to the value of y obtained on the supposition that $q = 0$.

Let $y = e^{mx}$, $\therefore e^{mx} (m^n + p_1 m^{n-1} + \dots + p_n)$ is the value of the first member; now this will vanish if m be any root of the equation

$$m^n + p_1 m^{n-1} + p_2 m^{n-2} + \dots + p_n = 0, \text{ or } M = f(m) = 0,$$

which is called the auxiliary equation of the proposed linear equation.

Hence the n real or imaginary roots of this equation, $m_1, m_2, m_3 \dots m_n$, provided they be all unequal, will give n different particular values of y , $e^{m_1 x}, e^{m_2 x}, \dots e^{m_n x}$, which satisfy the proposed equation; and therefore its complete integral is

$$y = a_1 e^{m_1 x} + a_2 e^{m_2 x} + \dots + a_n e^{m_n x}.$$

51. But if any of the roots are equal to one another, as, for instance, $m_1 = m_2$, the value of y becomes

$$y = (a_1 + a_2) e^{m_1 x} + a_3 e^{m_3 x} + \dots a_n e^{m_n x},$$

which contains only $n - 1$ arbitrary constants (because $a_1 + a_2$ can be reckoned only as a single constant), and therefore cannot be the complete integral of the proposed equation. In this case, in order to discover the complete integral, first suppose the two roots m_1, m_2 , to be only very nearly equal to one another, so that $m_2 = m_1 + h$, where h is a very small known quantity; then the part of the value of y corresponding to these roots is

$$\begin{aligned} a_1 e^{m_1 x} + a_2 e^{m_2 x} &= e^{m_1 x} (a_1 + a_2 e^{hx}) = e^{m_1 x} (a_1 + a_2 + a_2 \frac{hx}{1} + a_2 \frac{h^2 x^2}{1 \cdot 2} + \&c.) \\ &= e^{m_1 x} (c_1 + c_2 x + \frac{1}{2} c_2 h x^2 + \frac{1}{6} c_2 h^2 x^3 + \&c.), \end{aligned}$$

replacing the constants $a_1 + a_2$ and $a_2 h$, by c_1, c_2 respectively. Now let $h = 0$, then this becomes $e^{m_1 x} (c_1 + c_2 x)$; and the complete integral consequently is

$$y = (c_1 + c_2 x) e^{m_1 x} + a_3 e^{m_3 x} + \dots + a_n e^{m_n x}.$$

52. Generally, if we suppose r roots of the auxiliary equation to be nearly equal to one another, and therefore to be represented by

$$m_1 + h_1, \quad m_1 + h_2, \quad \dots \quad m_1 + h_r,$$

where $h_1, h_2, \dots h_r$ are very small quantities, the complete integral takes the form

$$y = e^{m_1 x} (a_1 e^{h_1 x} + a_2 e^{h_2 x} + \dots + a_r e^{h_r x}) + a_{r+1} e^{m_{r+1} x} + \&c.,$$

or, expanding $e^{h_1 x}, e^{h_2 x}, \&c.$,

$$y = e^{m_1 x} \left\{ \Sigma(a) + \dots + \frac{\Sigma(a h^{r-1})}{[r-1]} x^{r-1} + \frac{\Sigma(a h^r)}{[r]} x^r + \dots \right\} + a_{r+1} e^{m_{r+1} x} + \&c.,$$

or, replacing the constants

$$\Sigma(a), \quad \Sigma(a h), \quad \dots \quad \frac{\Sigma(a h^{r-1})}{[r-1]},$$

by $c_1, c_2 \dots c_r$,

$$y = e^{m_1 x} (c_1 + c_2 x + \dots + c_r x^{r-1} + \text{terms multiplied by } h_1, h_2, \&c.) \\ + a_{r+1} e^{m_{r+1} x} + \&c.$$

Now let $h_1 = 0 = h_2 = \dots = h_r$, in which case the auxiliary equation has r roots each $= m_1$; then the solution becomes

$$y = e^{m_1 x} (c_1 + c_2 x + c_3 x^2 + \dots + c_r x^{r-1}) + a_{r+1} e^{m_{r+1} x} + \dots + a_n e^{m_n x},$$

which contains n arbitrary constants, and is consequently the complete integral.

53. Of the correctness of the above modification for the case of equal roots, we may assure ourselves by the following reverse process. Let $y = e^{mx} u$, then since by Leibnitz's theorem

$$\frac{d^n (uv)}{dx^n} = \frac{d^n v}{dx^n} u + n \frac{d^{n-1} v}{dx^{n-1}} \frac{du}{dx} + \frac{n(n-1)}{1 \cdot 2} \frac{d^{n-2} v}{dx^{n-2}} \frac{d^2 u}{dx^2} + \&c., \\ \frac{d^n}{dx^n} (e^{mx} u) = e^{mx} \left\{ m^n \cdot u + n m^{n-1} \cdot \frac{du}{dx} + \frac{n(n-1)}{1 \cdot 2} m^{n-2} \frac{d^2 u}{dx^2} + \&c. \right\} \\ = e^{mx} \left(m + \frac{d}{dx} \right)^n u,$$

separating the symbol of operation from that of quantity.

Hence the first member of the equation, or $f\left(\frac{d}{dx}\right) e^{mx} u$, becomes

$$e^{mx} \left(m + \frac{d}{dx} \right)^n u + p_1 e^{mx} \left(m + \frac{d}{dx} \right)^{n-1} u \\ + p_2 e^{mx} \left(m + \frac{d}{dx} \right)^{n-2} u + \&c. + p_n e^{mx} u \\ = e^{mx} f\left(m + \frac{d}{dx}\right) u \\ = e^{mx} \left\{ f(m) u + \frac{1}{1} f'(m) \frac{du}{dx} + \frac{1}{1 \cdot 2} f''(m) \frac{d^2 u}{dx^2} + \dots + \frac{1}{[n]} f^{(n)}(m) \frac{d^n u}{dx^n} \right\}.$$

Now let $m = m_1$, then since $f(m) = 0$ has r roots equal to m_1 ,

each of the quantities $f(m)$, $f'(m)$, ... $f^{(r-1)}(m)$ becomes equal to zero, and the first r terms vanish; and if

$$u = a_1 + a_2x + a_3x^2 + \dots + a_rx^{r-1},$$

then $\frac{d^r u}{dx^r}$, $\frac{d^{r+1} u}{dx^{r+1}}$, ... $\frac{d^n u}{dx^n}$ likewise become zero, and all the remaining terms vanish. Hence for each group of r equal roots whether real or imaginary in the auxiliary equation, there will be in the value of y a term of the form

$$e^{m_1 x} (a_1 + a_2 x + a_3 x^2 + \dots + a_r x^{r-1}).$$

In the treatment of the linear equation with constant coefficients

$$\frac{d^n y}{dx^n} + p_1 \frac{d^{n-1} y}{dx^{n-1}} + \&c. + p_n y = \left\{ \left(\frac{d}{dx} \right)^n + p_1 \left(\frac{d}{dx} \right)^{n-1} + \dots + p_n \right\} y = X,$$

$$\text{or } f\left(\frac{d}{dx}\right) y = X,$$

great use will be made of the formulæ just obtained,

$$\left(\frac{d}{dx} + m \right)^n u = e^{-mx} \frac{d^n}{dx^n} (e^{mx} u), \quad f\left(\frac{d}{dx} + m \right) u = e^{-mx} f\left(\frac{d}{dx} \right) e^{mx} u.$$

When n is negative, since $\left(\frac{d}{dx} \right)^{-1}$ is equivalent to $\int dx$, the former leads to

$$\left(\frac{d}{dx} + m \right)^{-n} u = e^{-mx} \int^n dx^n (e^{mx} u).$$

54. Let $h \pm k\sqrt{-1}$ be a pair of imaginary roots in the auxiliary equation, then the corresponding terms in the value of y will be

$$C e^{hx+kx\sqrt{-1}} + C' e^{hx-kx\sqrt{-1}},$$

$$\text{or } e^{hx} \{ C (\cos kx + \sqrt{-1} \sin kx) + C' (\cos kx - \sqrt{-1} \sin kx) \}$$

$$= e^{hx} \{ c_1 \cos kx + c_2 \sin kx \} \text{ changing the arbitrary constants,}$$

$$\text{or } = \sqrt{c_1^2 + c_2^2} e^{hx} \left\{ \frac{c_1}{\sqrt{c_1^2 + c_2^2}} \cos kx + \frac{c_2}{\sqrt{c_1^2 + c_2^2}} \sin kx \right\}$$

$= \beta e^{hx} \cos(kx + \alpha)$, again changing the constants by putting

$$\sqrt{c_1^2 + c_2^2} = \beta, \text{ and } -\tan \alpha = \frac{c_2}{c_1}.$$

And if there should be r pairs of imaginary roots in the auxiliary equation, equal to $h \pm k\sqrt{-1}$, the corresponding part of the value of y will be

$$\begin{aligned} & e^{hx} (a_1 + a_2 x + \dots + a_r x^{r-1}) (\cos kx + \sqrt{-1} \sin kx) \\ & + e^{hx} (b_1 + b_2 x + \dots + b_r x^{r-1}) (\cos kx - \sqrt{-1} \sin kx), \end{aligned}$$

or, changing the arbitrary constants,

$$\begin{aligned} & e^{hx} (a_1 + a_2 x + \dots + a_r x^{r-1}) \cos kx. \\ & + e^{hx} (\beta_1 + \beta_2 x + \dots + \beta_r x^{r-1}) \sin kx. \end{aligned}$$

55. We shall now give some examples of integrating linear equations with constant coefficients, by the elementary method just investigated.

Ex. 1. $\frac{d^2 y}{dx^2} + \frac{dy}{dx} - 2y = 0.$

Let $y = e^{mx}$, $\therefore m^2 e^{mx} + m e^{mx} - 2e^{mx} = 0,$

or $m^2 + m - 2 = 0$; $\therefore m = 1, \text{ or } -2$;

$$\therefore y = C_1 e^x + C_2 e^{-2x}.$$

2. $a^2 \frac{d^4 y}{dx^4} = \frac{d^2 y}{dx^2}, \quad y = C_1 e^{\frac{x}{a}} + C_2 e^{-\frac{x}{a}} + C_3 x + C_4.$

3. $\frac{d^2 y}{dx^2} + n^2 y = 0.$

Let $y = e^{nx}$; $\therefore m^2 + n^2 = 0$, or $m = \pm n\sqrt{-1}$;

$$\therefore y = C_1 e^{nx\sqrt{-1}} + C_2 e^{-nx\sqrt{-1}}$$

$$= C_1 (\cos nx + \sqrt{-1} \sin nx) + C_2 (\cos nx - \sqrt{-1} \sin nx)$$

$$= (C_1 + C_2) \cos nx + (C_1 - C_2) \sqrt{-1} \sin nx$$

$$= a_1 \cos nx + a_2 \sin nx, \text{ changing the constants,}$$

$$= \beta (\cos nx \cos \alpha - \sin nx \sin \alpha) = \beta \cos (nx + \alpha).$$

This equation, $\frac{d^2 y}{dx^2} + n^2 y = 0$, of which the solution is

$$y = a_1 \cos nx + a_2 \sin nx, \quad \text{or } y = \beta \cos (nx + \alpha),$$

is of very frequent occurrence. Hence also the solution of

$$4. \quad \frac{d^2 y}{dx^2} + n^2 y + ax + b = 0, \quad \text{or}$$

$$\frac{d^2}{dx^2} \left(y + \frac{ax}{n^2} + \frac{b}{n^2} \right) + n^2 \left(y + \frac{ax}{n^2} + \frac{b}{n^2} \right) = 0, \quad \text{is}$$

$$y + \frac{ax}{n^2} + \frac{b}{n^2} = \beta \cos (nx + \alpha).$$

$$5. \quad \frac{d^2 y}{dx^2} - 2m \frac{dy}{dx} + (m^2 + n^2) y = 0;$$

$$y = e^{kx}, \quad \therefore k^2 - 2km + m^2 + n^2 = 0, \quad k = m \pm n \sqrt{-1};$$

$$\therefore y = e^{mx} (C \cos nx + C' \sin nx).$$

$$6. \quad \frac{d^2 y}{dx^2} + 2m \frac{dy}{dx} + n^2 y = 0;$$

$$y = e^{-mx} \beta \cos (x \sqrt{n^2 - m^2} + \alpha), \quad (n > m).$$

$$7. \quad \frac{d^4 y}{dx^4} - 2 \frac{d^3 y}{dx^3} + 2 \frac{d^2 y}{dx^2} - 2 \frac{dy}{dx} + y = 0;$$

$$y = e^{kx}, \quad (k^2 + 1)(k - 1)^2 = 0;$$

$$\therefore y = e^x (a_1 + a_2 x) + \beta \cos (x + \alpha).$$

$$8. \quad \frac{d^4 y}{dx^4} + 2n^2 \frac{d^2 y}{dx^2} + n^4 y = q;$$

$$y = \frac{q}{n^4} + (a + a_1 x) \cos nx + (b + b_1 x) \sin nx.$$

$$9. \quad \frac{d^4 y}{dx^4} + 4m \frac{d^3 y}{dx^3} + 2(n^2 + 3m^2) \frac{d^2 y}{dx^2} + 4m(n^2 + n^2) \frac{dy}{dx}$$

$$+ (m^2 + n^2)^2 y = q;$$

$$y = e^{kx}, \{ (k+m)^2 + n^2 \}^2 = 0;$$

$$\therefore y = \frac{q}{(m^2 + n^2)^2} + e^{-mx} (a + a_1 x) \cos nx + e^{-mx} (b + b_1 x) \sin nx.$$

$$10. \quad \frac{d^4 y}{dx^4} - 4 \frac{d^3 y}{dx^3} + 6 \frac{d^2 y}{dx^2} - 4 \frac{dy}{dx} + y = 4;$$

$$y - 4 = e^x (c_0 + c_1 x + c_2 x^2 + c_3 x^3).$$

Integration of linear equations with constant coefficients by separation of symbols.

56. Before proceeding to the general case of linear equations, we shall consider the case of a linear equation with all its coefficients constant but having a function of x , X , for the term independent of y . This equation can be shortly treated by the method of separation of symbols: for if we separate the symbol of operation $\frac{d}{dx}$ from its subject y , the equation becomes

$$\left\{ \left(\frac{d}{dx} \right)^n + p_1 \left(\frac{d}{dx} \right)^{n-1} + \dots + p_n \right\} y = X; \dots \dots \dots (1)$$

or if $a_1, a_2, \dots a_n$ be the n distinct roots of the auxiliary equation

$$f(z) = z^n + p_1 z^{n-1} + \dots + p_n = 0,$$

$$\left(\frac{d}{dx} - a_1 \right) \left(\frac{d}{dx} - a_2 \right) \dots \left(\frac{d}{dx} - a_n \right) y = X;$$

$$\therefore y = \left\{ A_1 \left(\frac{d}{dx} - a_1 \right)^{-1} + A_2 \left(\frac{d}{dx} - a_2 \right)^{-1} + \dots + A_n \left(\frac{d}{dx} - a_n \right)^{-1} \right\} X,$$

$A_1, A_2, \&c.$, having the same values as in the resolution of

$$(z^n + p_1 z^{n-1} + \dots + p_n)^{-1}$$

into a similar series of terms, and being therefore known in terms of $a_1, a_2, \&c.$ respectively. (*Integral Calculus*, Art. 34.) But by Leibnitz's Theorem (Art. 53) we have,

$$\left(\frac{d}{dx} + m \right)^{-n} u = e^{-mx} \left(\int dx \right)^n e^{mx} u;$$

and in the present case

$$\left(\frac{d}{dx} - a_1\right)^{-1} X = e^{a_1 x} \int dx e^{-a_1 x} X;$$

$$\therefore y = A_1 e^{a_1 x} \int dx e^{-a_1 x} X + A_2 e^{a_2 x} \int dx e^{-a_2 x} X + \dots + A_n e^{a_n x} \int dx e^{-a_n x} X,$$

the arbitrary constants being reserved under the signs of integration; or if they be expressed, we must add to the above value of y ,

$$C_1 e^{a_1 x} + C_2 e^{a_2 x} + \dots + C_n e^{a_n x},$$

which is called the complementary function, and is the value assumed by y when $X = 0$.

If we denote $\frac{d}{dz} f(z)$ by $f'(z)$, then since $A_1 = \frac{1}{f'(a_1)}$, the above result may be written

$$y = \Sigma \left\{ \frac{e^{ax}}{f'(a)} \int dx e^{-ax} X \right\}.$$

57. If the auxiliary equation have r roots equal to a_1 , then from (1),

$$y = \left(\frac{d}{dx} - a_1\right)^{-r} \left(\frac{d}{dx} - a_{r+1}\right)^{-1} \dots \left(\frac{d}{dx} - a_n\right)^{-1} X,$$

and this function of $\frac{d}{dx}$ may be resolved into partial fractions (*Integral Calculus*, Art. 36), so that

$$y = \left\{ A_1 \left(\frac{d}{dx} - a_1\right)^{-1} + A_2 \left(\frac{d}{dx} - a_1\right)^{-2} + \dots \right. \\ \left. + A_r \left(\frac{d}{dx} - a_1\right)^{-r} + A_{r+1} \left(\frac{d}{dx} - a_{r+1}\right)^{-1} + \dots + A_n \left(\frac{d}{dx} - a_n\right)^{-1} \right\} X.$$

$$\text{But } \left(\frac{d}{dx} - a_1\right)^{-m} X = e^{a_1 x} \left(\frac{d}{dx}\right)^{-m} e^{-a_1 x} X = e^{a_1 x} \int^m dx^m (e^{-a_1 x} X) +$$

$$e^{a_1 x} (c_0 + c_1 x + \dots + c_{m-1} x^{m-1}), \quad c_0, c_1 \text{ \&c.},$$

being the constants introduced after each integration,

$$\begin{aligned}
\therefore y = & A_1 e^{a_1 x} \int dx (e^{-a_1 x} X) + A_2 e^{a_1 x} \int^2 dx^2 (e^{-a_1 x} X) + \dots \\
& + A_r e^{a_1 x} \int^r dx^r (e^{-a_1 x} X). \\
& + A_{r+1} e^{x a_{r+1}} \int dx (e^{-x a_{r+1}} X) + \dots + A_n e^{x a_n} \int dx (e^{-x a_n} X) \\
& + (C_0 + C_1 x + \dots + C_{r-1} x^{r-1}) e^{a_1 x} + C_{r+1} e^{x a_{r+1}} + \dots + C_n e^{x a_n},
\end{aligned}$$

a single constant being substituted for the sum or product of several constants in forming the complementary function.

58. Again suppose

$$a_1 = \rho (\cos \theta + \sqrt{-1} \sin \theta) = \rho e^{\theta \sqrt{-1}}$$

to be an imaginary root of the auxiliary equation, then since A_1 is a function of a_1 it will be of the form

$$R (\cos \alpha + \sqrt{-1} \sin \alpha) = R e^{\alpha \sqrt{-1}},$$

and the term involving a_1 in the value of y will consequently be

$$\begin{aligned}
& R e^{x \rho \cos \theta} e^{(\alpha + x \rho \sin \theta) \sqrt{-1}} \int dx e^{-x \rho \cos \theta} e^{-x \rho \sin \theta \sqrt{-1}} X, \\
& \text{or } R e^{x \rho \cos \theta} \{ \cos (\alpha + x \rho \sin \theta) + \sqrt{-1} \sin (\alpha + x \rho \sin \theta) \} \\
& \times \int dx e^{-x \rho \cos \theta} X \{ \cos (x \rho \sin \theta) - \sqrt{-1} \sin (x \rho \sin \theta) \}.
\end{aligned}$$

Now the term in y involving the conjugate root to a_1 , will result from this by changing the sign of $\sqrt{-1}$; and therefore the sum of the two terms introduced into the value of y by the pair of imaginary roots $\rho (\cos \theta \pm \sqrt{-1} \sin \theta)$, will equal twice the real part of the foregoing expression, that is,

$$\begin{aligned}
& 2 R e^{x \rho \cos \theta} \cos (\alpha + x \rho \sin \theta) \{ \int dx e^{-x \rho \cos \theta} \cos (x \rho \sin \theta) X + C \} \\
& + 2 R e^{x \rho \cos \theta} \sin (\alpha + x \rho \sin \theta) \{ \int dx e^{-x \rho \cos \theta} \sin (x \rho \sin \theta) X + C' \} : \\
& \text{where } 2 R e^{x \rho \cos \theta} \{ C \cos (\alpha + x \rho \sin \theta) + C' \sin (\alpha + x \rho \sin \theta) \},
\end{aligned}$$

are the terms introduced into the complementary function, and may be replaced by

$$e^{x \rho \cos \theta} \{ c \cos (x \rho \sin \theta) + c' \sin (x \rho \sin \theta) \}.$$

59. Exactly in the same way, if the imaginary root

$$a_1 = \rho (\cos \theta + \sqrt{-1} \sin \theta) = \rho e^{\theta \sqrt{-1}}$$

occur r terms in the auxiliary equation, it will produce in the value of y , r terms of the form

$$A_m e^{a_1 x} \int^m dx^m e^{-a_1 x} X;$$

or since A_m is a function of a_1 and may be assumed

$$= R_m (\cos \alpha_m + \sqrt{-1} \sin \alpha_m) = R_m e^{a_m \sqrt{-1}},$$

of the form $R_m e^{x\rho \cos \theta} e^{i(\alpha_m + x\rho \sin \theta)\sqrt{-1}} (dx)^m e^{-x\rho \cos \theta} e^{-x\rho \sin \theta \sqrt{-1}} X$,

$$\text{or } R_m e^{x\rho \cos \theta} \{ \cos (\alpha_m + x\rho \sin \theta) + \sqrt{-1} \sin (\alpha_m + x\rho \sin \theta) \}$$

$$\times (dx)^m e^{-x\rho \cos \theta} \{ \cos (x\rho \sin \theta) - \sqrt{-1} \sin (x\rho \sin \theta) \} X.$$

But the root conjugate to a_1 will produce a term precisely the same as this except with $-\sqrt{-1}$ instead of $+\sqrt{-1}$; consequently the sum of these terms will produce twice the real part of the foregoing expression, that is,

$$2R_m e^{x\rho \cos \theta} \cos (\alpha_m + x\rho \sin \theta) (dx)^m e^{-x\rho \cos \theta} \cos (x\rho \sin \theta) X \\ + 2R_m e^{x\rho \cos \theta} \sin (\alpha_m + x\rho \sin \theta) (dx)^m e^{-x\rho \cos \theta} \sin (x\rho \sin \theta) X;$$

and to get all the terms introduced into the value of y by the pair of imaginary roots $\rho (\cos \theta \pm \sqrt{-1} \sin \theta)$ that occur r times in the auxiliary equation, m in the above formula must receive all values from 1 to r . Also we see that the part of the complementary function introduced by these roots, by substituting a single constant for the sum or product of other arbitrary constants, will take the form

$$(c_0 + c_1 x + \dots c_{r-1} x^{r-1}) e^{x\rho \cos \theta} \cos (x\rho \sin \theta) \\ + (c'_0 + c'_1 x + \dots c'_{r-1} x^{r-1}) e^{x\rho \cos \theta} \sin (x\rho \sin \theta).$$

60. We shall now give an example of each of the cases that have been examined. It may be observed that for equations capable of being reduced to either of the forms

$$f\left(\frac{d}{dx}\right) y = e^{mx}, \quad f\left(\frac{d}{dx}\right) y = p_0 x^m + p_1 x^{m-1} + \dots + p_m,$$

the process may be greatly simplified. For in the former case since $\frac{d}{dx} e^{mx} = me^x$, the symbol $\frac{d}{dx}$ is equivalent to the factor m ; so that the solution of the equation $f\left(\frac{d}{dx}\right)y = e^{mx}$ is $y = \frac{e^{mx}}{f(m)}$ + complementary function. And in the latter case if $\left\{f\left(\frac{d}{dx}\right)\right\}^{-1}$ can be expanded in a series of powers

$$A_0 + A_1 \frac{d}{dx} + A_2 \left(\frac{d}{dx}\right)^2 + \&c.,$$

then as every differential coefficient of X higher than the m^{th} is zero,

$$y = \left\{ A_0 + A_1 \frac{d}{dx} + \dots + A_m \left(\frac{d}{dx}\right)^m \right\} (p_0 x^m + p_1 x^{m-1} + \dots + p_m).$$

Also if $X = \cos(mx + \alpha)$ or $\sin(mx + \alpha)$, since in either case $\frac{d^2 X}{dx^2} = -m^2 X$, if the proposed equation contain only differential coefficients of an even order, that is, be of the form

$$f\left\{\left(\frac{d}{dx}\right)^2\right\}y = \cos(mx + \alpha),$$

then its solution is

$$y = \frac{\cos(mx + \alpha)}{f(-m^2)} + \text{complementary function.}$$

Ex. 1. $\frac{d^3 y}{dx^3} - 6n \frac{d^2 y}{dx^2} + nn^2 \frac{dy}{dx} - 6n^3 y = a^x,$

the roots of the auxiliary equations are $n, 2n, 3n$; and $a^x = e^{x \log a}$;

$$\therefore y = \frac{a^x}{(\log a - n)(\log a - 2n)(\log a - 3n)} + c_0 e^{nx} + c_1 e^{2nx} + c_2 e^{3nx}.$$

Ex. 2. $\frac{d^4 y}{dx^4} - 10 \frac{d^3 y}{dx^3} + 62 \frac{d^2 y}{dx^2} - 210 \frac{dy}{dx} + 261 y = e^x,$

the roots of the auxiliary equation are $2 \pm 5\sqrt{-1}, 3, 3$; and $f(1) = 104$,

$$\therefore y = \frac{e^x}{104} + (a + a'x) e^{3x} + (b \cos 5x + b' \sin 5x) e^{2x}.$$

Ex. 3. $\frac{d^2y}{dx^2} + 2 \frac{dy}{dx} + y = ax^3,$

$$\therefore y = \left(1 + \frac{d}{dx}\right)^{-2} ax^3 = \left\{1 - 2 \frac{d}{dx} + 3 \left(\frac{d}{dx}\right)^2 - 4 \left(\frac{d}{dx}\right)^3\right\} ax^3 + \&c.$$

$$= a (x^3 - 6x^2 + 18x - 24) + (c + c'x)e^{-x}.$$

Ex. 4. $\frac{d^4y}{dx^4} + 5 \left(\frac{dy}{dx}\right)^2 + 6y = \cos (mx + \alpha).$

$$\text{or } \left\{\left(\frac{d}{dx}\right)^2 + 2\right\} \left\{\left(\frac{d}{dx}\right)^2 + 3\right\} y = \cos (mx + \alpha).$$

$$\therefore y = \frac{\cos (mx + \alpha)}{m^4 - 5m^2 + 6} + C \cos (x\sqrt{2} + \beta) + C' \cos (x\sqrt{3} + \gamma).$$

Ex. 5. $\frac{d^n y}{dx^n} - a^n y = X.$

Let $b = a (\cos \phi + \sqrt{-1} \sin \phi) = ae^{\phi \sqrt{-1}}$ be a root of the auxiliary equation $z^n - a^n = 0$, where $\phi = \frac{2r\pi}{n}$, r being any integer, then if $B(z - b)^{-1}$ be the term corresponding to this root in the resolution of $(z^n - a^n)^{-1}$ into partial fractions,

$$B = \frac{1}{nb^{n-1}} = \frac{e^{-(n-1)\phi \sqrt{-1}}}{na^{n-1}}.$$

Hence the corresponding term in the value of $na^{n-1}y$ is

$$e^{-(n-1)\phi \sqrt{-1}} \left(\frac{d}{dx} - b\right)^{-1} X = e^{-(n-1)\phi \sqrt{-1}} e^{bx} \int dx e^{-bx} X$$

$$= e^{ax \cos \phi} e^{(ax \sin \phi - n\phi + \phi) \sqrt{-1}} \int dx e^{-ax \cos \phi} e^{-ax \sin \phi \sqrt{-1}} X$$

$$= e^{ax \cos \phi} \{\cos (ax \sin \phi - n\phi + \phi) + \sqrt{-1} \sin (ax \sin \phi - n\phi + \phi)\}$$

$$\times \int dx e^{-ax \cos \phi} \{\cos (ax \sin \phi) - \sqrt{-1} \sin (ax \sin \phi)\} X;$$

and as the root conjugate to b will produce a term exactly the same as this, only with $-\sqrt{-1}$ instead of $+\sqrt{-1}$, doubling the real part of this expression, we get

$$\frac{2e^{ax \cos \phi}}{na^{n-1}} \cos (ax \sin \phi - n\phi + \phi) \int dx \cos (ax \sin \phi) X e^{-ax \cos \phi} \\ + \frac{2e^{ax \cos \phi}}{na^{n-1}} \sin (ax \sin \phi - n\phi + \phi) \int dx \sin (ax \sin \phi) X e^{-ax \cos \phi}$$

for the general term in the value of y , where $\phi = \frac{1}{n} 2r\pi$, and to get all the terms, r must be taken from 0 to $\frac{1}{2}(n-1)$ or $\frac{1}{2}n$, according as n is odd or even; only the terms that result from this by making $r=0$ or $r=\frac{1}{2}n$, must be divided by 2.

Ex. 6. $\frac{d^n y}{dx^n} + a^n y = X$. The process, and the expression for the general term in the value of y , are exactly the same as in the preceding example, except that $\phi = \frac{1}{n} (2r+1)\pi$, where r is any integer; and to get all the terms we must take r from 0 to $\frac{1}{2}(n-1)$ or $\frac{1}{2}n-1$ according as n is odd or even; only the term that results from taking $r=\frac{1}{2}(n-1)$ must be divided by 2.

$$\text{Ex. 7. } \frac{d^{2n} y}{dx^{2n}} - 2 \cos \theta a^n \frac{d^n y}{dx^n} + a^{2n} y = X.$$

Let $b = a (\cos \phi + \sqrt{-1} \sin \phi) = ae^{\phi \sqrt{-1}}$ be a root of the auxiliary equation $z^{2n} - 2 \cos \theta a^n z^n + a^{2n} = 0$, where $\phi = \frac{1}{n} (2r\pi + \theta)$ r being any integer. Then if $B(z-b)^{-1}$ be the term corresponding to this root in the resolution of $(z^{2n} - 2 \cos \theta a^n z^n + a^{2n})^{-1}$, into partial fractions, since $b^n = a^n (\cos \theta + \sqrt{-1} \sin \theta)$,

$$B = \frac{1}{2n (b^n - a^n \cos \theta) b^{n-1}} = \frac{1}{2n \sqrt{-1} \sin \theta a^n b^{n-1}} = \frac{e^{-(n-1)\phi \sqrt{-1}}}{2n \sqrt{-1} \sin \theta a^{2n-1}}.$$

Hence the corresponding term in the value of $n \sin \theta a^{2n-1} y$ is

$$\frac{1}{2 \sqrt{-1}} e^{-(n-1)\phi \sqrt{-1}} \left(\frac{d}{dx} - b \right)^{-1} X = \frac{1}{2 \sqrt{-1}} e^{-(n-1)\phi \sqrt{-1}} e^{bx} \int dx e^{-bx} X,$$

$$\text{or } \frac{1}{2 \sqrt{-1}} e^{ax \cos \phi} e^{(ax \sin \phi - n\phi + \phi) \sqrt{-1}} \int dx e^{-ax \cos \phi} e^{-ax \sin \phi \sqrt{-1}} X,$$

$$\text{or } \frac{1}{2\sqrt{-1}} e^{ax \cos \phi} \{ \cos (ax \sin \phi - n\phi + \phi) + \sqrt{-1} \sin (ax \sin \phi - n\phi + \phi) \} \\ \times \int dx e^{-ax \cos \phi} \{ \cos (ax \sin \phi) - \sqrt{-1} \sin (ax \sin \phi) \} X.$$

Hence taking double the real part of this expression we get for the general term of the value of $n \sin \theta a^{2n-1} y$

$$e^{ax \cos \phi} \sin (ax \sin \phi - n\phi + \phi) \int dx e^{-ax \cos \phi} \cos (ax \sin \phi) X \\ - e^{ax \cos \phi} \cos (ax \sin \phi - n\phi + \phi) \int dx e^{-ax \cos \phi} \sin (ax \sin \phi) X,$$

when $\phi = \frac{1}{n} (2r\pi + \theta)$, and to get all the terms, r must be taken from 0 to $n - 1$.

61. The solution of $f\left(\frac{d}{dx}\right) y = e^{nx}$ in the form

$$y = \frac{e^{nx}}{f(m)} + c_1 e^{a_1 x} + \&c.,$$

fails, when $m = a_1$ a root of the auxiliary equation. Suppose that a_1 , the root to which m becomes equal, occurs twice in the auxiliary equation, so that $f(a_1) = f'(a_1) = 0$; then by altering the constants the value of y may be written

$$y = \frac{1}{f(m)} (e^{mx} - e^{a_1 x} - h x e^{a_1 x}) + (c + c' x) e^{a_1 x} + \&c.$$

Now let $m = a_1 + h$ where h is a small quantity, then

$$y = e^{a_1 x} \cdot \frac{\frac{1}{2} h^2 x^2 + \frac{1}{6} h^3 x^3 + \&c.}{\frac{1}{2} h^2 f''(a_1) + \&c.} + (c + c' x) e^{a_1 x} + \&c.,$$

or, making $h = 0$,

$$y = \frac{x^2 e^{a_1 x}}{f''(a_1)} + (c + c' x) e^{a_1 x} + \&c.$$

And if the root a_1 to which m becomes equal, occur r times,

$$y = \frac{x^r e^{a_1 x}}{f^{(r)}(a_1)} + (c_0 + c_1 x + \dots + c_{r-1} x^{r-1}) e^{a_1 x} + \&c.$$

These results are easily obtained by differentiating the

numerator and denominator of the vanishing fraction $\frac{e^{mx} - e^{a_1x}}{f(m)}$ in the usual way.

The principle employed here and in all similar cases is to suppose the arbitrary constant to have such a value as to make the expression which becomes infinite, assume the indeterminate form $\frac{0}{0}$; then its real value can be assigned by the ordinary rules. In the instances which follow, a negative value is assumed for the arbitrary constant, which becomes infinite; so that the difference between the two infinite quantities may be finite.

Ex. 1. $\frac{d^4y}{dx^4} - a^4y = e^{ax}$. First suppose the second member to be e^{mx} , then

$$y = \frac{e^{mx} - e^{ax}}{m^4 - a^4} + c_1e^{ax} + c_2e^{-ax} + \beta \cos(ax + \alpha);$$

hence making $m = a$,

$$y = \frac{xe^{ax}}{4a^3} + c_1e^{ax} + c_2e^{-ax} + \beta \cos(ax + \alpha).$$

Ex. 2. $\frac{d^4y}{dx^4} - a^4y = \cos ax$. Suppose the second member to be $\cos mx$, then

$$y = \frac{\cos mx - \cos ax}{m^4 - a^4} + c_1e^{ax} + c_2e^{-ax} + c_3 \cos ax + c_4 \sin ax;$$

hence making $m = a$,

$$y = -\frac{x \sin ax}{4a^3} + \text{complementary function.}$$

Ex. 3. $\frac{d^2y}{dx^2} + n^2y = \cos nx$,

$$y = \frac{x \sin nx}{2n} + c \cos nx + c' \sin nx.$$

62. To deduce the solution of the linear equation of the n^{th} order with variable coefficients

$$\frac{d^n y}{dx^n} + p_1 \frac{d^{n-1} y}{dx^{n-1}} + p_2 \frac{d^{n-2} y}{dx^{n-2}} + \dots + p_n y = X,$$

from the solution of the same equation when the term independent of y is supposed to become zero.

Let $u_1, u_2, \dots u_n$ be the n particular values of u which satisfy the equation

$$\frac{d^n u}{dx^n} + p_1 \frac{d^{n-1} u}{dx^{n-1}} + p_2 \frac{d^{n-2} u}{dx^{n-2}} + \dots + p_n u = 0 \dots\dots\dots (1),$$

with which the proposed coincides if we suppose its second member to become zero; then, as already proved,

$$u = c_1 u_1 + c_2 u_2 + \dots + c_n u_n.$$

If we now divide both sides by u_1 and differentiate, we shall eliminate c_1 ; next dividing both sides by the coefficient of c_2 in the new result which suppose v_1 , and differentiating, we shall eliminate c_2 ; again dividing by v_2 the coefficient which now affects c_3 and differentiating, we shall eliminate c_3 ; and proceeding in this manner till all the constants are eliminated, our final result will be of the form (where each symbol of differentiation affects the whole of the expression after it)

$$\frac{d}{dx} \frac{1}{v_{n-1}} \frac{d}{dx} \frac{1}{v_{n-2}} \dots \frac{d}{dx} \frac{1}{v_1} \frac{d}{dx} \frac{u}{u_1} = 0,$$

in which the coefficient of $\frac{d^n u}{dx^n}$ is evidently $\frac{1}{v_{n-1} v_{n-2} \dots v_1 u_1}$; therefore dividing by this, so as to make the coefficient of $\frac{d^n u}{dx^n}$ unity, we get

$$v_{n-1} v_{n-2} \dots v_1 u_1 \frac{d}{dx} \frac{1}{v_{n-1}} \frac{d}{dx} \frac{1}{v_{n-2}} \dots \frac{d}{dx} \frac{1}{v_1} \frac{d}{dx} \frac{u}{u_1},$$

which must be equivalent to the first member of equation (1). Therefore the same expression, only with y instead of u , must be equivalent to the first member of the proposed equation, and therefore equal to X . Hence equating these equals, and reversing the operations, we find the integral of the proposed equation,

$$y = u_1 \int dx v_1 \int dx v_2 \dots \int dx v_{n-1} \int dx \frac{X}{u_1 v_1 v_2 \dots v_{n-1}};$$

(each symbol of integration affecting the whole of the expression which follows it), which is a general formula for deducing the solution of a complete linear equation of the n^{th} order, from the

solution of the same equation when the term independent of y is supposed to become zero.

63. The application of this method is seen in the following examples.

Ex. 1. $\frac{d^2y}{dx^2} + 4\frac{dy}{dx} + 4y = ax^2.$

If we suppose u to denote the value of y in this equation when $a = 0$, since the roots of the auxiliary equation are $-2, -2$,

$$u = e^{-2x}(c_1 + c_2x); \quad \therefore \frac{d^2}{dx^2}(e^{2x}u) = 0,$$

in which expression the coefficient of $\frac{d^2u}{dx^2}$ is e^{2x} , therefore dividing by this coefficient, and changing u into y , we get

$$e^{-2x} \frac{d^2}{dx^2}(e^{2x}y) = ax^2;$$

$$\therefore y = e^{-2x} \int^2 dx^2 (ax^2 e^{2x}) = a \left(\frac{1}{4} x^2 - \frac{1}{2} x + \frac{3}{8} \right) + e^{-2x}(c_1 + c_2x).$$

Ex. 2. $\frac{d^2y}{dx^2} - 2m\frac{dy}{dx} + (m^2 + n^2)y = X.$

If we suppose u to be the value of y in this equation when $X = 0$, we have

$$u = C_1 e^{mx} \cos nx + C_2 e^{mx} \sin nx;$$

\therefore dividing by $e^{mx} \cos nx$, and differentiating, we find

$$\frac{d}{dx} e^{-mx} \sec nx \cdot u = C_2 n \sec^2 nx;$$

$$\therefore \frac{d}{dx} \cos^2 nx \frac{d}{dx} e^{-mx} \sec nx \cdot u = 0,$$

in which expression the coefficient of $\frac{d^2u}{dx^2}$ is $e^{-mx} \cos nx$, therefore dividing by this coefficient and changing u into y , we get

$$e^{mx} \sec nx \frac{d}{dx} \cos^2 nx \frac{d}{dx} e^{-mx} \sec nx \cdot y = X;$$

$$\therefore y = e^{mx} \cos nx \int dx \sec^2 nx \int dx e^{-mx} \cos nx X,$$

or, if we suppose the constants to be added after each integration,

$$y = e^{mx} (C \cos nx + C' \sin nx) + e^{mx} \cos nx \int dx \sec^2 nx \int dx e^{-mx} \cos nx X.$$

Hence the solution of $\frac{d^2y}{dx^2} + n^2y = X$ is

$$y = C \cos nx + C' \sin nx + \cos nx \int dx \sec^2 nx \int dx \cos nx X.$$

Also if we take $X = A \cos (mx + \alpha) + B \cos (nx + \beta)$,

then $\int dx \cos nx X$

$$= \frac{A}{m^2 - n^2} \{m \sin (mx + \alpha) \cos nx - n \cos (mx + \alpha) \sin nx\}$$

$$+ \frac{B}{2} \left\{ \frac{1}{2n} \sin (2nx + \beta) + x \cos \beta \right\};$$

$$\therefore \int dx \sec^2 nx \int dx \cos nx X = -\frac{A}{m^2 - n^2} \frac{\cos (mx + \alpha)}{\cos nx}$$

$$+ \frac{B}{2} \left\{ \frac{x \sin (nx + \beta)}{n \cos nx} - \frac{\sin \beta}{2n^2} \tan nx \right\};$$

$$\therefore y = C \cos nx + C' \sin nx + \frac{A}{n^2 - m^2} \cos (mx + \alpha)$$

$$+ \frac{B}{2n} x \sin (nx + \beta).$$

Ex. 3. $\frac{d^2y}{dx^2} + 2n \frac{dy}{dx} - \frac{2y}{x^2} = \frac{a}{x} \left(nx + \frac{1}{nx} - 1 \right);$

here $u = c_1 \left(1 - \frac{1}{nx} \right) + c_2 \left(1 + \frac{1}{nx} \right) e^{-2nx}$, (Ex. 2. Art. 64.)

and $y = \left(\frac{1}{2} ax + C' \right) \left(1 - \frac{1}{nx} \right) + C \left(1 + \frac{1}{nx} \right) e^{-2nx}.$

64. As this method cannot be applied till we have obtained a solution of the proposed equation deprived of the term independent of y , the following propositions are sometimes of use for equations of the second order.

If $y = u$ be a particular integral of the equation

$$\frac{d^2y}{dx^2} + P \frac{dy}{dx} + Qy = 0,$$

the other particular integral will be $u \int dx (u^{-2} e^{-\int dx P})$. For if we denote the two values of y by u and v , we have

$$\frac{d^2u}{dx^2} + P \frac{du}{dx} + Qu = 0, \quad \frac{d^2v}{dx^2} + P \frac{dv}{dx} + Qv = 0,$$

$$\therefore u \frac{d^2 v}{dx^2} - v \frac{d^2 u}{dx^2} + P \left(u \frac{dv}{dx} - v \frac{du}{dx} \right) = 0,$$

$$\therefore u \frac{dv}{dx} - v \frac{du}{dx} = C e^{-f dx P}, \text{ or } \frac{d}{dx} \left(\frac{v}{u} \right) = C u^{-2} e^{-f dx P};$$

$$\therefore v = C u \int dx (u^{-2} e^{-f dx P}).$$

Ex. 1. $\frac{d^2 y}{dx^2} + mP \frac{dy}{dx} - m^2 (P + 1) y = 0.$

It is easily seen that $y = u = e^{mx}$ is a solution;

$$\therefore y = e^{mx} (C' + C \int dx e^{-2mx - f dx P}).$$

Suppose $m = 1$, $P = -\frac{2}{x+3}$, $-f dx P = \log (x+3)^2$;

$$\therefore y = e^x \{ C' + C \int dx e^{-2x} (x+3)^2 \} = C' e^x + C e^{-x} (x^2 + 7x + 12\frac{1}{2})$$

is the solution of

$$(x+3) \frac{d^2 y}{dx^2} - 2 \frac{dy}{dx} - (x+1) y = 0.$$

Ex. 2. $\frac{d^2 y}{dx^2} + 2n \frac{dy}{dx} - \frac{2y}{x^2} = 0$, of which $y = u = 1 - \frac{1}{nx}$ is a solu-

tion, and thence we get $y = C' \left(1 - \frac{1}{nx} \right) + C \left(1 + \frac{1}{nx} \right) e^{-2nx}$.

Ex. 3. $\frac{d^2 y}{dx^2} + \frac{1}{x} \left(n + \frac{m}{1+x^m} \right) \frac{dy}{dx} - \frac{m+1}{4x^2} \left(m+2n-1 + \frac{2m}{1+x^m} \right) y = 0$;

$y = x^{\frac{m+1}{2}}$, from which the complete integral may be obtained.

Ex. 4. $\frac{d^2 y}{dx^2} + \frac{5+2x^3}{x+x^4} \frac{dy}{dx} - \frac{12+6x^3}{x^2+x^5} y = 0$;

of which $y = u = x^2$ is a solution.

$$P = -\frac{5x^{-6} - 2x^{-3}}{x^{-3} + x^{-2}}, \quad \therefore \int dx P = -\log (x^{-3} + x^{-2})$$

$$\therefore \frac{1}{u^2} e^{-f dx P} = x^{-3} + x^{-2}, \quad \therefore u \int dx \frac{e^{-f dx P}}{u^2} = -\frac{x^{-6}}{8} - \frac{x^{-3}}{5},$$

$$y = Cx^2 + C' \left(\frac{1}{x^6} + \frac{8}{5} \frac{1}{x^3} \right).$$

65. It is easily seen, by actual substitution, that if $y = u$ satisfy the equation

$$\frac{d^2y}{dx^2} + P \frac{dy}{dx} + n^2y = 0,$$

then the equation

$$\frac{d^2y}{dx^2} - P \frac{dy}{dx} + n^2y = 0$$

will be satisfied by the value

$$y = \frac{du}{dx} e^{\int dx P}.$$

Thus for the equation $\frac{d^2y}{dx^2} + \frac{4}{x} \frac{dy}{dx} + n^2y = 0$

$$y = u = Cx^{-3} \{ \cos (nx + \alpha) + nx \sin (nx + \alpha) \};$$

$$\therefore \frac{du}{dx} = Cx^{-3} \{ (n^2x^2 - 3) \cos (nx + \alpha) - 3nx \sin (nx + \alpha) \},$$

$$\text{and } e^{\int dx P} = x^4,$$

$$\therefore y = C \{ (n^2x^2 - 3) \cos (nx + \alpha) - 3nx \sin (nx + \alpha) \}$$

is the complete solution of $\frac{d^2y}{dx^2} - \frac{4}{x} \frac{dy}{dx} + n^2y = 0$.

66. If we know a particular integral of a linear equation of any order that has no term independent of y , we may reduce it to another equation of the same kind, of the order immediately inferior.

Let $y = u$ be a particular integral of $f\left(\frac{d}{dx}\right)y = 0$,

$$\text{or } \frac{d^n y}{dx^n} + p_1 \frac{d^{n-1} y}{dx^{n-1}} + p_2 \frac{d^{n-2} y}{dx^{n-2}} + \dots + p_n y = 0, \dots\dots\dots (1),$$

and assume $y = u \int dx z$, z being a new variable, a function of x ; then, separating the symbols of operation from those of quantity, we have

$$\frac{d^n y}{dx^n} = \left(\frac{d}{dx} + \frac{d'}{dx} \right)^n u \int dx z,$$

$\frac{d}{dx}$ being understood to affect u only, and $\frac{d'}{dx}$ to affect $\int dxz$ only.

Hence, substituting for the differential coefficients in the proposed, by this formula, and again separating the symbols of operation from those of quantity, we get successively

$$\begin{aligned} & \left(\frac{d}{dx} + \frac{d'}{dx}\right)^n u \int dxz + p_1 \left(\frac{d}{dx} + \frac{d'}{dx}\right)^{n-1} u \int dxz + \&c. + p_n u \int dxz \\ &= f\left(\frac{d}{dx} + \frac{d'}{dx}\right) u \int dxz \\ &= \int dxz \cdot f\left(\frac{d}{dx}\right) u + z f'\left(\frac{d}{dx}\right) u + \frac{1}{1 \cdot 2} \frac{dz}{dx} \cdot f''\left(\frac{d}{dx}\right) u + \dots \\ &+ \frac{1}{n-1} \frac{d^{n-2}z}{dx^{n-2}} f^{(n-1)}\left(\frac{d}{dx}\right) u + \frac{d^{n-1}z}{dx^{n-1}} \cdot u. \end{aligned}$$

But since u is a particular value of y , $f\left(\frac{d}{dx}\right)u = 0$; hence reversing the order of the terms and observing that $f^{(n)}\left(\frac{d}{dx}\right)$ denotes the same function of the symbol $\frac{d}{dx}$, that $\frac{d^r f(v)}{dv^r}$ does of v , we get for the depressed equation

$$\begin{aligned} & u \frac{d^{n-1}z}{dx^{n-1}} + \left(n \frac{du}{dx} + p_1 u\right) \frac{d^{n-2}z}{dx^{n-2}} \\ &+ \left\{ \frac{n(n-1)}{1 \cdot 2} \frac{d^2u}{dx^2} + (n-1)p_1 \frac{du}{dx} + p_2 u \right\} \frac{d^{n-3}z}{dx^{n-3}} + \dots + z f'\left(\frac{d}{dx}\right) u = 0. \quad (2) \end{aligned}$$

Similarly, if we know another particular solution u_1 of equation (1), then $\frac{d}{dx} \left(\frac{u_1}{u}\right)$ will be a value of z in equation (2), and we may depress this equation to another of the same form of the $(n-2)^{\text{th}}$ order; and if we know r particular solutions of equation (1), we may in this way depress it to an equation of the same form of the $(n-r)^{\text{th}}$ order.

Method of Parameters.

67. There is also another mode of integrating linear equations, which deserves to be mentioned, called the method of Parameters, which we shall now explain. It may be stated thus. The complete integral of the linear equation of the n^{th} order will be of the form

$$y = v_1 u_1 + v_2 u_2 + \dots + v_n u_n,$$

where u_1, u_2, \dots, u_n are the n particular integrals of the equation when the term independent of y becomes zero; and v_1, v_2, \dots, v_n are functions of x , determined by equations of the form

$$v_1 = \int dx f_1(x) + C_1.$$

Let the proposed equation be

$$\frac{d^n y}{dx^n} + p_1 \frac{d^{n-1} y}{dx^{n-1}} + \dots + p_n y = X,$$

and suppose $y = v_1 u_1 + v_2 u_2 + \dots + v_n u_n = \Sigma (vu)$,

and as we have made only one assumption respecting the n independent quantities v_1, v_2, \dots, v_n , we may make $n-1$ more;

$$\text{now } \frac{dy}{dx} = \Sigma \left(v \frac{du}{dx} \right) + \Sigma \left(u \frac{dv}{dx} \right); \quad \therefore \frac{dy}{dx} = \Sigma \left(v \frac{du}{dx} \right),$$

putting for the first of our additional assumptions $\Sigma \left(u \frac{dv}{dx} \right) = 0, (1);$

similarly $\frac{d^2 y}{dx^2} = \Sigma \left(v \frac{d^2 u}{dx^2} \right)$, putting $\Sigma \left(\frac{du}{dx} \frac{dv}{dx} \right) = 0, (2), \&c. = \&c.$

$$\text{and } \frac{d^n y}{dx^n} = \Sigma \left(v \frac{d^n u}{dx^n} \right) + X, \text{ putting } \Sigma \left(\frac{d^{n-1} u}{dx^{n-1}} \frac{dv}{dx} \right) = X, (n).$$

Substitute these values for $y, \frac{dy}{dx}, \frac{d^2 y}{dx^2}, \dots, \frac{d^n y}{dx^n}$, in the proposed equation, and the first member becomes

$$\begin{aligned} v_1 \left(\frac{d^n u_1}{dx^n} + p_1 \frac{d^{n-1} u_1}{dx^{n-1}} + \dots + p_n u_1 \right) &+ v_2 \left(\frac{d^n u_2}{dx^n} + p_1 \frac{d^{n-1} u_2}{dx^{n-1}} + \dots + p_n u_2 \right) \\ &+ \&c. + v_n \left(\frac{d^n u_n}{dx^n} + p_1 \frac{d^{n-1} u_n}{dx^{n-1}} + \dots + p_n u_n \right) + X, \end{aligned}$$

which manifestly reduces itself to X , since each of the quantities within brackets is equal to zero; hence if the parameters $v_1, v_2, \dots v_n$ be determined subject to the above n equations of condition, the assumed value of y will satisfy the equation; and since there are n equations in which the n quantities

$$\frac{dv_1}{dx}, \frac{dv_2}{dx}, \dots \frac{dv_n}{dx}$$

enter to the first degree, each of the parameters v_1, v_2 , &c. will be determined by an equation of the form $\frac{dv_1}{dx} = f_1(x)$, and so n arbitrary constants will be introduced.

Ex. $\frac{d^2y}{dx^2} + n^2y = \sec nx$.

The solution of $\frac{d^2y}{dx^2} + n^2y = 0$ is $y = a_1 \cos nx + a_2 \sin nx$.

Let $\therefore y = v_1 \cos nx + v_2 \sin nx$;

$$\therefore \frac{dy}{dx} = -v_1 n \sin nx + v_2 n \cos nx,$$

$$\text{making } \frac{dv_1}{dx} \cos nx + \frac{dv_2}{dx} \sin nx = 0.$$

$$\frac{d^2y}{dx^2} = -v_1 n^2 \cos nx - v_2 n^2 \sin nx + \sec nx,$$

$$\text{making } -\frac{dv_1}{dx} n \sin nx + \frac{dv_2}{dx} n \cos nx = \sec nx;$$

$$\therefore \frac{dv_1}{dx} = -\frac{1}{n} \frac{\sin nx}{\cos nx}, \quad \text{or } v_1 = \frac{1}{n^2} \log \cos nx + C_1.$$

$$\frac{dv_2}{dx} = \frac{1}{n}, \quad \text{or } v_2 = \frac{x}{n} + C_2;$$

$$\therefore y = \left(\frac{1}{n^2} \log \cos nx + C_1 \right) \cos nx + \left(\frac{x}{n} + C_2 \right) \sin nx.$$

Change of the Independent Variable.

68. Besides linear equations with constant coefficients, very few equations of the second and superior orders admit of being integrated; and those only by particular methods. Sometimes

the integration of an equation may be facilitated by changing the independent variable; and this is the method which has been attended with the greatest success. Thus in the example

$$\frac{d^2y}{dx^2} + a \left(\frac{dy}{dx} \right)^2 + bx \left(\frac{dy}{dx} \right)^3 = 0,$$

if we make y the independent variable, and consequently for

$$\frac{dy}{dx}, \frac{d^2y}{dx^2}, \text{ write } \frac{1}{\frac{dx}{dy}}, -\frac{d^2x}{dy^2} \div \left(\frac{dy}{dx} \right)^3,$$

the equation is transformed into the integrable shape

$$\frac{d^2x}{dy^2} - a \frac{dx}{dy} - bx = 0.$$

69. In the above example there is no difficulty in fixing upon the new independent variable. In other cases we must consider x and y as functions of a third variable t , and substitute the values of $\frac{dy}{dx}$, $\frac{d^2y}{dx^2}$, &c. corresponding to that supposition. When the equation is thus generalized, we must assume for x or y some known function of t , adapted to the particular form of the equation, so that there may arise for determining the function of t which expresses y or x , a differential equation simpler than the proposed one; between the integral of which and the assumed function, if we eliminate t , we obtain the required relation between x and y .

Ex. 1. $(1-x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} + n^2y = 0.$

The generalized equation is

$$(1-x^2) \frac{x'y'' - y'x''}{x'^3} - x \frac{y'}{x'} + n^2y = 0,$$

using x' , x'' , &c. to denote $\frac{dx}{dt}$, $\frac{d^2x}{dt^2}$, &c.

Let $x = \cos t$, $\therefore x' = -\sin t$, $x'' = -\cos t$, therefore by substitution $\frac{d^2y}{dt^2} + n^2y = 0$, which gives

$$y = A \cos nt + B \sin nt;$$

$$\therefore y = A \cos (n \cos^{-1} x) + B \sin (n \cos^{-1} x).$$

Ex. 2. $x^4 \frac{d^2 y}{dx^2} + cy = 0.$

The generalized equation is

$$x^4 \cdot \frac{x'y'' - y'x''}{x'^3} + cy = 0;$$

$$\text{let } x' = -x^2, \text{ or } x = \frac{1}{t}, \therefore x'' = 2x^3,$$

hence by substitution we get

$$y'' + \frac{2}{t} y' + cy = 0, \text{ or } \frac{d^2 (ty)}{dt^2} + cty = 0;$$

$$\therefore ty = \beta \cos (t\sqrt{c} + \alpha), \text{ or } y = \beta x \cos \left(\frac{\sqrt{c}}{x} + \alpha \right).$$

If $c = -n^2$, the solution is $y = x (c_1 e^{\frac{n}{x}} + c_2 e^{-\frac{n}{x}}).$

Ex. 3. $u^2 \frac{d^2 y}{dx^2} + \left(\frac{du}{dx} + a \right) u \frac{dy}{dx} + by = X,$

where u denotes a given function of x . This when generalized becomes

$$u^2 \cdot \frac{x'y'' - y'x''}{x'^3} + \left(\frac{du}{dx} + a \right) u \frac{y'}{x'} + by = X;$$

$$\text{let } x' = u, \text{ or } t = \int dx \frac{1}{u}, \therefore \frac{du}{dx} = \frac{x''}{x'},$$

hence by substitution we get the integrable form

$$\frac{d^2 y}{dt^2} + a \frac{dy}{dt} + by = X.$$

Ex. 4. $x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} - y = \frac{ax^2}{x^2 - 1}.$

Here the substitution $x = e^t$ gives

$$\frac{d^2 y}{dt^2} - y = a + \frac{a}{e^{2t} - 1};$$

$$\begin{aligned}\therefore y &= c_1 e^x + c_2 e^{-x} + \frac{1}{2}a + \frac{1}{4}a(e^{-x} - e^x) \log \left(\frac{e^x - 1}{e^x + 1} \right) \\ &= c_1 x + \frac{c_2}{x} + \frac{1}{2}a + \frac{1}{4}a \left(\frac{1}{x} - x \right) \log \left(\frac{x-1}{x+1} \right).\end{aligned}$$

Ex. 5. $x^3 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} - y = ax^n,$

$$y = c_1 x + \frac{c_2}{x} + \frac{ax^n}{n^2 - 1}.$$

Ex. 6. $\frac{d^2 y}{dx^2} + \frac{A}{a+bx} \frac{dy}{dx} + \frac{B}{(a+bx)^2} y = 0.$

The generalized equation is

$$\frac{x'y'' - y'x''}{x'^3} + \frac{A}{a+bx} \frac{y'}{x'} + \frac{B}{(a+bx)^2} y = 0.$$

Let $x' = a + bx$, or $e^x = a + bx$; $\therefore x'' = bx'$;

hence, substituting,

$$\frac{d^2 y}{dt^2} + (A - b) \frac{dy}{dt} + By = 0;$$

the solution of which is

$$y = c_1 e^{mt} + c_2 e^{m't} = c_1 (a + bx)^{\frac{m}{b}} + c_2 (a + bx)^{\frac{m'}{b}},$$

m and m' being the real roots of $m^2 + (A - b)m + B = 0.$

If the roots be impossible, and of the form $m \pm n\sqrt{-1}$, the solution is

$$y = \beta e^{mt} \cos (nt + \alpha) = \beta (a + bx)^{\frac{m}{b}} \cos \{ \log (a + bx)^{\frac{n}{b}} + \alpha \}.$$

The same substitution of course succeeds for the equation

$$(a + bx)^n \frac{d^n y}{dx^n} + A (a + bx)^{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \&c.$$

$$+ K (a + bx) \frac{dy}{dx} + Ly = f(x),$$

which, by putting $a + bx = z$, may first of all be reduced to the form

$$z^n \frac{d^n y}{dz^n} + a_1 z^{n-1} \frac{d^{n-1} y}{dz^{n-1}} + \&c. + a_{n-1} z \frac{dy}{dz} + a_n y = f_1(z);$$

and then to a linear equation with constant coefficients, by the formula

$$z^n \frac{d^n y}{dz^n} = \frac{d^n y}{dt^n} - p_1 \frac{d^{n-1} y}{dt^{n-1}} + \&c. \pm p_{n-1} y,$$

where $z = e^t$, and $p_1, p_2, \&c.$ are such that the roots of

$$k^n - p_1 k^{n-1} + p_2 k^{n-2} - \&c. \dots \pm p_{n-1} k = 0$$

are $0, 1, 2, 3 \dots (n-1)$; so that the preceding formula may be written

$$\begin{aligned} z^n \frac{d^n y}{dz^n} &= \left\{ \left(\frac{d}{dt} \right)^n - p_1 \left(\frac{d}{dt} \right)^{n-1} + \dots \pm p_n \right\} y \\ &= \left\{ \frac{d}{dt} \left(\frac{d}{dt} - 1 \right) \left(\frac{d}{dt} - 2 \right) \dots \left(\frac{d}{dt} - n + 1 \right) \right\} y. \end{aligned}$$

70. The result just noticed

$$x^n \frac{d^n y}{dx^n} = \frac{d}{dt} \left(\frac{d}{dt} - 1 \right) \dots \left(\frac{d}{dt} - n + 1 \right) y,$$

where $x = e^t$, may be written

$$x^n \frac{d^n y}{dx^n} = P_n \left(\frac{d}{dt} \right) y,$$

using the symbol P_n to denote the product of the n factors formed by subtracting from $\frac{d}{dt}$ the n numbers $0, 1, 2, \dots n-1$. Hence multiplying by $x^m = e^{mt}$, we get

$$x^{m+n} \frac{d^n y}{dx^n} = e^{mt} P_n \left(\frac{d}{dt} + m - m \right) y.$$

But by Art. 53 we have

$$f \left(\frac{d}{dt} \right) e^{mt} y = e^{mt} f \left(\frac{d}{dt} + m \right) y,$$

which shews that in the expression $f\left(\frac{d}{dt}\right) e^{mt} y$ we may remove the power of e^t from the operation of the symbol $f\left(\frac{d}{dt}\right)$, and prefix it as a factor of the whole expression, provided we add the index of the power to $\frac{d}{dt}$; and conversely in the expression $e^{mt} f\left(\frac{d}{dt} + m\right) y$ we may transfer the power of e^t to be a factor of y , and so bring it under the symbol of operation, by subtracting the index of the power from $\frac{d}{dt} + m$. Hence

$$x^{m+n} \frac{d^n y}{dx^n} = P_n \left(\frac{d}{dt} - m \right) e^{mt} y.$$

71. Hence an expression such as

$$(a + bx + cx^2 + \dots + lx^r) x^n \frac{d^n y}{dx^n}$$

is transformed by the substitution $x = e^t$ into

$$aP_n(d)y + bP_n(d-1)e'y + cP_n(d-2)e^2y + \dots + lP_n(d-r)e^ry,$$

where for convenience d is written for $\frac{d}{dt}$, the differentials of t being omitted; and consequently an equation such as

$$(a + bx + cx^2 + \&c.) x^n \frac{d^n y}{dx^n} + (a' + b'x + c'x^2 + \&c.) x^{n-1} \frac{d^{n-1} y}{dx^{n-1}} \\ + (a'' + b''x + \&c.) x^{n-2} \frac{d^{n-2} y}{dx^{n-2}} + \dots + (a^{(n)} + b^{(n)}x + \&c.) y = X$$

is transformed into

$$aP_n(d)y + a'P_{n-1}(d)y + a''P_{n-2}(d)y + \dots + a^{(n)}y \\ + bP_n(d-1)e'y + b'P_{n-1}(d-1)e'y + b''P_{n-2}(d-1)e'y + \dots + b^{(n)}e'y \\ + cP_n(d-2)e^2y + c'P_{n-1}(d-2)e^2y + c''P_{n-2}(d-2)e^2y + \dots + c^{(n)}e^2y \\ + \&c. = T \text{ (a function of } e^t),$$

which is evidently of the form (where $f_0(d)$, $f_1(d)$, &c. denote rational and integral functions of d)

$$f_0(d)y + f_1(d)e'y + f_2(d)e^2y + \&c. = T;$$

and in this form only differential coefficients affected with constant multipliers enter. By this transformation, the object of which is to bring all the variable quantities under the symbols of operation, we can in several cases effect the solution of the equation, or its reduction to a simpler form.

72. Thus suppose the first member reduced to a single term

$$f_r(d)e^ry = T, \text{ or } e^r f_r(d+r)y = T, \text{ or } f_r(d+r)y = e^{-r}T,$$

an ordinary linear equation with constant coefficients. This happens to the equation

$$(a+bx)^n \frac{d^n y}{dx^n} + A(a+bx)^{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \&c. + Ly = X,$$

which by replacing $a+bx$ by x is first transformed into

$$b^n x^n \frac{d^n y}{dx^n} + Ab^{n-1} x^{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \dots + Ly = X,$$

and then, making $x = e^t$, into

$$b^n P_n(d)y + Ab^{n-1}P_{n-1}(d)y + \dots + Ly = T,$$

which is of the form $f_0(d)y = T$.

$$\text{Ex. } x^3 \frac{d^3 y}{dx^3} + ax^2 \frac{d^2 y}{dx^2} + bx \frac{dy}{dx} + cy = X;$$

$$\text{this becomes } \{d^3 + (a-3)d^2 + (-a+b+2)d + c\}y = X,$$

$$\text{if therefore } a=3, b=1, c=-1, \text{ and } X=x^m, \frac{d^3 y}{dt^3} - y = e^{mt},$$

$$\begin{aligned} \text{and } y &= \frac{e^{mt}}{m^3-1} + c_0 + \beta \cos\left(\frac{2\pi t}{3} + \alpha\right) \\ &= \frac{x^m}{m^3-1} + c_0 + \beta \cos\left(\frac{2\pi}{3} \log x + \alpha\right). \end{aligned}$$

Or if $a = -3$, $b = 7$, $c = -8$, then $(d-2)^3 y = X$,

$$\text{and } y = e^{2t} (f dt)^3 e^{-2t} X = x^2 \left(f dx \frac{1}{x} \right)^3 \cdot \frac{X}{x^2},$$

because $f dt$ prefixed to any subject is equivalent to $f dx \frac{1}{x}$.

73. The meaning, in the above and similar cases, of such a notation as $\left(f dx \frac{1}{x} \right)^3 u$ is that the complex operation $f dx \frac{1}{x}$ (which consists in dividing the subject by x , and then integrating it with respect to x) is performed three times in succession; first upon u which gives a result u_1 , then upon u_1 which gives a result u_2 , and thirdly upon u_2 . Thus to get the complementary function in the above result, we must make $X = 0$; then

$$f dx \frac{1}{x} 0 = f dx 0 = C_0, \quad f dx \frac{1}{x} C_0 = C_0 \log x + C_1,$$

$$f dx \frac{1}{x} (C_0 \log x + C_1) = \frac{1}{2} C_0 (\log x)^2 + C_1 (\log x) + C_2;$$

and the complete value of y is

$$y = x^2 \left(f dx \frac{1}{x} \right)^3 \frac{X}{x^2} + x^2 \left\{ \frac{1}{2} C_0 (\log x)^2 + C_1 (\log x) + C_2 \right\}.$$

$$\text{Hence for } x^3 \frac{d^3 y}{dx^3} - 3x^2 \frac{d^2 y}{dx^2} + 7x \frac{dy}{dx} - 8y = \frac{x^3 (1-x)}{(1+x)^3},$$

$$y = x^2 \log(1+x) + x^2 \{c_0 + c_1 \log x + c_2 (\log x)^2\}.$$

74. Next suppose the first member of the transformed equation to consist only of two terms

$$f_m(d) e^{mt} y + f_n(d) e^{nt} y = T,$$

$$\text{or } e^{mt} f_m(d+m) y + e^{nt} f_n(d+m) e^{(n-m)t} y = T;$$

which by dividing by the coefficient of y , is reduced to the form

$$y + \phi(d) e^{rt} y = T_1.$$

Let $v + \psi(d) e^{rt} v = T_2$ be an auxiliary equation whose solution is known for any function T_2 of e^t , as the value of its second mem-

ber, and with which we wish to make the proposed equation coincide. Assume $y = f(d) v$ then

$$f(d) v + \phi(d) e^r f(d) v = T_1,$$

$$\text{or } v + \frac{\phi(d)}{f(d)} \cdot f(d-r) e^r v = \frac{T_1}{f(d)};$$

and in order that this may coincide with the auxiliary equation we must have the two conditions

$$\frac{T_1}{f(d)} = T_2, \quad \frac{\phi(d)}{f(d)} \cdot f(d-r) = \psi(d),$$

the latter of which gives

$$f(d) = \frac{\phi(d)}{\psi(d)} f(d-r) = \frac{\phi(d)}{\psi(d)} \cdot \frac{\phi(d-r)}{\psi(d-r)} \dots = P_r \frac{\phi(d)}{\psi(d)},$$

where P_r denotes the product of an infinite number of factors formed from $\frac{\phi(d)}{\psi(d)}$ by replacing d by $d-r$, $d-2r$, &c. Hence, $f(d)$ being known, $T_2 = \{f(d)\}^{-1} T_1$ can be computed; then v is known from $v + \psi(d) e^r v = T_2$ an equation which is supposed to admit of being solved when T_2 is any given function of e^r , and then $y = P_r \frac{\phi(d)}{\psi(d)} v$.

$$\text{Ex. 1.} \quad x^2 \frac{d^2 y}{dx^2} + (a+1) x \frac{dy}{dx} + (b+q^2 x^2) y = X;$$

$$\therefore d(d-1)y + (a+1) dy + (b+q^2 e^{2x}) y = T,$$

$$\text{or } (d^2 + ad + b) y + q^2 e^{2x} y = T;$$

$$\therefore y + \frac{q^2}{d^2 + ad + b} e^{2x} y = T_1.$$

As the proposed equation can be solved when $a = -1$, $b = 0$, let the auxiliary equation be

$$\frac{d^2 v}{dx^2} + q^2 v = X_2, \text{ or } v + \frac{q^2}{d^2 - d} e^{2x} v = T_2.$$

Assume $y = f(d) v$;

$$\therefore f(d) v + \frac{q^2}{d^2 + ad + b} e^{2x} f(d) v = T_1,$$

$$\text{or } v + \frac{q^2}{d^2 + ad + b} \frac{f(d-2)}{f(d)} e^{av} = \frac{T_1}{f(d)};$$

$$\therefore f(d) = \frac{d^2 - d}{d^2 + ad + b} f(d-2) = P_2 \frac{d^2 - d}{d^2 + ad + b};$$

and such values must be given to the constants a and b as will produce a value of $f(d)$ that will allow y to be determined from the equation $y = f(d) v = f(d) \beta \cos (qx + \alpha)$ when $X = 0$.

If q^2 be preceded by a negative sign, the value to be used for v in this and the following examples will be $\alpha e^{qx} + \beta e^{-qx}$.

$$2. \quad x^2 \frac{d^2 y}{dx^2} - 2nx \frac{dy}{dx} + (2n + q^2 x^2) y = 0;$$

$$f(d) = P_2 \frac{d}{d-2n} = d(d-2) \dots (d-2n+2);$$

$$\therefore y = d(d-2)(d-4) \dots (d-2n+2) \beta \cos (qx + \alpha);$$

but the symbol of operation $\frac{d}{dt} - r$ prefixed to any subject, is equivalent to

$$e^{rt} \frac{d}{dt} e^{-rt} = x^r \cdot x \frac{d}{dx} \frac{1}{x} = x^{r+1} \frac{d}{dx} \frac{1}{x^r};$$

$$\begin{aligned} \therefore y &= x \frac{d}{dx} \cdot x^3 \frac{d}{dx} \frac{1}{x^2} \cdot x^5 \frac{d}{dx} \frac{1}{x^4} \dots x^{2n-1} \frac{d}{dx} \frac{v}{x^{2n-2}} \\ &= \frac{1}{x^2} \left(x^3 \frac{d}{dx} \right)^n \frac{\beta \cos (qx + \alpha)}{x^{2n-2}}. \end{aligned}$$

Let $n = 2$, then the equation is

$$x^2 \frac{d^2 y}{dx^2} - 4x \frac{dy}{dx} + (4 + q^2 x^2) y = 0,$$

and its solution

$$y = \beta \{-q^2 x^2 \cos (qx + \alpha) + qx \sin (qx + \alpha)\}.$$

$$3. \quad x^2 \frac{d^2 y}{dx^2} - 2nx \frac{dy}{dx} + q^2 x^2 y = 0;$$

$$f(d) = P_2 \frac{d-1}{d-2n-1} = (d-1)(d-3) \dots (d-2n+1);$$

$$\therefore y = (d-1)(d-3) \dots (d-2n+1) \beta \cos(qx + \alpha);$$

and, as shewn in the preceding example, $\frac{d}{dt} - r$ prefixed to any subject is equivalent to $x^{r+1} \frac{d}{dx} x^{-r}$;

$$\begin{aligned} \therefore y &= x^2 \frac{d}{dx} \frac{1}{x} \cdot x^4 \frac{d}{dx} \frac{1}{x^3} \cdot x^6 \frac{d}{dx} \frac{1}{x^5} \dots x^{2n} \frac{d}{dx} \frac{v}{x^{2n-1}} \\ &= \frac{1}{x} \left(x^3 \frac{d}{dx} \right)^n \frac{\beta \cos(qx + \alpha)}{x^{2n-1}}. \end{aligned}$$

$$4. \quad x^2 \frac{d^2 y}{dx^2} - n(n+1)y + q^2 x^2 y = 0.$$

If in this equation we replace y by $x^{-n}y$, it is reduced to the preceding example; and consequently its solution is

$$y = \frac{1}{x^{n+1}} \left(x^3 \frac{d}{dx} \right)^n \frac{\beta \cos(qx + \alpha)}{x^{2n-1}}.$$

if we make n equal to 1, 2, we shall find the corresponding values of y to be

$$-\beta q \sin(qx + \alpha) - \frac{\beta}{x} \cos(qx + \alpha);$$

$$\text{and } -\beta \left(q^2 - \frac{3}{x^2} \right) \cos(qx + \alpha) + \frac{3\beta q}{x} \sin(qx + \alpha);$$

α and β denoting arbitrary constants.

$$5. \quad x^2 \frac{d^2 y}{dx^2} + (x + ax + qx^2) \frac{dy}{dx} + by = 0;$$

$$\therefore y + \frac{d-1}{d^2 + ad + b} qe^y = 0.$$

Making $a = -1$, $b = 0$, let the auxiliary equation be

$$v + \frac{qe^v}{d} = 0, \text{ where } v = \alpha + \beta e^{-qx},$$

$$\therefore f(d) = P_1 \frac{d^2 - d}{d^2 + ad + b};$$

and the constants a and b may receive suitable values as in the foregoing examples.

$$6. \quad x^3 \frac{d^2 y}{dx^2} + qx^3 \frac{dy}{dx} - n(n+1)y = 0, \text{ here } a = -1, b = -n(n+1),$$

$$\therefore f(d) = P_1 \frac{d(d-1)}{(d+n)(d-n-1)} = \frac{(d-1)(d-2)\dots(d-n)}{(d+1)(d+2)\dots(d+n)}.$$

But $d-r$ is equivalent to $x^{r+1} \frac{d}{dx} x^{-r}$, and $(d+r)^{-1}$ to $x^{-r} (f dx) x^{r-1}$;

$$\therefore y = \left(x^3 \frac{d}{dx}\right)^n \frac{1}{x^{n-1}} (x^{-2} f dx)^n x^{n-1} (\alpha + \beta e^{-qx}).$$

Let $n = 1$,

$$y = \left(x^3 \frac{d}{dx}\right) x^{-2} f dx (\alpha + \beta e^{-qx}) = x^2 \frac{d}{dx} (\alpha x^{-1} - \frac{\beta}{q} x^{-2} e^{-qx} + C x^{-2})$$

$$= -\alpha - \frac{2C}{x} + \beta \left(1 + \frac{2}{qx}\right) e^{-qx}. \text{ Hence, } y = -\alpha - \frac{2C}{x}$$

is a solution, and if we substitute it in the proposed equation to determine the supernumerary constant, we find $C = -\frac{\alpha}{q}$. The necessity of adding a constant in finding y arises from the disappearance of the factor $d-1$ from the auxiliary equation.

$$\text{Let } n = 2, \quad y = \left(x^3 \frac{d}{dx}\right)^2 \frac{1}{x} (x^{-2} f dx)^2 x (\alpha + \beta e^{-qx})$$

$$= \left(x^3 \frac{d}{dx}\right)^2 \left(\frac{\alpha}{2x^3} + \frac{\beta}{q^2 x^4} e^{-qx} - \frac{C}{x^4} + \frac{C'}{x^3}\right)$$

$$= \alpha + \frac{6C'}{x} - \frac{12C}{x^2} + \beta \left(1 + \frac{6}{qx} + \frac{12}{q^2 x^2}\right) e^{-qx}.$$

Since $y = \alpha + \frac{6C'}{x} - \frac{12C}{x^2}$ is a solution, the supernumerary constants C and C' must be determined by substituting this value in the proposed equation; and the exact solution will be found to be

$$y = \alpha \left(1 - \frac{6}{qx} + \frac{12}{q^2 x^2}\right) + \beta \left(1 + \frac{6}{qx} + \frac{12}{q^2 x^2}\right) e^{-qx}.$$

$$7. \quad (1-x^2) x^2 \frac{d^2 y}{dx^2} - (2m+x^2) x \frac{dy}{dx} + \{m(m+1) + q^2 x^2\} y = 0.$$

$$\text{or } y - \frac{(d-2)^2 - q^2}{(d-m)(d-m-1)} e^2 y = 0.$$

Making $m = 0$, let the auxiliary equation be

$$(1-x^2) \frac{d^2 v}{dx^2} - x \frac{dv}{dx} + q^2 v = 0,$$

where $v = \alpha \cos (q \cos^{-1} x) + \beta \sin (q \cos^{-1} x)$, (Art. 69)

$$\text{or } v - \frac{(d-2)^2 - q^2}{d(d-1)} e^2 v = 0;$$

$$\therefore f(d) = P_2 \frac{d(d-1)}{(d-m)(d-m-1)} = d(d-1) \dots (d-m+1);$$

$$\therefore y = d(d-1)(d-2) \dots (d-m+1) v$$

$$= x^m \left(\frac{d}{dx} \right)^m \{ \alpha \cos (q \cos^{-1} x) + \beta \sin (q \cos^{-1} x) \}.$$

If q^2 have a negative sign before it, the value of v is $\alpha e^{q \cos^{-1} x} + \beta e^{-q \cos^{-1} x}$.

75. Thirdly, suppose that the transformed equation can be put under the form

$$y + a_1 f(d) e' y + a_2 f(d) f(d-1) e^2 y \\ + a_3 f(d) f(d-1) f(d-2) e^3 y + \&c. = T;$$

then since $f(d-1) e^2 y = e' f(d) e' y$;

$$\therefore f(d) f(d-1) e^2 y = f(d) e' f(d) e' y = \{f(d) e'\}^2 y,$$

and generally

$$f(d) f(d-1) f(d-2) \dots f(d-n+1) e^n y = \{f(d) e'\}^n y,$$

consequently the proposed equation becomes, if we denote the symbol of operation $f(d) e'$ by ρ ,

$$\{1 + a_1 \rho + a_2 \rho^2 + \dots + a_m \rho^m\} y = T;$$

$$\therefore y = (1 + a_1 \rho + a_2 \rho^2 + \dots + a_m \rho^m)^{-1} T$$

$$= \left(\frac{N_1}{1 - q_1 \rho} + \frac{N_2}{1 - q_2 \rho} + \dots + \frac{N_m}{1 - q_m \rho} \right) T,$$

resolving the function of ρ with which T is affected into its partial fractions,

$$\text{or } y = N_1 u_1 + N_2 u_2 + \dots + N_m u_m, \text{ where } u_1, u_2, \&c.$$

are determined from the equations,

$$u_1 - q_1 f(d) e^u u_1 = T_1, \quad u_2 - q_2 f(d) e^u u_2 = T_1, \&c.$$

76. If $f(d) = d^{-1}$, the equation just treated of is the linear equation with constant coefficients :

$$x^n \frac{d^n y}{dx^n} + p_1 x x^{n-1} \frac{d^{n-1} y}{dx^{n-1}} + p_2 x^2 x^{n-2} \frac{d^{n-2} y}{dx^{n-2}} + \&c. = x^n X;$$

for this leads to

$$P_n(d) y + p_1 P_{n-1}(d-1) e^u y + p_2 P_{n-2}(d-2) e^u y + \&c. = T;$$

$$\text{or } y + \frac{p_1}{d} e^u y + \frac{p_2}{d(d-1)} e^u y + \&c. = \frac{T}{P_n(d)};$$

which is of the assumed form. For all other values of $f(d)$ the equation has variable coefficients. If the equation be

$$x^n \frac{d^n y}{dx^n} + p_1 x^{n-1} \frac{d^{n-1} y}{dx^{n-1}} + p_2 x^{n-2} \frac{d^{n-2} y}{dx^{n-2}} + \&c. = X;$$

$$f(d) = \frac{1}{d-n+1},$$

which leads to a linear equation of the first order for the determination of $u_1, u_2, \&c.$

$$x^n \frac{d}{dx} \left(\frac{u_1 - T_1}{x^{n-1}} \right) = q_1 x u_1; \text{ if } T_1 = 0, \text{ this gives } u_1 = C x^{n-1} e^{q_1 x}.$$

If $f(d) = a + b(d-r)^{-1}$, the particular values of y are determined from linear equations of the first order,

$$\frac{d}{dx} \{u_1 (1 - q_1 a x) x^{-r} - T_1 x^{-r}\} = q_1 b u_1 x^{-r}.$$

$$\begin{aligned} \text{Ex. } (1 + ax + bx^2) x^2 \frac{d^2 y}{dx^2} - \{2r + (r-2) ax - 4bx^2\} x \frac{dy}{dx} \\ + \{r(r+1) - rax + 2bx^2\} y = X. \end{aligned}$$

Changing the independent variable into $t = \log x$, this becomes

$$(1 + ae^t + be^{2t}) d(d-1)y - \{2r + (r-2)ae^t - 4be^{2t}\} dy \\ + \{r(r+1) - rae^t + 2be^{2t}\} y = T,$$

$$\text{or } (d-r)(d-r-1)y + ae^t(d+1)(d+1-r-1)y \\ + be^{2t}(d+2)(d+2-1)y = T,$$

$$\text{or } (d-r)(d-r-1)y + ad(d-r-1)e^ty + bd(d-1)e^{2t}y = T,$$

$$\text{or } y + a \frac{d}{d-r} e^ty + b \frac{d}{d-r} \cdot \frac{d-1}{d-r-1} e^{2t}y = T_1,$$

$$\text{or } (1 + a\rho + b\rho^2)y = T_1, \text{ putting } \frac{d}{d-r} e^t = \rho,$$

$$\text{or } (1 + a\rho + b\rho^2)y = (1 - q_1\rho)(1 - q_2\rho)y = T_1;$$

$$\therefore y = \left(\frac{N_1}{1 - q_1\rho} + \frac{N_2}{1 - q_2\rho} \right) T_1 = N_1u_1 + N_2u_2,$$

where u_1 is given by the equation

$$(1 - q_1\rho)u_1 = T_1, \text{ or } u_1 - q_1 \frac{d}{d-r} e^tu_1 = T_1,$$

$$\text{or } (d-r)(u_1 - T_1) = q_1 d e^tu_1,$$

$$\text{or } x^{r+1} \frac{d}{dx} \left(\frac{u_1 - T_1}{x^r} \right) = q_1 x \frac{d}{dx} (xu_1).$$

Further applications of this method may be seen in the *Philosophical Transactions*, where Mr Boole has treated this subject with great generality. As a concluding application of the method by changing the independent variable, let us solve the equation

$$(x^2 - 4) \frac{d^2y}{dx^2} + 2(x+1) \frac{dy}{dx} - n(n+1)y = X.$$

Taking z for independent variable where $z + \frac{1}{z} = x$, we find

$$(z-1)z^2 \frac{d^2y}{dz^2} + 2z^2 \frac{dy}{dz} - n(n+1)(z-1)y = Z.$$

Now assume $z = e^t$, then

$$(e^t - 1) d(d-1)y + 2e^t dy - n(n+1)(e^t - 1)y = Z,$$

$$\text{or } e^t(d+n+1)(d-n)y - (d-n-1)(d+n)y = Z,$$

$$\text{or } (d+n)(d-n-1)(e^t y - y) = Z;$$

$$\therefore (e^t - 1)y = \frac{1}{2n+1} \left(\frac{1}{d-n-1} - \frac{1}{d+n} \right) Z,$$

$$\text{or } (2n+1)(z-1)y = z^{n+1} \int dz z^{-n-2} Z - z^{-n} \int dz z^{n-1} Z.$$

If $Z = 0$, we have

$$y = \frac{cz^{n+1} + c'z^{-n}}{z-1};$$

and if for z we substitute its value $\frac{1}{2}(x + \sqrt{x^2 - 4})$, we shall find

$$y = cu + c' \sqrt{\frac{x+2}{x-2}} \left\{ u + 2(x-2) \frac{du}{dx} \right\},$$

$$\text{where } u = x^n + x^{n-1} - (n-1)x^{n-2} - (n-2)x^{n-3} + \&c.$$

the general term of the series for u being

$$(-1)^r \frac{(n-r-1)(n-r-2) \dots (n-2r+1)}{1.2.3 \dots r} (n-2r+nx-rx) x^{n-2r-1}.$$

Simultaneous Equations.

In these equations, which sometimes occur in the higher parts of Dynamics, instead of one equation between x , y , and the differential coefficients of y with respect to x , being given to determine the relation between x and y ; we have two equations containing x , y , and t (of which x and y are considered as functions) and the differential coefficients of x and y relative to t , to find that relation.

77. To integrate the simultaneous equations of the first order,

$$ax + by + \frac{dx}{dt} = T, \quad a'x + b'y + \frac{dy}{dt} = T',$$

T and T' denoting functions of t . Multiplying the latter by an indeterminate quantity m , and adding it to the former, we get

$$ax + by + m(a'x + b'y) + \frac{d}{dt}(x + my) = T + mT',$$

$$\text{or } \frac{d}{dt}(x + my) + (a + ma')(x + \frac{b + mb'}{a + ma'}y) = T + mT'.$$

Let $\frac{b + mb'}{a + ma'} = m$, which will give two values of m , m_1 and m_2 ; then the equation, under this condition, becomes a linear equation of the first order; and we obtain by integration

$$(x + my) e^{(a+ma')t} = \int dt (T + mT') e^{(a+ma')t};$$

and by substituting successively the two values of m , we obtain two primitive equations each containing an arbitrary constant, which will furnish values of x and y in terms of t , and the relation between x and y , if t be eliminated. If the two values of m are equal, we shall obtain only one equation between x , y , and t ; but if this can be solved with respect to x or y , and we substitute the value so found in one of the given equations, we shall obtain a second relation either between x and t , or between y and t ; and then t may be eliminated as before.

$$\text{Ex. 1. } 5x - 2y + \frac{dx}{dt} = e^t, \quad 6y - x + \frac{dy}{dt} = e^{2t};$$

$$\therefore (5 - m)x + (-2 + 6m)y + \frac{d}{dt}(x + my) = e^t + me^{2t},$$

$$\text{let } \frac{-2 + 6m}{5 - m} = m, \text{ or } m^2 + m - 2 = 0, \text{ or } m = 1, \text{ or } -2;$$

$$\therefore \frac{d}{dt}(x + y) + 4(x + y) = e^t + e^{2t};$$

$$\therefore x + y = \frac{e^t}{5} + \frac{e^{2t}}{6} + Ce^{-4t};$$

$$\text{similarly } x - 2y = \frac{e^t}{8} - \frac{2}{9}e^{2t} + C_1e^{-t},$$

which determine x and y in terms of t .

Ex. 2. $\frac{dx}{dt} + 5x + y = e^t, \quad \frac{dy}{dt} + 3y - x = e^{2t};$

$$\therefore \frac{d}{dt}(x + my) + (5 - m)x + (1 + 3m)y = e^t + me^{2t};$$

$$\therefore \frac{1 + 3m}{5 - m} = m, \quad \text{or } 1 - 2m + m^2 = (1 - m)^2 = 0.$$

Hence the values of m are each $= 1$, and integrating, we find

$$x + y = C_1 e^{-4t} + \frac{1}{5} e^t + \frac{1}{6} e^{2t}.$$

By means of this, eliminate y from the first equation, and we get

$$\frac{dx}{dt} + 4x = \frac{4}{5} e^t - \frac{e^{2t}}{6} - C_1 e^{-4t};$$

$$\therefore x = \frac{4}{25} e^t - \frac{1}{36} e^{2t} - C_1 t e^{-4t} + C_2 e^{-4t},$$

$$\text{and } y = \frac{1}{25} e^t + \frac{7}{36} e^{2t} + C_1 (1 + t) e^{-4t} - C_2 e^{-4t}.$$

The more general form

$$ax + by + A \frac{dx}{dt} + B \frac{dy}{dt} = T, \quad a'x + b'y + A' \frac{dx}{dt} + B' \frac{dy}{dt} = T',$$

may evidently be reduced to the above by successively eliminating $\frac{dy}{dt}$, $\frac{dx}{dt}$.

78. To integrate the simultaneous equations of the second order,

$$ax + by + c + \frac{d^2x}{dt^2} = 0, \quad a'x + b'y + c' + \frac{d^2y}{dt^2} = 0.$$

Multiplying the latter by an indeterminate quantity m , and adding it to the former, we get

$$(a + ma')x + (b + mb')y + c + mc' + \frac{d^2}{dt^2}(x + my) = 0,$$

$$\text{or } \frac{d^2}{dt^2}(x + my + c_1) + (a + ma')(x + my + c_1) = 0 \quad (1),$$

$$\text{if } \frac{b + mb'}{a + ma'} = m \quad (2), \quad \frac{c + mc'}{a + ma'} = c_1;$$

therefore, integrating equation (1), and substituting successively the two values of m given by equation (2), we obtain the two required primitives; or if the values of m be equal, we must proceed as in the former case.

$$\text{Ex. 1. } \frac{d^2x}{dt^2} - (3x + 4y - 3) = 0, \quad \frac{d^2y}{dt^2} + (x + y + 5) = 0;$$

$$\therefore \frac{d^2}{dt^2}(x + my) + (m - 3) \left(x + \frac{m - 4}{m - 3}y + \frac{3 + 5m}{m - 3} \right) = 0.$$

$$\text{Let } \frac{m - 4}{m - 3} = m, \quad \text{or } m^2 - 4m + 4 = (m - 2)^2 = 0;$$

$$\therefore \frac{d^2}{dt^2}(x + 2y) - (x + 2y - 13) = 0;$$

$$\therefore x + 2y - 13 = ce^t + c'e^{-t},$$

and eliminating x from the latter of the given equations, we find

$$\frac{d^2y}{dt^2} - y + 18 + ce^t + c'e^{-t} = 0;$$

$$\therefore y = 18 - \frac{c}{2} \left(t - \frac{1}{2} \right) e^t + \frac{c'}{2} t e^{-t} + ae^t + a'e^{-t},$$

$$\text{and } x = -23 + c \left(t + \frac{1}{2} \right) e^t - c' (t - 1) e^{-t} - 2ae^t - 2a'e^{-t}.$$

$$\text{Ex. 2. } \frac{d^2x}{dt^2} + a \frac{dx}{dt} = 0, \quad \frac{d^2y}{dt^2} + a \frac{dy}{dt} + b = 0.$$

79. If we have three variables x, y, z , which are functions of t , and if

$$\frac{dx}{dt} + ax + by + cz = T,$$

$$\frac{dy}{dt} + a_1x + b_1y + c_1z = T_1,$$

$$\frac{dz}{dt} + a_2x + b_2y + c_2z = T_2,$$

then if we multiply the second and third by indeterminate constants m, m' , and add them to the first, and assume

$$a + ma_1 + m'a_2 = s,$$

$$b + mb_1 + m'b_2 = ms, \quad c + mc_1 + m'c_2 = m's,$$

we get

$$\frac{d}{dt}(x + my + m'z) + s(x + my + m'z) = T + mT_1 + m'T_2.$$

But the three preceding equations give

$$s - a = ma_1 + m'a_2, \quad m(s - b_1) = m'b_2 + b, \quad m'(s - c_2) = mc_1 + c;$$

and if values of m and m' be obtained from the two latter and substituted in the former, we get to determine s the cubic equation

$$(s-a)(s-b_1)(s-c_2) - b_2c_1(s-a) - a_2c(s-b_1) - ba_1(s-c_2) - a_1b_2c - a_2bc_1 = 0.$$

Hence if $m_1, m'_1; m_2, m'_2; m_3, m'_3$; be the values of m, m' , corresponding to the three roots of this cubic, we have by solving the linear equations,

$$x + m_1y + m'_1z = F_1(t),$$

$$x + m_2y + m'_2z = F_2(t),$$

$$x + m_3y + m'_3z = F_3(t),$$

from which equations x, y, z may be found in terms of t . It may be observed that if $b_2 = c_1, c = a_2$, and $a_1 = b$, so that the cubic equation is

$$(s-a)(s-b_1)(s-c_2) - c_1^2(s-a) - a_2^2(s-b_1) - b^2(s-c_2) - 2c_1a_2b = 0,$$

then all its roots are real (*Theory of Algebr. Eq. Art. 58*).

80. This method readily leads to Jacobi's solution of the equation

$$(A + A'x + A''y) \left(x \frac{dy}{dx} - y \right) - (B + B'x + B''y) \frac{dy}{dx} + (C + C'x + C''y) = 0,$$

which, upon introducing the independent variable t , becomes

$$\begin{aligned} & \{(A + A'x + A''y)x - (B + B'x + B''y)\} \frac{dy}{dt} \\ & = \{(A + A'x + A''y)y - (C + C'x + C''y)\} \frac{dx}{dt}, \end{aligned}$$

and resolves itself into

$$\frac{dx}{dt} - (A + A'x + A''y)x + (B + B'x + B''y) = 0,$$

$$\frac{dy}{dt} - (A + A'x + A''y)y + (C + C'x + C''y) = 0;$$

or {putting $\frac{dz}{dt} = -(A + A'x + A''y)z$ into the system,

$$\frac{dz}{dt} + Az + A'xz + A''yz = 0,$$

$$\frac{d(xz)}{dt} + Bz + B'xz + B''yz = 0,$$

$$\frac{d(yz)}{dt} + Cz + C'xz + C''yz = 0.$$

Multiply the second and third by m and m' respectively, and add them to the first, and assume

$$A + mB + m'C = s, \quad A' + mB' + m'C' = ms, \quad A'' + mB'' + m'C'' = m's \quad (1);$$

$$\text{then } \frac{d}{dt} (z + mxz + m'yz) + s(z + mxz + m'yz) = 0;$$

$$\text{which gives } z(1 + mx + m'y)e^{\int s dt} = C, \text{ or } zue^{\int s dt} = C.$$

But equations (1) lead successively to

$$s - A = mB + m'C, \quad m(s - B') = A' + m'C', \quad m'(s - C'') = A'' + mB'';$$

the two latter of which determine m and m' from s by the equations

$$\left. \begin{aligned} m \{(s - B') (s - C'') - B'' C'\} &= A' (s - C'') + C' A'' \\ m' \{(s - C'') (s - B') - B'' C'\} &= A'' (s - B') + A' B'' \end{aligned} \right\} \quad (2),$$

and the values thence obtained of m and m' substituted in the former give for finding s the cubic equation

$$\begin{aligned} (s - A) (s - B') (s - C'') - B'' C' (s - A) - A'' C (s - B') \\ - A' B (s - C'') + A'' B C' + A' B'' C = 0. \end{aligned}$$

Let the three values of s in this equation be s_1, s_2, s_3 ; and let the corresponding values of m and m' , be m_1 and m'_1, m_2 and m'_2, m_3 and m'_3 ; then corresponding to the three roots of the cubic we get the three integrals.

$$zu_1e^{s_1t} = C_1, \quad zu_2e^{s_2t} = C_2, \quad zu_3e^{s_3t} = C_3,$$

and if we raise these equations respectively to the powers $s_2 - s_3$, $s_3 - s_1$, $s_1 - s_2$, and take their product, z and e^t are eliminated, and we find the required solution

$$u_1^{s_2-s_3} \times u_2^{s_3-s_1} \times u_3^{s_1-s_2} = C;$$

$$\text{or } (1 + m_1x + m'_1y)^{s_2-s_3} \times (1 + m_2x + m'_2y)^{s_3-s_1} \\ \times (1 + m_3x + m'_3y)^{s_1-s_2} = C,$$

where m_1 , m'_1 are given in terms of s_1 , m_2 , m'_2 in terms of s_2 , and m_3 , m'_3 in terms of s_3 , by equations (2). This includes as a particular case the equation (Ex. 7, Art. 18), in which Euler separated the variables,

$$(a + bx + cx^2 + y) \frac{dy}{dx} - (n + cx) y = 0,$$

$$\text{or } cx \left(x \frac{dy}{dx} - y \right) + (a + bx + y) \frac{dy}{dx} - ny = 0.$$

81. When the proposed equations are linear with constant coefficients, we may separate the symbols of operation from those of quantity and then obtain by the ordinary processes of elimination an equation containing only one of the unknown quantities: for as the symbols of operation here employed combine according to the same laws as ordinary algebraical quantities, they may be treated precisely as if they were symbols of quantity. Thus the equations of Art. 77 (suppressing the dt in $\frac{d}{dt}$) may be written

$$(d + a)x + by = T, \quad (d + b')y + a'x = T';$$

the latter gives $y = \frac{T' - a'x}{d + b'}$; and substituting this value in the former we get

$$(d + a)(d + b')x - a'bx = (d + b')T - bT',$$

an ordinary linear equation with constant coefficients, from which the value of x may be obtained, containing two arbitrary constants; and thence the value of y by the equation

$$y = \frac{1}{b} T - \frac{1}{b} (d + a)x.$$

$$\text{Ex. } \frac{dx}{dt} + 2cx - y = t^2, \quad \frac{dy}{dt} - 2cy + 5c^2x = e^{mt};$$

the latter gives for y the value $y = \frac{e^{mt} - 5c^2x}{d - 2c}$,

and substituting this in the former we get

$$(d^2 + c^2)x = e^{mt} + 2(t - ct^2).$$

$$\therefore x = \frac{e^{mt}}{m^2 + c^2} + \frac{2}{c^2}(t - ct^2) + \frac{4}{c^3} + \beta \cos(ct + \alpha);$$

$$\text{and } y = (d + 2c)x - t^2$$

$$= \frac{m + 2c}{m^2 + c^2}e^{mt} - 5t^2 + \frac{10}{c^2} - \beta c \sin(ct + \alpha) + 2\beta c \cos(ct + \alpha).$$

82. Again, suppose the equations of Art. 78 to be

$$(d^2 + ad + b)x = (md + n)y + T,$$

$$(d^2 - ad + b')y = (m'd + n')x + T';$$

obtaining a value of y from the latter, substituting it in the former and reducing, we find

$$\begin{aligned} \{d^4 - (a^2 - b - b' + mm')d^2 + (ab' - ab - mn' - m'n)d + bb' - nn'\}x \\ = (d^2 - ad + b')T + (md + n)T'; \end{aligned}$$

from which the value of x may be obtained; and if it be substituted in one of the proposed equations, we may deduce the value of y .

$$\text{Ex. 1. } (d - 1)^2 x = 2dy + \cos t$$

$$(d^2 + 2d + 6)y = -(d + 5)x + \sin t;$$

eliminating y , we find

$$(d^4 + 5d^2 + 6)x = 7 \cos t - 2 \sin t,$$

$$\therefore x = \frac{7}{2} \cos t - \sin t + c \cos(t\sqrt{2} + \alpha) + c' \cos(t\sqrt{3} + \beta);$$

and the value of y may be now obtained from the former of the proposed equations.

Ex. 2. $\{d^2 + \left(n + \frac{1}{n}\right)d + n^2\} x = dy + T,$

$$\left\{d^2 - \left(n + \frac{1}{n}\right)d + \frac{1}{n^2}\right\} y = \left\{m^2 d - \left(n + \frac{1}{n}\right)\left(n^2 - \frac{1}{n^2}\right)\right\} x,$$

eliminating y , we find

$$\{d^4 - (2 + m^2)d^2 + 1\} x = \left\{d^2 - \left(n + \frac{1}{n}\right)d + \frac{1}{n^2}\right\} T,$$

$$\text{or } (d^2 - md - 1)(d^2 + md - 1)x = \left\{d^2 - \left(n + \frac{1}{n}\right)d + \frac{1}{n^2}\right\} T,$$

which gives x ; and then y can be obtained from the former of the given equations. In these examples d is used for $\frac{d}{dt}$.

Solutions expressed by Definite Integrals.

83. Sometimes the complete integral of a differential equation may be expressed by a definite integral, as in the following instances.

Ex. 1. $\frac{d^{n-1}y}{dx^{n-1}} = xy + m.$

Let $v_1 = \alpha_1 \int_0^\infty dt e^{a_1 t x} \cdot e^{-\frac{t^n}{n}},$ α_1 being a constant quantity ;

$$\text{then } \frac{d^{n-1}}{dx^{n-1}} v_1 = \alpha_1^n \int_0^\infty dt t^{n-1} e^{a_1 t x} e^{-\frac{t^n}{n}} = \alpha_1^n (1 + x v_1);$$

change α_1 into $\alpha_2, \alpha_3,$ &c., and let $v_2, v_3,$ &c. be the corresponding values of $v_1,$

$$\therefore \frac{d^{n-1}}{dx^{n-1}} v_2 = \alpha_2^n (1 + x v_2),$$

.....

$$\frac{d^{n-1}}{dx^{n-1}} v_n = \alpha_n^n (1 + x v_n).$$

Now suppose $\alpha_1, \alpha_2, \dots \alpha_n$ to be the n^{th} roots of unity; then multiplying each of these equations by arbitrary constants

$c_1, c_2, \&c.$ such that $c_1 + c_2 + \dots + c_n = m$, and adding them together, we find

$$\frac{d^{n-1}}{dx^{n-1}} (c_1 v_1 + c_2 v_2 + \dots + c_n v_n) = m + x (c_1 v_1 + c_2 v_2 + \dots + c_n v_n).$$

Hence $y = c_1 v_1 + c_2 v_2 + \dots + c_n v_n$ is the complete integral of the proposed equation, containing only $n-1$ arbitrary constants by reason of the equation of condition; or it may be written

$$y = \int_0^\infty dt e^{-\frac{t^n}{n}} \{ c e^{tx} + c_1 \alpha e^{\alpha t x} + c_2 \alpha^2 e^{\alpha^2 t x} + \dots + c_{n-1} \alpha^{n-1} e^{\alpha^{n-1} t x} \},$$

α being a primitive root of $k^n - 1 = 0$.

Let $m = 0, n = 3$, then $\frac{d^2 y}{dx^2} = xy$, the solution of which is

$$y = \int_0^\infty dt e^{-\frac{1}{3} t^3 - \frac{1}{2} x t} \left\{ B e^{\frac{3 t x}{2}} + \frac{1}{2} (B + A \sqrt{3}) \cos \frac{\sqrt{3} t x}{2} - \frac{1}{2} (A + B \sqrt{3}) \sin \frac{\sqrt{3} t x}{2} \right\}.$$

Let $m = 0, n = 2$, then $\frac{dy}{dx} = xy$, and

$$\begin{aligned} y &= C \int_0^\infty dt e^{-\frac{t^2}{2}} (e^{xt} + e^{-xt}) = \frac{2C}{\sqrt{-1}} \int_0^\infty dz e^{\frac{z^2}{2}} \cos xz, \text{ putting } t = z \sqrt{-1}, \\ &= \frac{2C}{\sqrt{-1}} \cdot \frac{\sqrt{\pi} e^{\frac{x^2}{2}}}{\sqrt{-2}} = C \sqrt{2\pi} e^{\frac{x^2}{2}}, \quad (\text{Integ. Cal. Art. 112}). \end{aligned}$$

which agrees with the result obtained by direct integration.

Ex. 2. If the more general form were proposed,

$$\frac{d^{n-1} y}{dx^{n-1}} = (\alpha + \beta x) y,$$

$$\text{assume } \alpha + \beta x = \beta^{\frac{n-1}{n}} t; \quad \therefore \frac{dy}{dx} = \frac{dy}{dt} \frac{dt}{dx} = \beta^{\frac{1}{n}} \frac{dy}{dt},$$

$$\text{and } \frac{d^{n-1} y}{dx^{n-1}} = \beta^{\frac{n-1}{n}} \frac{d^{n-1} y}{dt^{n-1}} = \beta^{\frac{n-1}{n}} t y; \quad \therefore \frac{d^{n-1} y}{dt^{n-1}} = t y;$$

and to this form may the still more general case

$$\frac{d^{n-1}y}{dx^{n-1}} = axy + bx + cy + e$$

be reduced. (The above solution is taken from Crelle's Journal, Vol. x.)

Approximate Solutions of Differential Equations.

84. When all the known methods of integrating a proposed differential equation fail, we must endeavour to resolve it approximately, that is, to obtain from it the value of y in terms of x , in the form of a series. The first mode which presents itself of effecting this, is to assume for y a series arranged according to powers of x , with both its coefficients and exponents undetermined; for in most cases it happens that the exponents do not follow the progression of the natural numbers, and that particular artifices are requisite for discovering their law. When the form of this series is known, we may determine its coefficients by substituting it and its differential coefficients, for y , $\frac{dy}{dx}$, &c. in the proposed equation. The following application of the method to Riccati's Equation will give an idea of the mode of obtaining both the exponents and the coefficients.

Ex. 1. $\frac{dy}{dx} + by^2 - ax^n = 0,$

let $by = \frac{1}{z} \frac{dz}{dx}$; $\therefore \frac{d^2z}{dx^2} - abzx^n = 0$, or putting c for $-ab$,

$$\frac{d^2z}{dx^2} + czx^n = 0.$$

Assume $z = x^a (A + Bx^\beta + Cx^{2\beta} + \&c.)$; then

$$\left. \begin{aligned} &+ (\alpha + \beta) (\alpha + \beta - 1) Bx^{a+\beta-2} + (\alpha + 2\beta) (\alpha + 2\beta - 1) Cx^{a+2\beta-2} + \&c. \\ &+ cAx^{a+n} \end{aligned} \right\} = 0.$$

Hence β must equal $n+2$; and then to determine α , A , B , &c., we have

$$\begin{aligned}\alpha(\alpha-1)A &= 0, \\ (\alpha+n+2)(\alpha+n+1)B + cA &= 0, \\ (\alpha+2n+4)(\alpha+2n+3)C + cB &= 0, \\ (\alpha+3n+6)(\alpha+3n+5)D + cC &= 0, \\ &\dots\dots\dots\end{aligned}$$

Hence α may be either zero or unity, and A remains undetermined; calling therefore A and A' the two values of A corresponding to $\alpha=0$, $\alpha=1$, we get

$$\begin{aligned}z &= A \left\{ 1 - \frac{cx^{n+2}}{(n+1)(n+2)} + \frac{c^2x^{2n+4}}{(n+1)(n+2)(2n+3)(2n+4)} - \&c. \right\} \\ &+ A'x \left\{ 1 - \frac{cx^{n+2}}{(n+2)(n+3)} + \frac{c^2x^{2n+4}}{(n+2)(n+3)(2n+4)(2n+5)} - \&c. \right\}\end{aligned}$$

and substituting this value of z in the expression $y = \frac{1}{bz} \frac{dz}{dx}$, we shall obtain the value of y , involving only one arbitrary constant $\frac{A}{A'}$.

As the terms of the above series have divisors of the forms $(n+2)i \mp 1$, where i is an integer; if n be such that

$$(n+2)i \mp 1 = 0,$$

one or other of the series will be illusory, and we shall only obtain a particular value of z ; and if $n+2=0$, both series become infinite, but in that case the equation may be exactly integrated by Art. 16.

$$\text{Ex. 2. } (x^2-4) \frac{d^2y}{dx^2} + x \frac{dy}{dx} - n^2y = 0.$$

$$\text{Assume } y = x^\alpha (a_0 + a_1x^\beta + a_2x^{2\beta} + a_3x^{3\beta} + \&c.),$$

$$\begin{aligned}\therefore y &= (\alpha^2 - n^2) a_0 x^\alpha + \{(\alpha+\beta)^2 - n^2\} a_1 x^{\alpha+\beta} - 4\alpha(\alpha-1) a_0 x^{\alpha-2} + \&c. \\ &+ \{(\alpha+r\beta)^2 - n^2\} a_r x^{\alpha+r\beta} - 4(\alpha-2r+2)(\alpha-2r+1) a_{r-1} x^{\alpha-2r} + \&c.,\end{aligned}$$

the first term shews that α may be either $+n$ or $-n$, α_0 being the arbitrary constant; the next terms shew that $\beta = -2$, i. e. the series is a descending one; and that the coefficients are derived from one another by the law (taking $\alpha = n$),

$$\alpha_r = -\frac{(n-2r+2)(n-2r+1)}{r(n-r)} \alpha_{r-1};$$

$$\begin{aligned} \therefore y &= \alpha_0 \{x^n - nx^{n-2} + \frac{n(n-3)}{1 \cdot 2} x^{n-4} - \&c.\} \\ &+ \alpha'_0 \{x^{-n} + nx^{-n-2} + \frac{n(n+3)}{1 \cdot 2} x^{-n-4} + \&c.\}. \end{aligned}$$

As the solution of the proposed equation is (Art. 48)

$$y = \alpha_0 \cos \left(n \cos^{-1} \frac{x}{2} \right) + \alpha'_0 \sin \left(n \cos^{-1} \frac{x}{2} \right),$$

if we make $x = 2 \cos \theta$, we get the well-known result,

$$\cos n\theta = (2 \cos \theta)^n - n (2 \cos \theta)^{n-2} + \frac{n(n-3)}{1 \cdot 2} (2 \cos \theta)^{n-4} - \&c.$$

85. The transformation of Art. 71 may be employed in the case of linear equations to find the solution in the form of a series: for suppose the transformed equation to consist of three terms

$$f_0(d)y + f_1(d)dy + f_2(d)d^2y = 0.$$

$$\text{Let } y = u_m x^m + u_{m+1} x^{m+1} + \dots + u_{r-2} x^{r-2} + u_{r-1} x^{r-1} + u_r x^r + \&c.,$$

where u_r in the general term $u_r x^r$ is an invariable function of the index r , and x^m is the lowest power of x that can enter into the development of y . Now substitute this value of y (having first replaced x by e^t) in the proposed equation: then since

$$f(d)e^{rt} = f(r)e^{rt},$$

the first member will become

$$\begin{aligned} f_0(m) u_m e^{mt} + \{f_0(m+1) u_{m+1} + f_1(m+1) u_m\} e^{(m+1)t} + \dots \\ + \{f_0(r) u_r + f_1(r) u_{r-1} + f_2(r) u_{r-2}\} e^{rt} + \&c. \end{aligned}$$

which will vanish, provided

$$f_0(m) u_m = 0, \quad f_0(m+1) u_{m+1} + f_1(m+1) u_m = 0,$$

and if for every value of r from $m+2$ upwards, we have

$$f_0(r) u_r + f_1(r) u_{r-1} + f_2(r) u_{r-2} = 0.$$

The first condition $f_0(m) = 0$, determines the values of m , u_m being an arbitrary constant: the second condition gives u_{m+1} in terms of u_m ; and the third determines any coefficient u_r from the values of the two preceding. The third condition comprises the other two when we make $r = m$, $r = m+1$, if we consider that as $u_m x^m$ is the lowest power of x that can enter, $u_{m-1} = 0$, $u_{m-2} = 0$. As many real unequal values as exist for m , so many ascending series there are for y : if m have equal or imaginary values, the method fails.

$$\text{Ex. 1.} \quad (1-x^2) \frac{d^2 y}{dx^2} - x \frac{dy}{dx} + n^2 y = 0,$$

$$\text{or } d(d-1)y + \{n^2 - (d-2)^2\} e^{2y} = 0.$$

Assume $y = u_m x^m + u_{m+1} x^{m+1} + \dots + u_r x^r + \&c.$; then the first member of the equation becomes

$$m(m-1) u_m e^{m^t} + m(m+1) u_{m+1} e^{(m+1)t} + \dots \\ + [r(r-1) u_r + \{n^2 - (r-2)^2\} u_{r-2}] e^{r^t} + \&c.$$

which will vanish, provided we have

$$m(m-1) u_m = 0, \quad u_{m+1} = 0;$$

$$\text{and } u_r = -\frac{n^2 - (r-2)^2}{r(r-1)} u_{r-2}$$

for all values of r from $m+2$ upwards. Hence since the second term vanishes, the 4th, 6th, &c. term vanishes; and taking m successively equal to 0, and to 1, we get the complete value of y ;

$$y = C \left(1 - \frac{n^2 x^2}{1.2} + \frac{n^2 (n^2 - 2^2)}{1.2.3.4} x^4 - \&c. \right) \\ + C' \left\{ x - \frac{n^2 - 1}{2.3} x^3 + \frac{(n^2 - 1)(n^2 - 3^2)}{2.3.4.5} x^5 - \&c. \right\},$$

which agrees with well-known results (Trig. Art. 134), as the solution of the equation is

$$y = C \cos (n \sin^{-1} x) + C' \sin (n \sin^{-1} x).$$

$$\text{Ex. 2. } (x^2 - 4) \frac{d^2 y}{dx^2} + 2(x+1) \frac{dy}{dx} - n(n+1)y = 0;$$

$$\therefore e^{2x} (d+n+1)(d-n)y + 2e^x dy - 4d(d-1)y = 0,$$

and dividing by e^{2x} in order to obtain y in a descending series, we get

$$(d+n+1)(d-n)y + 2e^{-x}dy - 4d(d-1)y = 0,$$

$$\text{or } (d-n)(d+n+1)y + 2(d+1)e^{-x}y - 4(d+1)(d+2)e^{-2x}y = 0.$$

Since the values of m are n and $-(n+1)$, we may assume

$$y = u_n e^{nx} + u_{n-1} e^{(n-1)x} + u_{n-2} e^{(n-2)x} + u_{n-3} e^{(n-3)x} + \&c.$$

then the first member of the equation becomes

$$-2nu_{n-1}e^{(n-1)x} - 2(2n-1)u_{n-2}e^{(n-2)x} - 3(2n-2)u_{n-3}e^{(n-3)x} + \&c.$$

$$+ 2nu_n e^{(n-1)x} + (2n-2)u_{n-1}e^{(n-2)x} + (2n-4)u_{n-2}e^{(n-3)x} + \&c.$$

$$-4(n-1)nu_n e^{(n-2)x} - 4(n-2)(n-1)u_{n-1}e^{(n-3)x} + \&c.$$

Hence, equating the coefficients of the powers of e^x to zero, we get

$$u_{n-1} = u_n, \quad u_{n-2} = -(n-1)u_n, \quad u_{n-3} = -(n-2)u_n, \quad \&c.;$$

and since $u_n = c$, an arbitrary constant,

$$\frac{y}{c} = x^n + x^{n-1} - (n-1)x^{n-2} - (n-2)x^{n-3} + \dots$$

$$+ \frac{(-1)^r}{[r]} (n-r-1)(n-r-2) \dots (n-2r+1) \{n-2r+(n-r)x\} x^{n-2r-1} + \&c.$$

Similarly the series corresponding to $m = -n-1$ may be found.

86. In the preceding instances we arrive immediately at a complete result; but it often happens that the solution we obtain by the method of indeterminate coefficients involves no arbitrary constant. To supply this defect, we must introduce, instead of

the arbitrary constant, a value of y corresponding to a given value of x ; that is, supposing these to be b and a , we must substitute in the given equation,

$$y = b + u, \quad x = a + t;$$

then determine u in a series all whose terms vanish when $t=0$; and replace u and t by their values $y-b$ and $x-a$; in this way it is evident the arbitrary constant will be involved implicitly; for, from the complete integral

$$f(x, y, C) = 0,$$

C may be expressed in terms of a and b .

Ex. $\frac{dy}{dx} + y = gx^m,$

$$\text{this becomes } \frac{du}{dx} + b + u = g(a+t)^m;$$

$$\text{assume } u = t^\alpha (A + Bt^\beta + Ct^\gamma + \&c.);$$

$$\therefore 0 = \alpha A t^{\alpha-1} + (\alpha + \beta) B t^{\alpha+\beta-1} + (\alpha + \gamma) C t^{\alpha+\gamma-1} + \&c.$$

$$+ b \quad + \quad A t^\alpha \quad + \quad B t^{\alpha+\beta} + \&c.$$

$$- g a^m \quad - \quad m g a^{m-1} t \quad - \frac{m(m-1)}{1 \cdot 2} g a^{m-2} t^2 - \&c.$$

$$\therefore \alpha = 1, \quad \beta = 1, \quad \gamma = 2, \quad \&c.$$

$$A = g a^m - b, \quad 2B = g m a^{m-1} - g a^m + b,$$

$$6C = g m(m-1) a^{m-2} - g m a^{m-1} + g a^m - b, \quad \&c.$$

87. The approximate solution of a differential equation may sometimes be obtained in the form of a continued fraction by assuming

$$y = \frac{Ax^\alpha}{1+} \frac{Bx^\beta}{1+} \frac{Cx^\gamma}{1+ \&c.}.$$

First, suppose x to be very small, and for y substitute Ax^α in the given equation; then, retaining only the terms of lowest dimensions in x , A and α become known by equating the coefficients and exponents of those terms. Next, write $y = \frac{Ax^\alpha}{1+z}$

in the proposed equation, and in the result put $z = Bx^\beta$, and determine B , β , as before, by supposing x to be very small; then in the transformed equation in z , put $z = \frac{Bx^\beta}{1+t}$; and so on for the rest.

Ex. $my + (1+x) \frac{dy}{dx} = 0.$

Let $y = Ax^a$; $\therefore (m+a) Ax^a + Aax^{a-1} = 0,$

or $Aax^{a-1} = 0$; $\therefore a = 0,$

and A remains undetermined. Next, put

$$y = \frac{A}{1+z}, \text{ and } z = Bx^\beta,$$

and we get successively $m(1+z) = (1+x) \frac{dz}{dx},$

$$m + Bx^\beta (m - \beta) = \beta Bx^{\beta-1}, \text{ or } m = \beta Bx^{\beta-1};$$

$$\therefore \beta = 1, \quad B = m;$$

similarly, putting $z = \frac{mx}{1+t}$, and $t = Cx^\gamma,$

we determine C and γ ; and thus we obtain

$$y = \frac{A}{1 + \frac{mx}{1 - \frac{m-1}{1 + \frac{x}{2} \cdot \frac{m+1}{3} \cdot \frac{x}{2} \cdot \frac{m-2}{3} \cdot \frac{x}{2} \cdot \frac{m+2}{5} \cdot \frac{x}{2} \cdot \frac{m-3}{5} \cdot \frac{x}{2} \dots}}}$$

Since the proposed equation when integrated gives

$$y = A(1+x)^{-m},$$

the above continued fraction is the development of $A(1+x)^{-m}.$

88. We may approximate to the integral of a differential equation by successive substitutions, in a manner similar to that employed by Newton for the solution of algebraical equations, as in the following instance.

Ex. $\frac{d^2y}{dx^2} + n^2y + \alpha y^2 + a = 0$, where α is a very small quantity.

We may assume

$$y = u + \alpha u_1 + \alpha^2 u_2 + \alpha^3 u_3 + \&c.$$

which gives

$$\begin{aligned} \frac{d^2 u}{dx^2} + n^2 u + a + \alpha \left(\frac{d^2 u_1}{dx^2} + n^2 u_1 + u^2 \right) \\ + \alpha^2 \left(\frac{d^2 u_2}{dx^2} + n^2 u_2 + 2u u_1 \right) \\ + \alpha^3 \left(\frac{d^2 u_3}{dx^2} + n^2 u_3 + 2u u_2 \right) + \&c. = 0. \end{aligned}$$

Hence, equating the coefficient of each power of α to zero, we get

$$\begin{aligned} \frac{d^2 u}{dx^2} + n^2 u + a &= 0, \\ \frac{d^2 u_1}{dx^2} + n^2 u_1 + u^2 &= 0, \\ \frac{d^2 u_2}{dx^2} + n^2 u_2 + 2u u_1 &= 0, \&c. \dots\dots\dots (3). \end{aligned}$$

The first give $u = -\frac{a}{n^2} + c \cos nx + c' \sin nx$; and this value substituted in the second reduces it to the form

$$\frac{d^2 u_1}{dx^2} + n^2 u_1 = X_1,$$

the integral of which by Art. (67) may be shewn to be

$$u_1 = \cos nx \left(c_1 - \frac{1}{n} \int dx X_1 \sin nx \right) + \sin nx \left(c'_1 + \frac{1}{n} \int dx X_1 \cos nx \right).$$

Similarly, these values of u and u_1 substituted in (3) reduce it to the form

$$\frac{d^2 u_2}{dx^2} + n^2 u_2 = X_2,$$

which may be in like manner integrated; and in this way the coefficients of the powers of α may be deduced one from the other by a uniform process.

89. We have seen (Art. 63) that the solution of

$$\frac{d^2 y}{dx^2} + n^2 y = A \cos (mx + \alpha) + B \cos (nx + \beta),$$

is $y = c_1 \cos nx + c_2 \sin nx$

$$+ \frac{A}{n^2 - m^2} \cos (mx + \alpha) + \frac{B}{2n} x \sin (nx + \beta).$$

Hence, if from the proposed equation we had to determine y approximately, we could not neglect the term

$$A \cos (mx + \alpha)$$

even when A is exceedingly small, provided m and n are nearly equal to one another; because in the value of y this term is divided by $n^2 - m^2$ which is very small. With respect to the last term in the value of y , we remark, that it is not periodical, but may increase indefinitely, as x increases. The equations which present themselves for solution in physical Astronomy are usually of the above form; and upon the peculiarities just noticed depend some of the most interesting results in that subject.

The following example supplies an omission at the end of Art. 83. When m is an integer we have obtained (Art. 74, Ex. 4) the solution of the equation

$$\frac{d^2 y}{dx^2} + n^2 y = \frac{m(m+1)}{x^2} y.$$

When m is a fraction, assume $y = x^{m+1} u$, then

$$\frac{d^2 u}{dx^2} + \frac{2m+2}{x} \frac{du}{dx} + n^2 u = 0,$$

the complete integral of which is

$$u = \beta \int_{-\infty}^x dt (t^2 - n^2)^m \cos (xt + \alpha),$$

α and β being the arbitrary constants; for this gives

$$\begin{aligned} \frac{du}{dx} &= -\beta \int_{-\infty}^x dt (t^2 - n^2)^m \sin (xt + \alpha) \cdot t \\ &= \frac{\beta x}{2m+2} \int_{-\infty}^x dt (t^2 - n^2)^{m+1} \cos (xt + \alpha), \end{aligned}$$

$$\frac{d^2 u}{dx^2} = -\beta \int_{-\infty}^x dt (t^2 - n^2)^m \cos (xt + \alpha) \cdot t^2$$

$$= -\beta \int_{-\infty}^x dt (t^2 - n^2)^m (t^2 - n^2 + n^2) \cos (xt + \alpha);$$

$$\therefore \frac{d^2 u}{dx^2} + n^2 u = -\beta \int_{-\infty}^x dt (t^2 - n^2)^{m+1} \cos (xt + \alpha) = -\frac{2m+2}{x} \frac{du}{dx}.$$

SECTION V.

ON DIFFERENTIAL EQUATIONS INVOLVING TWO OR MORE INDEPENDENT VARIABLES.

90. IF u be a function of any number of independent variables x, y, z, t , &c., and if $u + \delta u$ be the value of u when these variables simultaneously become $x + \delta x, y + \delta y, z + \delta z$, &c.,

$$\text{then } \delta u = \frac{du}{dx} \cdot \delta x + \frac{du}{dy} \cdot \delta y + \frac{du}{dz} \cdot \delta z + \dots$$

+ terms of two dimensions in $\delta x, \delta y$, &c.

If we now suppose all terms to be neglected of more than one dimension in $\delta x, \delta y$, &c., and denote the value of δu corresponding to that supposition by du , we have

$$du = \frac{du}{dx} \cdot \delta x + \frac{du}{dy} \cdot \delta y + \frac{du}{dz} \cdot \delta z + \dots;$$

according to which definition, it appears that du denotes that part of the increment of u as given by Taylor's Theorem, which involves only the first powers of the arbitrary increments of the variables on which it depends; hence, in conformity with this definition, $\delta x, \delta y, \delta z$, &c., must be represented by dx, dy, dz , &c.; for if $f(x) = x$, then

$$f(x + \delta x) = x + \delta x;$$

consequently that part of the increment of $f(x)$ which involves the first power of δx is, in this case, the whole of it δx , therefore $dx = \delta x$; and so on for the others;

$$\therefore du = \frac{du}{dx} \cdot dx + \frac{du}{dy} \cdot dy + \frac{du}{dz} \cdot dz + \&c.,$$

where dx, dy, dz , &c., are the arbitrary increments of the independent variables x, y, z , &c., and are entirely independent of

those variables; and du is that part of the corresponding increment of the dependent variable u which involves only simple dimensions of dx, dy, dz , &c.; and which approaches nearer and nearer to the value of the whole increment of u , the smaller dx, dy, dz , &c. are taken.

The quantities $\frac{du}{dx} \cdot dx, \frac{du}{dy} \cdot dy$, &c., are called the partial differentials of u with respect to x, y , &c. respectively; and du is called the total or complete differential of u , or the differential of u merely.

91. According to the above definition, the differential of du , or the second differential of u , will be, supposing it to involve only two independent variables,

$$\begin{aligned} d^2u &= \frac{d}{dx}(du) \cdot dx + \frac{d}{dy}(du) \cdot dy \\ &= \frac{d}{dx} \left(\frac{du}{dx} \cdot dx + \frac{du}{dy} \cdot dy \right) dx + \frac{d}{dy} \left(\frac{du}{dx} \cdot dx + \frac{du}{dy} \cdot dy \right) dy \\ &= \frac{d^2u}{dx^2} \cdot dx^2 + 2 \frac{d}{dx} \frac{d}{dy} u \cdot dx dy + \frac{d^2u}{dy^2} \cdot dy^2, \end{aligned}$$

because in this process dx and dy are independent of x and y ; being, in fact, the arbitrary increments of those variables, which might have been denoted by h and k , did not a due regard to the precision and symmetry of the notation require it otherwise. And in general, if u be a function of any number of independent variables x, y, z, t , &c., the n^{th} differential of u will be given by the formula, (where the symbols of operation are separated from those of quantity,)

$$d^n u = \left(dx \cdot \frac{d}{dx} + dy \cdot \frac{d}{dy} + dz \cdot \frac{d}{dz} + \&c. \right)^n u.$$

92. Hence we see that the differential of a function of several variables has no reference to one variable rather than to another; but in its formation, all the variables of which it is a function are supposed to undergo simultaneous unconnected alterations; and the value of the differential depends upon the

values of all those alterations. Whereas in forming a differential coefficient, one particular independent variable only is changed, the rest remaining unaffected; and a quantity is produced, wholly independent of the value of the alteration which that variable may have received.

It is evident that the rules for deriving differential coefficients suffice for finding the differentials of functions.

Having, for instance, formed all the partial differential coefficients of the first order, if we multiply each by the arbitrary increment or differential of the corresponding independent variable and take the sum, we shall obtain the first differential of the function; and similarly, the differentials of the second, and higher orders, may be formed by means of the formula at the end of the preceding Article.

Integration of Differential Functions of two or more Variables.

93. Total integration of a proposed differential is the finding a function whose differential is the quantity proposed. This operation is denoted by the symbol \int ; the process, like the former of total differentiation, having reference to all and not to one in particular of the variables. When an expression presented for total integration can be reduced to the form $f(v) dv$, its integral $= \int df(v)$, and can be obtained immediately.

Ex. 1. If $u = x^2 + y^2$,

$du = 2(xdx + ydy)$ is the differential of u ;

also $\int (xdx + ydy) = \frac{1}{2}(x^2 + y^2)$,

since the latter quantity differentiated produces the quantity under the symbol of total integration.

$$\begin{aligned} \text{Ex. 2. } du &= \frac{y(xdy - ydx)}{x^2 \sqrt{x^2 + y^2}} = \frac{\frac{y}{x}}{\sqrt{1 + \frac{y^2}{x^2}}} \cdot \frac{xdy - ydx}{x^2} \\ &= \frac{v}{\sqrt{1 + v^2}} dv, \quad \text{putting } \frac{y}{x} = v; \end{aligned}$$

$$\therefore u = \sqrt{1+v^2} + C = \frac{1}{x} \sqrt{x^2+y^2} + C.$$

94. To integrate $du = Pdx + Qdy$, u being a function of two independent variables x and y , and P and Q functions of x and y . This amounts to finding the integral of a differential function of two variables of the first order; and involves the same process as was used at Art. 7 for exact differential equations of the first order between two variables; since all such equations by being multiplied by an integrating factor are rendered differential functions.

Since $Pdx + Qdy$ must be identical with $\frac{du}{dx} dx + \frac{du}{dy} dy$, we have

$$\frac{du}{dx} = P, \quad \frac{du}{dy} = Q, \quad \text{with the condition} \quad \frac{dP}{dy} = \frac{dQ}{dx};$$

$$\therefore u = \int dx P + f(y), \quad f(y) \text{ involving } y \text{ only};$$

$$\therefore Q = \frac{du}{dy} = \frac{d}{dy} (\int dx P) + \frac{d}{dy} f(y);$$

$$\therefore f(y) = \int dy \left\{ Q - \frac{d}{dy} (\int dx P) \right\};$$

consequently u is known; and we may observe that in finding the value of $f(y)$ it is only necessary to integrate those terms of Q which involve y only. If we had begun by integrating with respect to y , the result would have been

$$u = \int dy Q + \int dx \left\{ P - \frac{d}{dx} (\int dy Q) \right\};$$

and as before it is only necessary to integrate those terms of P which involve x only.

Ex. 1.
$$du = \frac{y}{(x+y)^2} dx - \frac{x}{(x+y)^2} dy;$$

here
$$\frac{du}{dx} = \frac{y}{(x+y)^2}, \quad \frac{du}{dy} = -\frac{x}{(x+y)^2},$$

$$\therefore u = -\frac{y}{x+y} + f(y);$$

and differentiating with respect to y

$$\frac{-1}{x+y} + \frac{y}{(x+y)^2} + \frac{d}{dy}f(y) = \frac{du}{dy} = -\frac{x}{(x+y)^2};$$

$$\therefore \frac{d}{dy}f(y) = 0, \text{ and } f(y) = C,$$

as we might have foreseen, since there is no term in Q that involves y only;

$$\therefore u = C - \frac{y}{x+y}.$$

Ex. 2.
$$du = \frac{ydy + xdx - 2xdy}{(x-y)^2},$$

$$u = \log(x-y) - \frac{y}{x-y} + C.$$

95. To integrate $du = Pdx + Qdy + Rdz$, u being a function of three independent variables x, y, z .

Since $Pdx + Qdy + Rdz$ is identical with

$$\frac{du}{dx} \cdot dx + \frac{du}{dy} \cdot dy + \frac{du}{dz} \cdot dz,$$

we have
$$\frac{du}{dx} = P, \quad \frac{du}{dy} = Q, \quad \frac{du}{dz} = R;$$

together with the equations of condition

$$\frac{dP}{dy} = \frac{dQ}{dx}, \quad \frac{dQ}{dz} = \frac{dR}{dy}, \quad \frac{dR}{dx} = \frac{dP}{dz},$$

which are found by supposing each of the quantities z, x, y to be constant in succession (as we are evidently at liberty to do, since those variables are independent of one another), and then taking the corresponding equation of condition for du being the exact differential of a function of two independent variables. Hence

$$u = \int dx P + w, \quad w \text{ being a function of } y \text{ and } z;$$

$$\text{also } Q = \frac{d}{dy}(\int dx P) + \frac{dw}{dy},$$

$$R = \frac{d}{dz}(\int dx P) + \frac{dw}{dz},$$

which two equations giving the values of the partial differential coefficients of w , its value may be found by the preceding Article; and so the value of u completely determined.

Ex. 1. $du = yzdx + xzdy + xydz$;

here $\frac{du}{dx} = yz$, $\frac{du}{dy} = xz$, $\frac{du}{dz} = xy$; $\therefore u = xyz + w$;

$$xz = xz + \frac{dw}{dy}, \quad \text{or } \frac{dw}{dy} = 0;$$

$$xy = xy + \frac{dw}{dz}, \quad \text{or } \frac{dw}{dz} = 0;$$

$$\therefore w = C, \text{ and } u = xyz + C.$$

Ex. 2. $du = \frac{y}{a-z} dx + \frac{x}{a-z} dy + \frac{xy}{(a-z)^2} dz$;

$$u = \frac{xy}{a-z} + C.$$

96. Differential equations involving more than two variables admit of division into two classes, Total and Partial. A total differential equation is one which expresses the relation between the differential of the dependent variable and the other variables and their differentials, and sometimes also the dependent variable itself; it is consequently equivalent to a system of equations in which each differential coefficient of the dependent variable is given explicitly. Thus z being a function of the independent variables x and y , the total differential equation

$$Pdx + Qdy + Rdz = 0,$$

in which P , Q , and R are functions of x , y and z , amounts to the same as the two equations

$$P + R \frac{dz}{dx} = 0, \quad Q + R \frac{dz}{dy} = 0.$$

A Partial differential equation, on the contrary, is a relation between all or certain of the partial differential coefficients of the dependent variable, and the variables: as in the instances

$$(x-a) \frac{dz}{dx} + (y-b) \frac{dz}{dy} = z - c,$$

$$x^2 \frac{d^2z}{dx^2} + 2xy \frac{d}{dx} \frac{dz}{dy} + y^2 \frac{d^2z}{dy^2} = 0,$$

z in both being a function of the independent variables x and y .

Total Differential Equations.

97. We may first consider the equation

$$Pdx + Qdy + Rdz = 0,$$

one of the three variables x, y, z , being a function of the other two, which are independent.

Since the proposed equation may arise from combining the results of differentiating two separate equations, we have first to examine whether it can be satisfied by a single primitive relation between x, y and z . If x, y, z be co-ordinates of a point, the cases will be distinguished according as the proposed equation is the differential equation to a series of surfaces, or to a series of curves in space.

For example, the equation

$$(z-c) x dx + (z-c) y dy = \{x(x-a) + y(y-b)\} dz,$$

arises from combining the results of differentiating the equations

$$z = f(x^2 + y^2), \quad \frac{y-b}{z-c} = \phi\left(\frac{x-a}{z-c}\right);$$

for these equations give respectively

$$y \frac{dz}{dx} - x \frac{dz}{dy} = 0, \quad (x-a) \frac{dz}{dx} + (y-b) \frac{dz}{dy} = z - c;$$

from which if $\frac{dz}{dx}$ and $\frac{dz}{dy}$ be determined, and substituted in

$$dz = \frac{dz}{dx} \cdot dx + \frac{dz}{dy} \cdot dy,$$

we get the proposed equation; which, consequently, cannot in general be satisfied by a single relation between x, y, z . It is the

analytical expression of the conditions of the problem, to find a surface belonging at the same time to surfaces of revolution about the axis of z , and to conical surfaces.

98. To find the equation of condition for

$$Pdx + Qdy + Rdz = 0,$$

admitting a solution of the form $f(x, y, z) = C$.

If the proposed equation can be satisfied by a single relation between x , y and z , then

$$dz = -\frac{P}{R}dx - \frac{Q}{R}dy$$

is the differential of a function of two independent variables ;

$$\therefore \frac{dz}{dx} = -\frac{P}{R}, \quad \frac{dz}{dy} = -\frac{Q}{R},$$

$$\text{with the condition } \frac{d}{dy} \left(\frac{P}{R} \right) = \frac{d}{dx} \left(\frac{Q}{R} \right);$$

or, since P , Q , R may contain z which is a function of x and y ,

$$\frac{d}{dy} \left(\frac{P}{R} \right) + \frac{d}{dz} \left(\frac{P}{R} \right) \frac{dz}{dy} = \frac{d}{dx} \left(\frac{Q}{R} \right) + \frac{d}{dz} \left(\frac{Q}{R} \right) \frac{dz}{dx};$$

therefore, substituting for $\frac{dz}{dx}$, $\frac{dz}{dy}$ their values,

$$R \frac{d}{dy} \left(\frac{P}{R} \right) - Q \frac{d}{dz} \left(\frac{P}{R} \right) = R \frac{d}{dx} \left(\frac{Q}{R} \right) - P \frac{d}{dz} \left(\frac{Q}{R} \right);$$

and performing the differentiations and reducing we get

$$P \left(\frac{dR}{dy} - \frac{dQ}{dz} \right) + Q \left(\frac{dP}{dz} - \frac{dR}{dx} \right) + R \left(\frac{dQ}{dx} - \frac{dP}{dy} \right) = 0,$$

the equation of condition for

$$Pdx + Qdy + Rdz = 0$$

admitting a solution of the form $f(x, y, z) = C$.

99. If the above equation of condition be not satisfied, then the equation

$$Pdx + Qdy + Rdz = 0$$

cannot, by being multiplied by any factor, become susceptible of a solution of the form

$$f(x, y, z, C) = 0.$$

For suppose V to be a factor which renders

$$Pdx + Qdy + Rdz$$

the immediate differential of some function u of x, y, z considered as independent of one another ;

$$\therefore \frac{d}{dy}(VP) = \frac{d}{dx}(VQ), \quad \frac{d}{dz}(VQ) = \frac{d}{dy}(VR),$$

$$\frac{d}{dx}(VR) = \frac{d}{dz}(VP);$$

$$\text{or } \left. \begin{aligned} V \frac{dP}{dy} + P \frac{dV}{dy} &= V \frac{dQ}{dx} + Q \frac{dV}{dx} \\ V \frac{dQ}{dz} + Q \frac{dV}{dz} &= V \frac{dR}{dy} + R \frac{dV}{dy} \\ V \frac{dR}{dx} + R \frac{dV}{dx} &= V \frac{dP}{dz} + P \frac{dV}{dz} \end{aligned} \right\} \dots\dots\dots (1).$$

Hence, multiplying the first of these equations by R , the second by P , and the third by Q , and taking their sum, the factor V disappears, and we find

$$P \left(\frac{dR}{dy} - \frac{dQ}{dz} \right) + Q \left(\frac{dP}{dz} - \frac{dR}{dx} \right) + R \left(\frac{dQ}{dx} - \frac{dP}{dy} \right) = 0, \dots\dots (2)$$

the same equation of condition as in the preceding Article.

100. If this equation be satisfied, which will be the case only when the proposed admits a primitive of the form

$$f(x, y, z, C) = 0,$$

equations (1) afford a means of determining V . Then

$$du = VPdx + VQdy + VRdz,$$

whence u can be found by the method of Art. 95, and $u + C = 0$ is the required relation between x, y , and z .

Or, without determining V , we may integrate considering one of the variables constant, and add an arbitrary function of that variable; then differentiate the result with respect to that variable and compare it with the proposed equation, and so the function added will become known.

In the majority of cases which present themselves, the factor V is capable of being determined by inspection.

Ex. 1. $(ay - bz) dx + (cz - ax) dy + (bx - cy) dz = 0.$

Divide by $(ay - bz)(bx - cy)$, and the resulting equation

$$\frac{dx}{bx - cy} + \frac{(cz - ax) dy}{(ay - bz)(bx - cy)} + \frac{dz}{ay - bz} = 0,$$

satisfies the conditions, considering x, y, z as independent,

$$\frac{dP}{dy} = \frac{dQ}{dx}, \quad \frac{dQ}{dz} = \frac{dR}{dy}, \quad \frac{dR}{dx} = \frac{dP}{dz};$$

and therefore the first member may be regarded as the differential of some function, u , of x, y, z considered as independent;

$$\therefore \frac{du}{dx} = \frac{1}{bx - cy}; \quad \text{and } u = \frac{1}{b} \log (bx - cy) + w;$$

$$\therefore \frac{1}{ay - bz} = \frac{dw}{dz},$$

$$\frac{cz - ax}{(ay - bz)(bx - cy)} = -\frac{c}{b} \cdot \frac{1}{bx - cy} + \frac{dw}{dy},$$

$$\text{or } \frac{dw}{dy} = \frac{a}{b} \cdot \frac{-1}{ay - bz}; \quad \text{hence } dw = -\frac{1}{b} \cdot \frac{ady - bdz}{ay - bz};$$

$$\therefore w = -\frac{1}{b} \log (ay - bz) + C;$$

$$\therefore u = \frac{1}{b} \log \left(\frac{bx - cy}{ay - bz} \right) + C = 0, \quad \text{or } \frac{bx - cy}{ay - bz} = C.$$

Ex. 2. $xa^2x + ydy + (x^2 + y^2)(1 - z^{-1})dz = 0, \quad x^2 + y^2 = Cz^2e^{-2z}.$

When in an equation of this sort, the differentials enter above the first degree, it is not integrable unless it can be resolved

into rational factors of the form $Pdx + Qdy + Rdz$; for whatever be the integral, it must upon differentiation produce a result of that form.

101. If the equation

$$Pdx + Qdy + Rdz = 0$$

be susceptible of a primitive of the form $f(x, y, z, C) = 0$, and be homogeneous and of n dimensions with respect to x, y, z ; then, putting $x = vz$, $y = wz$, and dividing by z^n , it becomes

$$S(vdz + zdv) + T(wdz + zdw) + Udz = 0,$$

S, T, U being functions of v and w ;

$$\text{or } -\frac{dz}{z} = \frac{Sdv + Tdw}{Sv + Tw + U};$$

hence the second member is an exact differential since the first is so, and it may be sometimes integrated by inspection, or by the method of Art. 94.

$$\text{Ex. 1. } (y+z)dx + (x+z)dy + (x+y)dz = 0;$$

$$\text{put } x = vz, \quad y = wz,$$

$$\therefore \frac{dz}{z} + \frac{(w+1)dv + (v+1)dw}{v(w+1) + w(v+1) + v+w} = 0,$$

$$\text{or } \frac{dz}{z} + \frac{1}{2} \frac{d(v+w+vw)}{v+w+vw} = 0;$$

$$\therefore \log z^2 (v+w+vw) = \log C, \quad \text{or } xz + yz + xy = C.$$

$$\text{Ex. 2. } (y^2 + yz + z^2)dx + (x^2 + xz + z^2)dy$$

$$+ (x^2 + xy + y^2)dz = 0.$$

$$\text{Make } x = vz, \quad y = wz,$$

$$\therefore \frac{dz}{z} + \frac{(w^2 + w + 1)dv + (v^2 + v + 1)dw}{v(w^2 + w + 1) + w(v^2 + v + 1) + v^2 + vw + w^2} = 0;$$

but of the latter fraction, the denominator

$$= vw(w+v+1) + v(w+1+v) + w(v+1+w)$$

$$= (vw + v + w)(v + w + 1),$$

and the numerator

$$= (1 + v + w) d(v + w + vw) - (v + w + vw) d(v + w);$$

$$\therefore \frac{dz}{z} + \frac{d(v + w + vw)}{v + w + vw} - \frac{d(1 + v + w)}{1 + v + w} = 0.$$

$$\therefore \log \left\{ \frac{z(v + w + vw)}{1 + v + w} \right\} = \log C;$$

$$\therefore zx + zy + xy = C(z + x + y).$$

Ex. 3. $2(y + z) dx + (x + 3y + 2z) dy + (x + y) dz = 0.$

By the same substitution this becomes

$$\frac{3dz}{z} + 2 \frac{dv + dw}{v + w} + \frac{dw}{w + 1} = 0; \quad \therefore (x + y)^2 (y + z) = C.$$

Ex. 4. $(x^2 - y^2 + z^2) dx - z^2 dy + (y - x)(z^2 + xy + x^2) \frac{dz}{z} = 0;$

putting $x = vz$, $y = wz$, this is reduced to

$$dw = (1 + v^2 - w^2) dv;$$

$$\therefore \frac{e^{-v^2}}{w - v} = \int dv e^{-v^2} + C, \quad (\text{Ex. 4. Art. 18})$$

$$\text{or } \frac{z^2 e^{-\frac{x^2}{z^2}}}{y - x} = \int dx e^{-\frac{x^2}{z^2}} + Cz,$$

z being constant under the sign of integration.

102. We have seen that the equation

$$Pdx + Qdy + Rdz = 0,$$

when the equation of condition (Art. 98) is not satisfied, does not admit of being derived from a single primitive equation involving two independent variables. The integral in this case will be exhibited by a system of two equations; and the proposed equation cannot be regarded as the differential equation to a surface, but to a system of curves in space all endowed with some common property.

Ex. 1. $dz = aydx + bdy$.

Since the equation of condition in this case is not satisfied, x and y cannot be independent, and we may assume $y = f(x)$;

$$\therefore dz = af(x) dx + b \frac{d}{dx} f(x) dx,$$

$$\therefore z = a \int dx f(x) + bf(x),$$

which, with $y = f(x)$, constitutes the integral of the proposed equation.

In general, if V be a factor which makes $Pdx + Qdy$ an exact differential considering z as constant, and we find

$$\int (VPdx + VQdy) = w + \phi(z),$$

it is evident that

$$w + \phi(z) = 0, \text{ together with } \frac{dw}{dz} + \frac{d}{dz} \phi(z) - VR = 0,$$

satisfy the proposed equation, where $\phi(z)$ denotes any function of z .

Ex. 2. $zdx + xdy + ydz = 0$,

$$y + z \log x + \phi(z) = 0, \quad \log x + \frac{d}{dz} \phi(z) = \frac{y}{x}.$$

Ex. 3. $\{x(x-a) + y(y-b)\} dz = (z-c)(xdx + ydy)$,

$$x^2 + y^2 + 2\phi(z) = 0, \quad x(x-a) + y(y-b) + (z-c) \frac{d}{dz} \phi(z) = 0.$$

Partial Differential Equations. Equations of the first order.

103. In partial differential equations of two independent variables, the differential coefficients of the first order $\frac{dz}{dx}$, $\frac{dz}{dy}$, of the dependent variable z , are usually denoted by the symbols p and q ; and $\frac{d^2z}{dx^2}$, $\frac{d^2z}{dx dy}$, $\frac{d^2z}{dy^2}$, the differential coefficients of the second order, by r , s , t , respectively. A partial differential equation is said to be of the n^{th} order, when it involves one or more of the partial differential coefficients of the dependent variable of the n^{th} order; but none of a superior

order. To be the general equation of the n^{th} order, it ought to contain the independent variables, and the dependent variable together with all its partial differential coefficients from the first order to the n^{th} order inclusive. To integrate a partial differential equation, is to find for the dependent variable an expression between the differential coefficients of which that relation exists which is indicated by the proposed equation; and under the most general form possible.

104. The complete integral of $f(x, y, z, p, q) = 0$, the general equation of the first order, will involve one arbitrary or general function.

For let $u = F\{x, y, z, \phi(v)\} = 0$, be an equation by virtue of which z is a function of the independent variables x and y , v being a known function of x, y , and z . Then p and q are given by the equations $\frac{du}{dx} = 0$, $\frac{du}{dy} = 0$, each of which will involve $\frac{d}{dv} \phi(v)$ or $\phi'(v)$, and may involve $\phi(v)$; consequently, between the three equations $u = 0$, $\frac{du}{dx} = 0$, $\frac{du}{dy} = 0$, it will be possible to eliminate $\phi(v)$ and $\phi'(v)$, and there will result a relation $f(x, y, z, p, q) = 0$, wholly independent of the form of the function $\phi(v)$; and it is evident that in general more than one function cannot thus be eliminated. Conversely, an equation of the form

$$f(x, y, z, p, q) = 0$$

being proposed, its complete integral to have all the generality possible must be of the form $F\{x, y, z, \phi(v)\} = 0$, where the form of $\phi(v)$ is arbitrary.

For example, let

$$u = z + mx + ny - \phi(v) = 0,$$

$$\text{where } v = (x - a)^2 + (y - b)^2 + (z - c)^2;$$

$$\text{then } \frac{du}{dx} = p + m - \phi'(v) \{2(x - a) + 2(z - c)p\} = 0,$$

$$\frac{du}{dy} = q + n - \phi'(v) \{2(y - b) + 2(z - c)q\} = 0.$$

Hence, transposing and dividing one result by the other to eliminate $\phi'(v)$, we find

$$\frac{p+m}{q+n} = \frac{(x-a) + (z-c)p}{(y-b) + (z-c)q},$$

$$\begin{aligned} \text{or } p \{y-b-n(z-c)\} - q \{x-a-m(z-c)\} \\ = n(x-a) - m(y-b), \end{aligned}$$

the partial differential equation of which the complete integral is

$$z + mx + ny = \phi \{ (x-a)^2 + (y-b)^2 + (z-c)^2 \}.$$

105. To integrate an equation in which only one of the differential coefficients of the first order enters with x , y , and z .

Let the equation be

$$f(x, y, z, p) = 0.$$

Integrate it, considering y as a constant, and in place of the arbitrary constant C , add a function of y of arbitrary form. The resulting solution, containing one arbitrary function, will have all the generality that can be attained.

$$\text{Ex} \quad (x^2 + y^2) \frac{dz}{dx} = z^2 + y^2;$$

$$\therefore \frac{1}{z^2 + y^2} \frac{dz}{dx} - \frac{1}{x^2 + y^2} = 0;$$

$$\therefore \tan^{-1} \frac{z}{y} - \tan^{-1} \frac{x}{y} = \tan^{-1} \phi(y),$$

$$\text{or } z - x = (y^2 + xz) \phi(y),$$

$\phi(y)$ being arbitrary in form.

The equation $f(x, y, z, q) = 0$ is similarly integrated, the correction in this case being an arbitrary function of x .

$$\text{Ex.} \quad xy \frac{dz}{dy} + nz = 0,$$

$$z^x y^n = \phi(x).$$

106. To integrate the equation of the first order,

$$Pp + Qq = R,$$

P , Q , and R being functions of x , y , and z .

Let the primitive be $F(x, y, z) = 0$; therefore, denoting $\frac{d}{dx} F(x, y, z)$ by $F'(x)$, and so on for the other differential coefficients, we get

$$F'(x) + F'(z) \cdot p = 0,$$

$$F'(y) + F'(z) \cdot q = 0;$$

$$\therefore PF'(x) + QF'(y) + RF'(z) = 0.$$

$$\text{But } dF(x, y, z) = F'(x) dx + F'(y) dy + F'(z) dz = 0,$$

$$\therefore PF'(x) dx + PF'(y) dy + PF'(z) dz = 0;$$

$$\text{and } PF'(x) dx + QF'(y) dx + RF'(z) dx = 0;$$

$$\therefore F'(y) \{Pdy - Qdx\} + F'(z) \{Pdz - Rdx\} = 0, \quad (1)$$

which is satisfied by

$$\left. \begin{aligned} Pdy - Qdx &= 0, \\ Pdz - Rdx &= 0. \end{aligned} \right\} \quad (2).$$

Suppose that by integrating these equations either separately or conjointly, we obtain $M = a$, $N = b$, two relations between the three variables and the arbitrary constants a and b , which satisfy them. By means of the equations $M = a$, $N = b$, any two of the variables as y and z can be expressed in terms of a and b and the third variable x . The complete primitive then becomes

$$F(x, y, z) = \phi(x, a, b) = 0,$$

and the differential of $\phi(x, a, b)$ must by virtue of equations (2) be identically equal to zero, therefore $\phi(x, a, b)$ cannot contain x , and consequently

$$0 = \phi(a, b) = \phi(M, N), \text{ or } M = f(N).$$

The assumptions (2) satisfy equation (1) independently of the forms of $F'(y)$, $F'(x)$, that is, independently of the form of F ; therefore the form of ϕ , and consequently of f , is arbitrary.

Hence $M=f(N)$ being a solution of the proposed equation and containing one arbitrary function, is the general primitive.

107. For the success of the method, it is in general necessary that of the three equations

$$Pdy - Qdx = 0, \quad Pdz - Rdx = 0, \quad Qdz - Rdy = 0,$$

one at least should contain those two variables only whose increments or differentials appear in it, or that by their combination such an equation should be produced. For by integrating it we shall obtain an equation by means of which one of the variables may be eliminated from either of the other auxiliary equations. They are called the reducing equations of

$$Pp + Qq = R;$$

and may be easily remembered under the form

$$\frac{dx}{P} = \frac{dy}{Q} = \frac{dz}{R}.$$

108. By a similar process, partial differential equations of three or more independent variables can sometimes be integrated. If z be a function of x, y, t , and the equation be

$$Tn + Pp + Qq = R,$$

where $n = \frac{dz}{dt}$, we have

$$n = \frac{R - Pp - Qq}{T};$$

which, substituted in

$$dz = ndt + pdx + qdy,$$

$$\text{gives } Tdz - Rdt = p(Tdx - Pdt) + q(Tdy - Qdt).$$

And if the equations $\frac{dx}{P} = \frac{dy}{Q} = \frac{dz}{R} = \frac{dt}{T}$, or

$$Tdx - Pdt = 0, \quad Tdy - Qdt = 0,$$

$$\text{give } L = a, \quad M = b, \quad N = c,$$

then the solution is

$$\phi(L, M, N) = 0, \quad \text{or } L = f(M, N).$$

109. To explain the geometrical meaning of the solution of a partial differential equation of the first order.

If we regard x, y, z as the co-ordinates of a point, and the proposed differential equation as the equation to a system of surfaces, $M=a$, $N=b$ are the equations to two surfaces which, being satisfied by the same values of x, y, z that satisfy the differential equation, conjointly represent a line situated on one of the surfaces represented by the primitive. By giving any alterations to a and b we obtain other lines in space; but in order that these lines may always lie on the surface in question, these alterations must not be independent but connected, according to some law expressed by the relation $a=f(b)$, or $M=f(N)$.

For example, the equation $px+qy=z$ is the differential equation to a conical surface, and

$$\frac{y}{z}=a, \quad \frac{x}{z}=b,$$

are the equations to any straight line through the vertex. If a and b undergo all possible variations consistently with the restriction imposed by the nature of the directrix, the assemblage of straight lines thus formed is the conical surface.

110. We shall now apply the preceding method, which is due to Lagrange, to some examples.

Ex. 1. $px + qz + y = 0.$

This compared with $Pp + Qq = R$, gives in the place of

$$Qdz - Rdy = 0, \quad Pdz - Rdx = 0,$$

the reducing equations

$$zdz + ydy = 0, \quad xdz + ydx = 0.$$

From the former, which alone is immediately integrable, we get $z^2 + y^2 = a^2$; then by substitution in the latter

$$xdz + \sqrt{a^2 - z^2}dx = 0, \quad \text{or} \quad \frac{dz}{\sqrt{a^2 - z^2}} + \frac{dx}{x} = 0;$$

$$\therefore \sin^{-1} \frac{z}{a} + \log x = \log b, \quad \text{or} \quad xe^{\sin^{-1} \frac{z}{a}} = b;$$

but $a = \sqrt{z^2 + y^2}$, therefore $\sin^{-1} \frac{z}{a} = \tan^{-1} \frac{z}{y}$; and $b = f(a)$,

$$\text{hence } xe^{\tan^{-1} \frac{z}{y}} = f(\sqrt{z^2 + y^2}),$$

is the integral of the proposed equation.

Ex. 2. $p(x-a) + q(y-b) = z-c.$

Here the reducing equations are

$$(x-a) dz - (z-c) dx = 0, \quad (y-b) dz - (z-c) dy = 0;$$

$$\therefore \frac{x-a}{z-c} = \alpha, \quad \frac{y-b}{z-c} = \beta,$$

$$\therefore \frac{y-b}{z-c} = f\left(\frac{x-a}{z-c}\right).$$

Ex. 3. $mp + nq = 1;$

$$\therefore mdz - dx = 0, \quad ndz - dy = 0;$$

$$\therefore x - mz = \alpha, \quad y - nz = \beta;$$

$$\therefore y - nz = f(x - mz).$$

Ex. 4. $(x - mz)p + (y - nz)q = 0;$

$$\therefore (x - mz) dz = 0; \quad \therefore z = a,$$

$$(x - mz) dy - (y - nz) dx = 0,$$

$$\text{or } (x - ma) dy - (y - na) dx = 0;$$

$$\frac{x - ma}{y - na} = b; \quad \text{but } a = f(b), \quad \therefore z = f\left(\frac{x - mz}{y - nz}\right).$$

5. $px + qy = nz, \quad z = x^n f\left(\frac{y}{x}\right).$

6. $\frac{p}{y} + \frac{q}{x} = \frac{1}{z}, \quad z^2 = xy + f\left(\frac{y}{x}\right).$

7. $p\dot{x}^2 + q\dot{y}^2 = z^2, \quad \frac{1}{z} = \frac{1}{x} + f\left(\frac{1}{y} - \frac{1}{x}\right).$

8. $z - px - qy = n\sqrt{x^2 + y^2 + z^2},$

$$x^{n-1} (z + \sqrt{x^2 + y^2 + z^2}) = f\left(\frac{y}{x}\right).$$

$$9. \quad p(y+x) + q(y-x) = z,$$

$$\tan^{-1} \frac{x}{y} - \log(x^2 + y^2)^{\frac{1}{2}} = f\left(\frac{z}{\sqrt{x^2 + y^2}}\right).$$

$$10. \quad x \frac{dz}{dx} + y \frac{dz}{dy} + t \frac{dz}{dt} + u \frac{dz}{du} = nz, \quad z = t^n f\left(\frac{x}{t}, \frac{y}{t}, \frac{u}{t}\right).$$

111. The following examples require artifices in the combination of the reducing equations.

$$\text{Ex. 1.} \quad (2az + cx)p + (2bz - cy)q = (ay + bx)c.$$

The reducing equations are

$$(ay + bx)cdx = (2az + cx)dz,$$

$$(ay + bx)c dy = (2bz - cy)dz.$$

Multiply the first by y , and the second by x , and add them together, then

$$cydx + cxdy = 2zdz, \text{ or } z^2 - cxy = a.$$

Multiply the first by $-b$, and the second by a , and add them together, then

$$-b dx + a dy = -dz, \text{ or } z + ay - bx = \beta;$$

$$\therefore z + ay - bx = \phi(z^2 - cxy).$$

$$\begin{aligned} \text{Ex. 2.} \quad \{y - b - n(z - c)\}p - \{x - a - m(z - c)\}q \\ = n(x - a) - m(y - b), \end{aligned}$$

the equation to surfaces of revolution. Here the reducing equations are

$$\{n(x - a) - m(y - b)\}dx - \{y - b - n(z - c)\}dz = 0, \quad (1).$$

$$\{y - b - n(z - c)\}dy + \{x - a - m(z - c)\}dx = 0, \quad (2).$$

$$\{x - a - m(z - c)\}dz + \{n(x - a) - m(y - b)\}dy = 0, \quad (3).$$

Multiply (1) by $x - a$, and (3) by $y - b$, and add, then

$$\{n(x - a) - m(y - b)\}\{(x - a)dx + (y - b)dy + (z - c)dz\} = 0;$$

$$\therefore (x - a)dx + (y - b)dy + (z - c)dz = 0;$$

$$\text{or } (x - a)^2 + (y - b)^2 + (z - c)^2 = \alpha.$$

Multiply (1) by m , and (3) by n , and add, then

$$\{n(x-a) - m(y-b)\} (mdx + ndy + dz) = 0;$$

$$\therefore mdx + ndy + dz = 0,$$

$$\text{or } mx + ny + z = \beta = f(x);$$

$$\therefore z + mx + ny = f\{(x-a)^2 + (y-b)^2 + (z-c)^2\}.$$

Ex. 3. $(y+z)p + (x+z)q = x+y.$

The reducing equations are

$$(y+z) dz = (x+y) dx,$$

$$(x+z) dz = (x+y) dy;$$

$$\therefore (y-x) dz = -(x+y) (dy - dx)$$

$$(x+y) (dx + dy + dz) = 2(x+y+z) dz;$$

$$\therefore \frac{dx + dy + dz}{x + y + z} = \frac{2dz}{x + y} = -\frac{2(dy - dx)}{y - x};$$

$$\therefore (x+y+z) (y-x)^2 = \alpha.$$

Similarly,

$$(x+y+z) (y-z)^2 = \beta;$$

$$\therefore F\{(x+y+z) \cdot (y-x)^2, (x+y+z) \cdot (z-y)^2\} = 0.$$

Ex. 4. $(x+y+z) \frac{dz}{dt} + (t+y+z) \frac{dz}{dx}$

$$+ (t+x+z) \frac{dz}{dy} = t+x+y.$$

The reducing equations are

$$(x+y+z) dz = (t+x+y) dt,$$

$$(x+y+z) dx = (t+y+z) dt,$$

$$(x+y+z) dy = (t+x+z) dt;$$

$$\therefore (x+y+z) (dz - dt) = (t-z) dt,$$

$$(x+y+z) (dx - dt) = (t-x) dt,$$

$$(x+y+z) (dy - dt) = (t-y) dt,$$

$$(x+y+z) (dx + dy + dz + dt) = 3(t+x+y+z) dt;$$

$$\text{hence } \frac{dz - dt}{t - z} = \frac{dt}{x + y + z} = \frac{dx + dy + dz + dt}{3(x + y + z + t)};$$

$$\therefore (x + y + z + t) \cdot (z - t)^3 = \alpha; \text{ similarly}$$

$$(x + y + z + t) \cdot (x - t)^3 = \beta, \quad (x + y + z + t) \cdot (y - t)^3 = \gamma;$$

$$\therefore (x + y + z + t) \cdot (z - t)^3 =$$

$$F\{(x + y + z + t) \cdot (x - t)^3, (x + y + z + t) \cdot (y - t)^3\}.$$

Method of solving linear Partial Differential Equations by separation of symbols.

112. As the symbols of operation here employed $\frac{d}{dx}$, $\frac{d}{dy}$ are independent of one another, and as they are subject in their combinations to the same laws as algebraical quantities, we may by separating the symbols of operation from those of quantity readily effect the solution of many partial differential equations by the same processes as were employed for differential equations between two variables of the like class. In particular the whole class of Linear Partial Differential Equations bears a strict analogy to equations of the same sort between two variables, both in the mode of treatment which they admit, and in the form of the solution obtained.

Great use will be here made of the symbolical expressions for Taylor's Theorem, applied to a function of one or more variables: viz.

$$f(x + h) = e^{h \frac{d}{dx}} f(x), \quad f(x + h, y) = e^{h \frac{d}{dx}} f(x, y),$$

$$f(x + h, y + k) = e^{h \frac{d}{dx} + k \frac{d}{dy}} f(x, y);$$

and this use of the symbols $e^{h \frac{d}{dx}}$, $e^{k \frac{d}{dy}}$, must be particularly attended to, as having an important bearing on the interpretation of the results obtained. For since, when only two independent variables x and y are involved, after every integration relative to x an arbitrary function of y , $\phi(y)$, must be added instead of an arbitrary constant, if an operating factor such as $\left(\frac{d}{dx} - a \frac{d}{dy}\right)^{-1}$

be applied to a subject $f(x, y)$, we shall have by Art. 53, (taking notice that in operating relative to x , we must treat y and the symbol $\frac{d}{dy}$ like simple constants; and the same holds relative to x and $\frac{d}{dx}$ when integrating or differentiating relative to y),

$$\begin{aligned} \left(\frac{d}{dx} - a \frac{d}{dy}\right)^{-1} f(x, y) &= e^{ax \frac{d}{dy}} \{ \int dx e^{-ax \frac{d}{dy}} f(x, y) + \phi(y) \} \\ &= e^{ax \frac{d}{dy}} \int dx f(x, y - ax) + \phi(y + ax), \end{aligned}$$

where we see that an arbitrary function of a binomial, viz. $\phi(y + ax)$ here takes the place of the term Ce^{ax} consisting of an arbitrary constant multiplied by the exponential quantity e^{ax} , which is always met with in the solutions of linear equations between two variables. In the following examples we shall often for the sake of convenience write d , d' , d'' instead of $\frac{d}{dx}$, $\frac{d}{dy}$, $\frac{d}{dt}$, suppressing the differentials; and arbitrary functions of x and y will sometimes be denoted by c_x , c_y .

113. We shall begin with the linear equation of the first order having constant coefficients,

$$a \frac{dz}{dx} + b \frac{dz}{dy} + cz = f(x, y),$$

$$\text{or } \left(a \frac{d}{dx} + b \frac{d}{dy} + c\right) z = f(x, y);$$

$$\therefore az = \left(\frac{d}{dx} + \frac{c + bd'}{a}\right)^{-1} f(x, y) = e^{-\frac{x}{a}(c + bd')} \{ \int dx e^{\frac{x}{a}(c + bd')} f(x, y) + \phi(y) \},$$

$$\text{or } e^{\frac{cx}{a}} az = e^{-\frac{bx}{a}} \int dx e^{\frac{cx}{a}} f\left(x, y + \frac{bx}{a}\right) + \phi\left(y - \frac{bx}{a}\right).$$

Suppose $f(x, y)$ to be a rational integral algebraic function of n dimensions;

$$\begin{aligned} \text{then } z &= \frac{1}{c} \left\{ 1 + \frac{1}{c} (ad + bd') \right\}^{-1} f(x, y) \\ &= \frac{1}{c} \left\{ 1 - \frac{1}{c} (ad + bd') + \frac{1}{c^2} (ad + bd')^2 - \&c. \pm \frac{1}{c^n} (ad + bd')^n \right\} f(x, y) \\ &\quad + e^{-\frac{cx}{a}} \phi \left(y - \frac{bx}{a} \right); \end{aligned}$$

where we stop with the n^{th} term of the development of the operating function, because any power of d or d' higher than the n^{th} , applied to $f(x, y)$, produces zero. Hence if $f(x, y) = g$, a constant,

$$z = \frac{g}{c} + e^{-\frac{cx}{a}} \phi \left(y - \frac{bx}{a} \right);$$

unless $c = 0$, when $z = \frac{gx}{a} + \phi \left(y - \frac{bx}{a} \right)$; and if $f(x, y) = gxy$,

$$z = g \frac{xy}{c} - g \frac{ay + bx}{c^2} + \frac{2gab}{c^3} + e^{-\frac{cx}{a}} \phi \left(y - \frac{bx}{a} \right).$$

Again, supposing $f(x, y)$ to be equal to ge^{mx+ny} , or to $g \cos(mx + ny)$, the values of z will be found to be, respectively,

$$z = \frac{ge^{mx+ny}}{am + bn + c} + e^{-\frac{cx}{a}} \phi \left(y - \frac{bx}{a} \right);$$

$$z = g \cdot \frac{c \cos(mx + ny) + (ma + nb) \sin(mx + ny)}{c^2 + (ma + nb)^2} + e^{-\frac{cx}{a}} \phi(ay - bx).$$

114. If the proposed equation be

$$a \frac{dz}{dx} + b \frac{dz}{dy} + cz^2 = zf(x, y),$$

then assuming $z = e^v$, or $v = \log z$, where v denotes a new function of x and y , the equation is reduced to the same form as the preceding, viz.:

$$a \frac{dv}{dx} + b \frac{dv}{dy} + cv = f(x, y),$$

hence the solution is

$$ae^{\frac{cx}{a}} \log z = e^{-\frac{bx}{a}} \int dx e^{\frac{cx}{a}} f\left(x, y + \frac{bx}{a}\right) + \phi\left(y - \frac{bx}{a}\right).$$

$$\text{Thus let } f(x, y) = \frac{\alpha y + \beta x}{(\alpha y - bx)y} = \frac{1}{b} \cdot \frac{ab + \beta a}{\alpha y - bx} - \frac{\beta}{by};$$

$$\therefore f\left(x, y + \frac{bx}{a}\right) = \frac{ab + \beta a}{ab} \cdot \frac{1}{y} - \frac{a\beta}{b} \cdot \frac{1}{\alpha y + bx};$$

$$\therefore a \log z = \frac{ab + \beta a}{b} \cdot \frac{x}{\alpha y - bx} - \frac{a\beta}{b^2} \log(\alpha y) + \phi\left(y - \frac{bx}{a}\right)$$

is the solution (where $c=0$) of

$$a \frac{dz}{dx} + b \frac{dz}{dy} = \frac{\alpha y + \beta x}{\alpha y - bx} \cdot \frac{z}{y}.$$

A particular case of this is

$$\frac{dz}{dx} - m \frac{dz}{dy} = \frac{az}{y} + \frac{bz}{mx + y},$$

$$\text{for which } z = y^{-\frac{a}{m}} \cdot e^{\frac{bx}{y+mx}} \cdot \phi(y + mx).$$

$$\text{Again, let } f(x, y) = \frac{\alpha}{x} + \frac{\beta}{y} + \frac{\gamma}{\sqrt{xy}},$$

$$\begin{aligned} \text{then } \int dx f\left(x, y + \frac{bx}{a}\right) &= \alpha \log x + \frac{a\beta}{b} \log\left(y + \frac{bx}{a}\right) \\ &\quad + 2\gamma \sqrt{\frac{a}{b}} \log\left(\sqrt{x} + \sqrt{\frac{\alpha y}{b} + x}\right); \end{aligned}$$

$$\begin{aligned} \therefore a \log z &= \alpha \log x + \frac{a\beta}{b} \log y + 2\gamma \sqrt{\frac{a}{b}} \log\left(\sqrt{x} + \sqrt{\frac{\alpha y}{b}}\right) \\ &\quad + \phi\left(y - \frac{bx}{a}\right) \end{aligned}$$

is the solution (where $c=0$) of

$$a \frac{dz}{dx} + b \frac{dz}{dy} = z \left(\frac{\alpha}{x} + \frac{\beta}{y} + \frac{\gamma}{\sqrt{xy}} \right).$$

A particular case of this is

$$\frac{dz}{dx} + m \frac{dz}{dy} = z \left(\frac{\alpha}{x} + \frac{\beta}{y} + \frac{\gamma}{\sqrt{xy}} \right),$$

$$\text{for which } z = x^{\frac{\alpha}{m}} y^{\frac{\beta}{m}} \left(\sqrt{x} + \sqrt{\frac{y}{m}} \right)^{\frac{2\gamma}{\sqrt{m}}} \cdot \phi(y - mx).$$

115. Next let an equation involving three independent variables be proposed,

$$\frac{dz}{dx} + a \frac{dz}{dy} + b \frac{dz}{dt} = x^2 y t,$$

$$\text{or } (d + ad' + bd'') z = x^2 y t;$$

$$\begin{aligned} \therefore z &= \frac{1}{d} \left(1 + \frac{ad' + bd''}{d} \right)^{-1} x^2 y t \\ &= \left(\frac{1}{d} - \frac{ad' + bd''}{d^2} + \frac{2ab d' d''}{d^3} \right) x^2 y t, \end{aligned}$$

these being the only terms necessary to be preserved in the development of the operating function, since $d'^2 y t = 0$; therefore, effecting the operations indicated,

$$z = \frac{1}{3} x^2 y t - \frac{1}{12} (at + by) x^4 + \frac{1}{30} ab x^5 + \phi(y - ax, t - bx);$$

the complementary function being

$$e^{-x(ad' + bd'')} \phi(y, t) = \phi(y - ax, t - bx).$$

116. We come next to the case of the complete linear equation of the second order with constant coefficients

$$(ad^2 + bdd' + cd'^2 + ad' + \beta d + \gamma) z = f(x, y).$$

This may be integrated whenever the operating function can be resolved into two factors rational with respect to d and d' ; the condition for which may be shewn, by putting the operating function equal to zero and solving it relative to d , to be

$$(b^2 - 4ac) \gamma = b\alpha\beta - a\alpha^2 - c\beta^2.$$

Thus, suppose $b = 1$, $a = c = 0$, then $\gamma = \alpha\beta$, and

$$(dd' + ad' + \beta d + \alpha\beta) z = f(x, y),$$

$$\text{or } \left(\frac{d}{dx} + \alpha\right) \left(\frac{d}{dy} + \beta\right) z = f(x, y);$$

$$\therefore \left(\frac{d}{dy} + \beta\right) z = \left(\frac{d}{dx} + \alpha\right)^{-1} f(x, y) = e^{-\alpha x} \int dx e^{\alpha x} f(x, y) + e^{-\alpha x} C_v,$$

$$\therefore z = e^{-\alpha x - \beta y} \int dy e^{\beta y} \{ \int dx e^{\alpha x} f(x, y) \} + e^{-\alpha x - \beta y} \int dy e^{\beta y} C_v' + e^{-\beta y} C_x'.$$

Hence, changing the arbitrary functions, we find

$$z = e^{-\alpha x - \beta y} \int dy e^{\beta y} \{ \int dx e^{\alpha x} f(x, y) \} + e^{-\alpha x} c_v + e^{-\beta y} c_x',$$

for the solution of

$$\frac{d^2 z}{dx dy} + \alpha \frac{dz}{dy} + \beta \frac{dz}{dx} + \alpha \beta z = f(x, y).$$

Hence if $f(x, y) = e^{mx+ny}$, since in this case $\frac{d}{dx}$ and $\frac{d}{dy}$ are equivalent respectively to factors m and n ,

$$z = \frac{e^{mx+ny}}{(m+\alpha)(n+\beta)} + e^{-\alpha x} c_v + e^{-\beta y} c_x'.$$

Again, let $f(x, y) = e^{mx} \cos ny$, then

$$\int dx e^{\alpha x} f(x, y) = \frac{e^{(m+\alpha)x} \cos ny}{m+\alpha};$$

$$\begin{aligned} \therefore z &= \frac{e^{mx-\beta y}}{m+\alpha} \int dy e^{\beta y} \cos ny \\ &= \frac{e^{mx}}{m+\alpha} \frac{n \sin ny + \beta \cos ny}{n^2 + \beta^2} + e^{-\alpha x} c_v + e^{-\beta y} c_x'. \end{aligned}$$

Similarly, if we suppose $a = 1$, $b = \gamma = 0$, then $c = -\frac{\alpha^2}{\beta^2}$, and the equation is

$$(d^2 - \frac{\alpha^2}{\beta^2} d'^2 + \alpha d' + \beta d) z = f(x, y),$$

$$\text{or } (d + \frac{\alpha}{\beta} d') (d - \frac{\alpha}{\beta} d' + \beta) z = f(x, y).$$

If we assume $f(x, y) = 0$, this gives

$$z = e^{-\beta x} e^{\frac{\alpha x}{\beta}} \phi(y) + e^{-\frac{\alpha x}{\beta}} \psi(y) = e^{-\beta x} \phi\left(y + \frac{\alpha x}{\beta}\right) + \psi\left(y - \frac{\alpha x}{\beta}\right)$$

for the solution of

$$\frac{d^2 z}{dx^2} - \frac{\alpha^2}{\beta^2} \frac{d^2 z}{dy^2} + \beta \frac{dz}{dx} + \alpha \frac{dz}{dy} = 0.$$

If $b^2 = 4ac$, we must have

$$b\alpha\beta = a\alpha^2 + c\beta^2, \quad \text{or} \quad b\beta = 2a\alpha;$$

then the proposed equation becomes

$$\left\{ \left(\sqrt{a}d + \sqrt{c}d' + \frac{\beta}{2\sqrt{a}} \right)^2 - \left(\frac{\beta^2}{4a} - \gamma \right) \right\} z = f(x, y);$$

and the operating function can be separated into two real factors, if $\beta^2 > 4a\gamma$.

117. Next, let the equation be of the n^{th} order

$$\left(\frac{d}{dx} \right)^n z + p_1 \left(\frac{d}{dx} \right)^{n-1} \frac{d}{dy} z + p_2 \left(\frac{d}{dx} \right)^{n-2} \left(\frac{d}{dy} \right)^2 z + \dots + p_n \left(\frac{d}{dy} \right)^n z = V,$$

the index of differentiation being the same in every term; and V denoting a function of x and y ; then the equation may be written

$$\left\{ \left(\frac{d}{dx} \right)^n + p_1 \left(\frac{d}{dx} \right)^{n-1} + p_2 \left(\frac{d}{dx} \right)^{n-2} + \dots + p_n \right\} d^n z = V.$$

Let a be a root that occurs singly, and b a root that occurs m times, in the auxiliary equation

$$v^n + p_1 v^{n-1} + p_2 v^{n-2} + \dots + p_n = 0;$$

then by resolution into partial fractions we shall have

$$\frac{1}{v^n + p_1 v^{n-1} + \dots + p_n} = \frac{A}{v-a} + \frac{B_m}{(v-b)^m} + \frac{B_{m-1}}{(v-b)^{m-1}} + \dots + \frac{B_1}{v-b} + \&c.$$

Hence substituting $\frac{d}{d'}$ for v , we get

$$z = \left\{ \frac{Ad'}{d-ad'} + \frac{B_m d'^m}{(d-bd')^m} + \frac{B_{m-1} d'^{m-1}}{(d-bd')^{m-1}} + \dots + \frac{B_1 d'}{d-bd'} + \&c. \right\} d'^n V;$$

$$\therefore z = A e^{axd'} \int dx e^{-axd'} d'^{n+1} V + B_m e^{bx d'} \int (dx)^m e^{-bx d'} d'^{n+m} V + \&c.$$

Instead of reserving the complementary functions under the sign of integration, we may obtain them explicitly by supposing $V=0$; then since

$$\int dx 0 = c_y, \quad (\int dx)^m 0 = c_y^0 + x c_y' + x^2 c_y'' + \dots + x^{m-1} c_y^{(m-1)},$$

the part of the complementary function due to the root a and the m roots b is, changing the arbitrary functions,

$$Ac_{y+ax} + B_m (c^0_{y+bx} + xc'_{y+bx} + x^2c''_{y+bx} + \dots + x^{m-1}c^{(m-1)}_{y+bx});$$

and similar terms will be introduced by the other roots of the auxiliary equation.

$$\text{Ex. 1. } \frac{d^2z}{dx^2} - n^2 \frac{d^2z}{dy^2} = cxy;$$

$$\therefore z = \frac{cxy}{d^2 - n^2 d'^2} = d^{-2} \left\{ 1 + n^2 \left(\frac{d'}{d} \right)^2 + \&c. \right\} cxy = d^{-2} cxy,$$

all the other terms being neglected, because when the operations are performed they vanish of themselves;

$$\therefore z = \frac{c}{6} x^3 y + c_{y+nx} + c'_{y-nx}.$$

$$\text{Ex. 2. } \left(\frac{d}{dx} \right)^3 z - 4 \left(\frac{d}{dx} \right)^2 \frac{d}{dy} z + 5 \frac{d}{dx} \left(\frac{d}{dy} \right)^2 z - 2 \left(\frac{d}{dy} \right)^3 z = e^{mx+ny},$$

$$\text{or } (d - 2d') (d - d')^2 z = e^{mx+ny};$$

$$\therefore z = \frac{e^{mx+ny}}{(m-2n)(m-n)^2} + c_{y+2x} + c^0_{y+x} + xc'_{y+x},$$

since d and d' operating upon e^{mx+ny} are equivalent respectively to the factors m and n .

If the second member of the equation be $x^2 + y^2$, then

$$z = \frac{x^2 + y^2}{d^3 - 4d^2d' + 5dd'^2 - 2d'^3} = d^{-3} \left(1 + 4 \frac{d'}{d} + 11 \frac{d'^2}{d^2} \right) (x^2 + y^2),$$

these being the only terms in the development of the operating function which it is necessary to retain;

$$\therefore z = \frac{x^3 y^2}{6} + \frac{x^4 y}{3} + \frac{x^5}{5} + c_{y+2x} + c^0_{y+x} + xc'_{y+x}.$$

$$\text{Ex. 3. } \frac{d^2z}{dx^2} + (a+b) \frac{d^2z}{dx dy} + ab \frac{d^2z}{dy^2} = f(x, y),$$

$$\text{or } (d + ad') (d + bd') z = f(x, y);$$

$$\therefore z = \frac{d'^{-1}}{a-b} \left(\frac{1}{d+bd'} - \frac{1}{d+ad'} \right) f(x, y),$$

$$= \frac{d'^{-1}}{a-b} \{ e^{-bx d'} \int dx f(x, y+bx) - e^{-ax d'} \int dx f(x, y+ax) \}.$$

Let $f(x, y) = 0$, then $(d+ad')(d+bd')z=0$ gives

$$z = e^{-ax d'} \phi(y) + e^{-bx d'} \psi(y) = \phi(y-ax) + \psi(y-bx),$$

which shews the form of the complementary functions, whatever be the second member of the equation. If $f(x, y)$ be a rational integral algebraic function of x and y , then

$$z = [d'^2 - d'^4 \{ (a+b)dd' + abd'^2 \} + \&c.] f(x, y) + \phi(y-ax) + \psi(y-bx).$$

Thus let $f(x, y) = c + m(x+y) + nxy$,

then $z = \{d'^2 - (a+b)d'^3 d'\} \{c + m(x+y) + nxy\}$

$$= \frac{1}{2} (c + my) x^2 + \frac{1}{6} (m + ny) x^3 - \frac{1}{24} (a+b) (4m + nx) x^3$$

$$+ \phi(y-ax) + \psi(y-bx).$$

Change of the Independent Variables.

118. The method by changing the independent variables is applicable to partial differential equations, and will generally lead to the solution of any equation of a form analogous to one of those for which the method succeeds in the case of differential equations between two variables. The change of the independent variables that has the most extensive application is to assume $x = e^t$, $y = e^v$, as we proceed to shew in the following instances, making use of the formulæ of Art. 70.

Ex. 1. $ax \frac{dz}{dx} + by \frac{dz}{dy} + cz = V;$

this becomes

$$a \frac{dz}{dt} + b \frac{dz}{dv} + cz = U,$$

the integral of which has already been found (Art. 113).

Let $V = x^m y^n$, then $U = e^{mt+nv};$

$$\therefore z = \frac{e^{mt+nv}}{am + bn + c} + e^{-\frac{ct}{a}} \phi\left(v - \frac{bt}{a}\right),$$

or, restoring the values of t and v , and changing the arbitrary function,

$$z = \frac{x^m y^n}{am + bn + c} + x^{-\frac{c}{a}} \phi\left(\frac{y^n}{x^b}\right).$$

$$\begin{aligned} \text{Ex. 2. } \quad ax^2 \frac{d^2 z}{dx^2} + (a+b) xy \frac{d^2 z}{dxdy} + by^2 \frac{d^2 z}{dy^2} + cx \frac{dz}{dx} \\ + cy \frac{dz}{dy} - cz = 0. \end{aligned}$$

This, by the same substitutions as in the foregoing example, becomes

$$ad(d-1)z + (a+b)dd'z + bd'(d'-1)z + c(d+d'-1)z = 0,$$

$$\text{or } (ad + bd' + c)(d + d' - 1)z = 0;$$

$$\begin{aligned} \therefore az &= e^{-\frac{ct}{a}} e^{-\frac{bt}{a'}} \phi(v) + e^{t(1-a')} \psi(v) \\ &= e^{-\frac{ct}{a}} \phi\left(v - \frac{bt}{a}\right) + e^t \psi(v - t); \end{aligned}$$

hence, restoring the values of v and t ,

$$z = x^{-\frac{c}{a}} \phi\left(yx^{-\frac{b}{a}}\right) + x\psi\left(\frac{y}{x}\right).$$

$$\begin{aligned} \text{Ex. 3. } \quad ax^3 \frac{d^3 z}{dx^3} + (2a+b)x^2 y \frac{d^3 z}{dx^2 dy} \\ + (a+2b)xy^2 \frac{d^3 z}{dxdy^2} + by^3 \frac{d^3 z}{dy^3} = 0; \end{aligned}$$

$$\begin{aligned} \therefore \{ad(d-1)(d-2) + (2a+b)d(d-1)d' \\ + (a+2b)dd'(d'-1) + bd'(d'-1)(d'-2)\}z = 0, \end{aligned}$$

$$\text{or } (ad + bd')(d + d' - 1)(d + d' - 2)z = 0;$$

$$\begin{aligned} \therefore z &= e^{-\frac{bt}{a}} \phi(v) + e^{-t(a-1)} \psi(v) + e^{-t(a-2)} \chi(v) \\ &= \phi\left(v - \frac{bt}{a}\right) + e^t \psi(v - t) + e^{2t} \chi(v - t) \\ &= \phi\left(yx^{-\frac{b}{a}}\right) + x\psi\left(\frac{y}{x}\right) + x^2 \chi\left(\frac{y}{x}\right). \end{aligned}$$

$$\text{Ex. 4. } x^n \frac{d^n z}{dx^n} + nx^{n-1}y \frac{d^n z}{dx^{n-1}dy} + \frac{n(n-1)}{1.2} x^{n-2}y^2 \frac{d^n z}{dx^{n-2}dy^2} + \&c. \\ + y^n \frac{d^n z}{dy^n} = 0. \quad \text{Here we have (Art. 70)}$$

$$\{d(d-1) \dots (d-n+1) + nd(d-1) \dots (d-n+2) d' \\ + \frac{n(n-1)}{1.2} d(d-1) \dots (d-n+3) d' (d'-1) + \&c.\} z = 0,$$

$$\text{or } (d+d')(d+d'-1) \dots (d+d'-n+1) z = 0,$$

as is easily seen by equating the coefficients of u^n in the product of the expansions of $(1+u)^d$, $(1+u)^{d'}$, and in the equivalent expansion of $(1+u)^{d+d'}$. Hence

$$z = e^{-td} f_0(v) + e^{-t(d'-1)} f_1(v) + \dots + e^{-t(d'-n+1)} f_{n-1}(v) \\ = f_0(v-t) + e^t f_1(v-t) + \dots + e^{(n-1)t} f_{n-1}(v-t) \\ = f_0\left(\frac{y}{x}\right) + x f_1\left(\frac{y}{x}\right) + \dots + x^{n-1} f_{n-1}\left(\frac{y}{x}\right).$$

$$\text{Ex. 5. } \left(x \frac{d}{dx} + y \frac{d}{dy}\right)^n z + p_1 \left(x \frac{d}{dx} + y \frac{d}{dy}\right)^{n-1} z + \dots + p_n z = 0.$$

$$\text{Since } \left(x \frac{d}{dx} + y \frac{d}{dy}\right)^n = (d+d')(d+d'-1) \dots (d+d'-n+1),$$

the equation may be transformed into

$$\{(d+d')^n + q_1(d+d')^{n-1} + \dots + q_n\} z = 0, \\ \text{or } (d+d'-a_1)(d+d'-a_2) \dots (d+d'-a_n) z = 0; \\ \therefore z = e^{a_1 t} e^{-td} f_1(v) + e^{a_2 t} e^{-td} f_2(v) + \dots + e^{a_n t} e^{-td} f_n(v) \\ = x^{a_1} f_1(v-t) + x^{a_2} f_2(v-t) + \&c. \\ = x^{a_1} f_1\left(\frac{y}{x}\right) + x^{a_2} f_2\left(\frac{y}{x}\right) + \dots + x^{a_n} f_n\left(\frac{y}{x}\right).$$

119. Other substitutions either for the dependent variable or for the independent variables, that may be sometimes employed, are the following.

Ex. 1. $P \frac{dz}{dx} + Q \frac{dz}{dy} = Rf(z)$ where P , Q , and R are functions of x and y .

Let v denote a function of z determined by the equation

$$\frac{dv}{dz} = \frac{1}{f(z)};$$

then the equation becomes

$$P \frac{dv}{dx} + Q \frac{dv}{dy} = R.$$

Thus let the equation be

$$(y^2 + mxy) \left(\frac{dz}{dx} - m \frac{dz}{dy} \right) = (max + by)(z^2 + 1),$$

then $v = \tan^{-1} z$, and the solution of

$$\frac{dv}{dx} - m \frac{dv}{dy} = \frac{max + by}{y^2 + mxy} = \frac{a}{y} - \frac{a-b}{y+mx}$$

$$\text{is } v = \phi(y + mx) - \frac{a}{m} \log y - \frac{(a-b)x}{y+mx};$$

$$\therefore z = \tan \left\{ \phi(y + mx) - \frac{a}{m} \log y - \frac{(a-b)x}{y+mx} \right\}.$$

$$\text{Ex. 2. } \frac{1}{x^{n-1}} \frac{dz}{dx} + \frac{m}{y^{n-1}} \frac{dz}{dy} = \frac{az}{x^n} + \frac{bz}{y^n} + \frac{cz}{(xy)^{\frac{n}{2}}},$$

$$\text{let } x^n = t, \quad y^n = v;$$

$$\therefore \frac{dz}{dt} + m \frac{dz}{dv} = \left(\frac{a}{nt} + \frac{b}{nv} + \frac{c}{n\sqrt{tv}} \right) z;$$

$$\therefore z = t^{\frac{a}{n}} \cdot v^{\frac{b}{n}} \cdot \left(\sqrt{t} + \sqrt{\frac{1}{m}v} \right)^{\frac{2c}{n\sqrt{m}}} \phi(v - mt);$$

$$\text{or } z = x^a y^{\frac{b}{m}} \left(x^{\frac{n}{2}} + \frac{1}{\sqrt{m}} y^{\frac{n}{2}} \right)^{\frac{2c}{n\sqrt{m}}} \phi(y^n - mx^n).$$

120. Since when operating relative to x we may treat y and the symbol $\frac{d}{dy}$ as simple constants, and *vice versa* when operating

relative to y , we may often solve a partial differential equation by integrating it with respect to one of the independent variables, and introducing arbitrary functions of the other instead of constants; and then effecting the operations indicated relative to the second variable, and interpreting the result in the usual way, which will be attended with no difficulty when only the exponential symbols $e^{a \frac{d}{dx}}$ or $e^{a \frac{d}{dy}}$ are involved.

Ex. 1. $\frac{d^2 z}{dx^2} - a^2 \frac{d^2 z}{dy^2} = 0$; this may be written

$$\frac{d^2 z}{dx^2} - (ad')^2 z = 0, \text{ or } \left(\frac{d}{dx} - ad' \right) \left(\frac{d}{dx} + ad' \right) z = 0;$$

$$\therefore z = e^{axd} \phi(y) + e^{-axd} \psi(y) = \phi(y + ax) + \psi(y - ax).$$

$$\text{Hence } x \frac{d^2 z}{dx^2} + 2 \frac{dz}{dx} = a^2 x \frac{d^2 z}{dy^2},$$

$$\text{which may be written } \frac{d^2 (xz)}{dx^2} - a^2 \frac{d^2 (xz)}{dy^2} = 0, \text{ gives}$$

$$xz = \phi(y + ax) + \psi(y - ax).$$

$$\text{Ex. 2. } x^4 \frac{d^2 z}{dx^2} - a^2 \frac{d^2 z}{dy^2} = 0, \text{ or } x^4 \frac{d^2 z}{dx^2} - (ad')^2 z = 0;$$

$$\therefore \frac{z}{x} = e^{\frac{a}{x}d} \phi(y) + e^{-\frac{a}{x}d} \psi(y) = \phi\left(y + \frac{a}{x}\right) + \psi\left(y - \frac{a}{x}\right). \\ (\text{Ex. 2, Art. 69.})$$

$$\text{Ex. 3. } \frac{d^2 z}{dx^2} + a \frac{d^2 z}{dx dy} - \frac{2z}{x^2} = 0,$$

$$\text{or } \frac{d^2 z}{dx^2} + ad' \cdot \frac{dz}{dx} - \frac{2z}{x^2} = 0;$$

$$\therefore z = \left(1 - \frac{2}{axd'}\right) \phi'(y) + \left(1 + \frac{2}{axd'}\right) e^{-axd} \psi'(y), (\text{Ex. 6, Art. 74})$$

$$= \phi'(y) - \frac{2}{ax} \phi(y) + \psi'(y - ax) + \frac{2}{ax} \psi(y - ax).$$

$$\text{Ex. 4. } \frac{d^2 z}{dx^2} - \frac{m(m+1)}{x^2} z - a^2 \frac{d^2 z}{dy^2} = 0,$$

$$\text{or } \frac{d^2 z}{dx^2} - \frac{m(m+1)}{x^2} z - (ad')^2 z = 0 ;$$

$$\therefore z x^{m+1} = \left(x^3 \frac{d}{dx} \right)^m \frac{e^{axd'} \phi(y) + e^{-axd'} \psi(y)}{x^{2m-1}}, \quad (\text{Ex. 4, Art. 74})$$

$$= \left(x^3 \frac{d}{dx} \right)^m \frac{\phi(y+ax) + \psi(y-ax)}{x^{2m-1}}.$$

$$\text{Ex. 5. } \frac{d^2 z}{dx^2} - \frac{2m}{x} \frac{dz}{dx} - a^2 \frac{d^2 z}{dy^2} = 0,$$

$$xz = \left(x^3 \frac{d}{dx} \right)^m \frac{\phi(y+ax) + \psi(y-ax)}{x^{2m-1}}. \quad (\text{Ex. 3, Art. 74.})$$

$$\text{Ex. 6. } x^2 \frac{d^2 z}{dx^2} - 2mx \frac{dz}{dx} + 2mz - q^2 x^2 \frac{d^2 z}{dy^2} = 0,$$

$$x^2 z = \left(x^3 \frac{d}{dx} \right)^m \frac{\phi(y+qx) + \psi(y-qx)}{x^{2m-2}}. \quad (\text{Ex. 2, Art. 74.})$$

If $m = -1$, in Ex. 4, we get the solution of

$$\frac{d^2(xz)}{dx^2} - \frac{2xz}{x^2} - q^2 \frac{d^2(xz)}{dy^2} = 0,$$

$$\text{or } x^2 \frac{d^2 z}{dx^2} + 2x \frac{dz}{dx} - 2z - q^2 x^2 \frac{d^2 z}{dy^2} = 0, \text{ viz.}$$

$$x^2 z = qx \{ \phi'(y+qx) - \psi'(y-qx) \} - \phi(y+qx) - \psi(y-qx).$$

$$\text{Ex. 7. } \frac{d^2 z}{dx dy} + \frac{1}{x+y} \left(\frac{dz}{dx} + \frac{dz}{dy} \right) - \frac{2z}{(x+y)^2} = 0.$$

Make $y+x=t$, $y-x=v$, then

$$t^2 \frac{d^2 z}{dt^2} + 2t \frac{dz}{dt} - 2z - t^2 \frac{d^2 z}{dv^2} = 0,$$

$$\therefore t^2 z = t\phi'(t+v) - t\psi'(v-t) - \phi(t+v) - \psi(v-t);$$

$$\therefore z(x+y)^2 = (x+y) \{ \phi'(2y) - \psi'(-2x) \} - \phi(2y) - \psi(-2x).$$

$$\text{Ex. 8. } \frac{d^2z}{dx^2} + \frac{2n}{x} \frac{dz}{dx} - a^2 \frac{d^2z}{dy^2} = 0. \quad (\text{Ex. 3. Art. 74.})$$

$$x^{2n}z = \frac{d}{dx} \left\{ \frac{1}{x} \left(x^3 \frac{d}{dx} \right)^n \frac{\phi(y+ax) + \psi(y-ax)}{x^{2n-1}} \right\}. \quad (\text{Art. 65.})$$

Hence if $n=1$, we get, as in Ex. 1,

$$xz = \phi(y+ax) + \psi(y-ax);$$

and if $n=2$, we find

$$x^3z = \phi(y+ax) + \psi(y-ax) - ax\phi'(y+ax) + ax\psi'(y-ax).$$

$$\text{Ex. 9. } x^3 \frac{d^2z}{dx^2} + qx^2 \frac{d^2z}{dx dy} - 6z = 0. \quad (\text{Ex. 6. Art. 74.})$$

$$z = \frac{12}{q^2x^3} \phi(y) - \frac{6}{qx} \phi'(y) + \phi''(y) \\ + \frac{12}{q^2x^3} \psi(y-qx) + \frac{6}{qx} \psi'(y-qx) + \psi''(y-qx).$$

$$\text{Ex. 10. } \left(x \frac{d}{dx} + y \frac{d}{dy} \right)^2 z - 2 \left(x \frac{d}{dx} + y \frac{d}{dy} \right) z + 2z = x^m y^n.$$

$$z = \frac{x^m y^n}{(m+n-1)(m+n-2)} + x\phi\left(\frac{y}{x}\right) + x^2\psi\left(\frac{y}{x}\right).$$

$$\text{Ex. 11. } f(x) \cdot \frac{d^2z}{dx^2} + \frac{1}{2}f'(x) \cdot \frac{dz}{dx} - a^2 \frac{d^2z}{dy^2} = 0, \quad (\text{Ex. 1. Art. 48})$$

$$z = \phi[y + a \int dx (fx)^{-\frac{1}{2}}] + \psi[y - a \int dx \{f(x)\}^{-\frac{1}{2}}],$$

where $f(x)$ denotes a given function of x .

$$\text{Ex. 12. } \frac{d^2z}{dx^2} - a^2 x^{-\frac{1}{2}} \frac{d^2z}{dy^2} = 0.$$

$$z = x^{\frac{1}{2}} \{ \phi'(y + 3ax^{\frac{1}{2}}) + \psi'(y - 3ax^{\frac{1}{2}}) \} \\ - \frac{1}{3a} \{ \phi(y + 3ax^{\frac{1}{2}}) - \psi(y - 3ax^{\frac{1}{2}}) \}.$$

$$\text{Ex. 13. } \frac{d^2z}{dx^2} + f(x) \cdot \frac{d^2z}{dx dy} - \frac{f(x)}{x} \frac{dz}{dy} = 0, \quad (\text{Ex. 2. Art. 48})$$

$$z = x \int dx \frac{1}{x^2} \phi[y - \int dx f(x)] + x\psi(y).$$

Partial Differential Equations of a higher degree than the first.

121. To integrate $F(x, y, z, p, q) = 0$, when it contains terms of more than one dimension in p and q .

In order that $dz = p dx + q dy$ may be a perfect differential, we must have

$$\frac{dp}{dy} + q \frac{dp}{dz} = \frac{dq}{dx} + p \frac{dq}{dz}, \quad (2).$$

If from the proposed equation we determine

$$q, \frac{dq}{dx}, \frac{dq}{dz},$$

equation (2) becomes an equation for the determination of p of the form

$$\frac{dp}{dx} + M \frac{dp}{dy} + N \frac{dp}{dz} = L,$$

the integration of which depends on the integration of one of the equations

$$dp - L dx = 0, \quad dy - M dx = 0, \quad dz - N dx = 0.$$

Let $p = f(x, y, z, a)$ be found from these equations, a being an arbitrary constant. This value of p and the corresponding value of q found from the proposed equation, being substituted in $dz = p dx + q dy$ render it an exact differential; and thus a value of z will be obtained involving two arbitrary constants a and b ; and this will consequently be the complete primitive. The general primitive may be obtained by putting $b = \psi(a)$, differentiating the equation with regard to a , and eliminating a . The result containing one arbitrary function is as general as any solution which the equation admits.

Ex. 1. $p^2 + q^2 = 1$,

$$q = \sqrt{1 - p^2}, \quad \frac{dq}{dx} = -\frac{p}{\sqrt{1 - p^2}} \frac{dp}{dx}, \quad \frac{dq}{dz} = -\frac{p}{\sqrt{1 - p^2}} \frac{dp}{dz};$$

$$\therefore \frac{dp}{dy} + \sqrt{1 - p^2} \frac{dp}{dz} + \frac{p}{\sqrt{1 - p^2}} \frac{dp}{dx} + \frac{p^2}{\sqrt{1 - p^2}} \frac{dp}{dz} = 0,$$

$$\text{or } \frac{dp}{dz} + p \cdot \frac{dp}{dx} + \sqrt{1 - p^2} \frac{dp}{dy} = 0;$$

which is integrable if we can integrate the system of equations

$$dp = 0, \quad pdz - dx = 0, \quad \sqrt{1-p^2} \cdot dz - dy = 0.$$

The first gives $p = a$, whence $q = \sqrt{1-a^2}$,

$$\therefore dz = adx + \sqrt{1-a^2} dy,$$

$$\text{or } z = ax + \sqrt{1-a^2} y + \phi(a).$$

Differentiating this with respect to a , we find

$$0 = x - \frac{a}{\sqrt{1-a^2}} y + \phi'(a);$$

between which and the preceding we may eliminate a , when the form of the function $\phi(a)$ is assigned. This is a simple case of the Problem of finding surfaces of equivalent area to a given surface, that is, such that any cylindrical surface parallel to the axis of z , may always intercept equal areas in the required and given surface. If P, Q denote the values of $\frac{dz}{dx}, \frac{dz}{dy}$ in the given surface, the equation of condition is

$$p^2 + q^2 = P^2 + Q^2;$$

which, if the given surface be a plane, leads to the equation of this example.

Ex. 2. $p^m q^n = c^n.$

The equation in p to be integrated is

$$\frac{dp}{dx} + \frac{n}{mc} p^{\frac{m}{n}+1} \frac{dp}{dy} + \left(\frac{n}{m} + 1\right) p \frac{dp}{dz} = 0;$$

$$\therefore dp = 0, \quad \text{and } p = a^n, \quad \therefore q = ca^{-m},$$

$$\therefore dz = a^n dx + \frac{c}{a^m} dy,$$

$$\text{and } z = a^n x + \frac{c}{a^m} y + \phi(a),$$

$$0 = na^{n-1}x - \frac{mc}{a^{m+1}}y + \phi'(a);$$

and it remains to eliminate a between these equations when the form of $\phi(a)$ is assigned.

If the proposed equation be

$$p^m q^n = c^n x^\alpha y^\beta z^\gamma,$$

and we take new independent variables x' and y' such that

$$\frac{dx'}{dx} = x^{\frac{\alpha}{m}}, \quad \frac{dy'}{dy} = y^{\frac{\beta}{n}}.$$

$$\text{Then} \quad \left(\frac{dz}{dx}\right)^m \cdot \left(\frac{dz}{dy}\right)^n = c^n z^\gamma.$$

Now assume z' such that $\frac{dz'}{dz} = z^{\frac{-\gamma}{m+n}}$, then the equation becomes

$$\left(\frac{dz'}{dx'}\right)^m \cdot \left(\frac{dz'}{dy'}\right)^n = c^n,$$

and is reduced to the preceding.

$$\text{Ex. 3.} \quad z = pq;$$

here, when the form of $\phi(a)$ is assigned, a must be eliminated between the equations,

$$\frac{z}{y+a} = x + \phi(a), \quad \frac{z}{(y+a)^2} = -\phi'(a).$$

Thus if $\phi(a) = \frac{c}{a}$, then it will be found that $z = (c + \sqrt{xy})^2$.

Non-linear Equations of the Second and Higher Orders.

122. Here, besides the coefficients p and q of the first order which may be involved together with x , y , and z , the equation must contain one or more of the coefficients of the second order r , s , t ; so that in its most general form it will be

$$F(x, y, z, p, q, r, s, t) = 0.$$

In partial differential equations of the second order, we cannot be certain of the form of the solution, nor pronounce before-

hand how many arbitrary functions it ought to contain. For let

$$u = f\{x, y, z, \phi(v), \psi(w)\} = 0,$$

be an equation containing two arbitrary functions of v and w two known functions of x, y and z ; then p and q will be given by the equations

$$\frac{du}{dx} = 0, \quad \frac{du}{dy} = 0;$$

and r, s , and t by the equations

$$\frac{d^2u}{dx^2} = 0, \quad \frac{dd}{dx dy} u = 0, \quad \frac{d^2u}{dy^2} = 0.$$

These together with $u = 0$ make six equations, into which the six quantities

$$\phi(v), \quad \psi(w), \quad \phi'(v), \quad \psi'(w), \quad \phi''(v), \quad \psi''(w),$$

may enter. Consequently, it will not generally be possible to eliminate these six quantities and obtain a relation between x, y, z, p, q, r, s, t , independent of the forms of the functions ϕ and ψ ; although particular cases do occur in which this can be effected.

In general, if $u = 0$ contain n independent variables and m functions of the form $\phi(v)$, where v is a determinate function of the variables; these functions can certainly be eliminated between the equations obtained after r differentiations, if

$$1 + n + \frac{n(n+1)}{1 \cdot 2} + \dots + \frac{n(n+1) \dots (n+r-1)}{r!} > m(r+1),$$

the former quantity expressing the number of equations, and the latter the number of functions.

123. As an example of forming a partial differential equation of the second order, and of the sort we are now considering, by the elimination of two arbitrary functions, we may take

$$\begin{aligned} u &= y - x\phi(z) - \psi(z) = 0; \\ \therefore \frac{du}{dx} &= -\phi(z) - x\phi'(z)p - \psi'(z)p = 0, \\ \frac{du}{dy} &= 1 - x\phi'(z)q - \psi'(z)q = 0; \end{aligned}$$

$$\begin{aligned}
&\therefore p + q\phi(z) = 0; \\
&\therefore r + s\phi(z) + pq\phi'(z) = 0, \\
&\quad s + t\phi(z) + q^2\phi'(z) = 0; \\
&\therefore qr - ps + (qs - pt)\phi(z) = 0, \\
&\quad \text{or } q^2r - 2pqs + p^2t = 0.
\end{aligned}$$

124. The equations

$$\frac{d^2z}{dx^2} + P \frac{dz}{dx} + Q = 0,$$

$$\frac{d^2z}{dy^2} + P \frac{dz}{dy} + Q = 0,$$

where P and Q are functions of x , y and z , must be integrated as equations between two variables, y being regarded as constant in the former, and x in the latter; and arbitrary functions of those variables respectively, being introduced instead of constants.

$$\begin{aligned}
\text{Ex. } (x - x^2) \frac{d^2z}{dx^2} + (y - xy - 2x) \frac{dz}{dx} - yz &= 0, \\
(1 - x)z &= x^{1-y} \phi(y) + \psi(y).
\end{aligned}$$

125. The equations

$$\frac{dd}{dx dy} z + P \frac{dz}{dx} = Q,$$

$$\frac{dd}{dx dy} z + P \frac{dz}{dy} = Q,$$

where P and Q do not contain z , are reducible to the case of Art. 105, by considering $\frac{dz}{dx}$ or $\frac{dz}{dy}$ respectively as a simple quantity v .

$$\text{Ex. } \frac{dd}{dx dy} z + \frac{y}{1-y^2} \frac{dz}{dx} = ay^3.$$

This being a linear equation in $\frac{dz}{dx}$ which is made integrable by the factor $\frac{1}{\sqrt{1-y^2}}$, we have

$$\frac{1}{\sqrt{1-y^2}} \frac{dz}{dx} = a \int dy \frac{y^3}{\sqrt{1-y^2}} = -a \{y^2 \sqrt{1-y^2} + \frac{2}{3}(1-y^2)^{3/2}\} + \psi'(x);$$

$$\therefore z = \{\psi(x) + \phi(y)\} \sqrt{1-y^2} - \frac{ax}{3} (2+y^2)(1-y^2).$$

126. Similarly, equations of the forms

$$f\left(x, y, z, \frac{dz}{dx}, \frac{d^2z}{dx^2}, \&c., \frac{d^nz}{dx^n}\right) = 0,$$

$$f\left(x, y, z, \frac{dz}{dy}, \frac{d^2z}{dy^2}, \&c., \frac{d^nz}{dy^n}\right) = 0,$$

may be treated as if they contained only two variables; arbitrary functions of y instead of constants being introduced into the solution of the first, and arbitrary functions of x into the solution of the second. To this case may also be reduced the equation

$$f\left(x, y, \frac{d^nz}{dy^n}, \frac{dd^nz}{dx dy^n}, \frac{d^2d^nz}{dx^2 dy^n}, \&c., \frac{d^m d^nz}{dx^m dy^n}\right) = 0;$$

for by putting $\frac{d^nz}{dy^n} = v$, it becomes

$$f\left(x, y, v, \frac{dv}{dx}, \frac{d^2v}{dx^2}, \&c., \frac{d^mv}{dx^m}\right) = 0,$$

which will give a value of v containing m arbitrary functions of y ; and then $\frac{d^nz}{dy^n} = v$ will give z involving n arbitrary functions of x .

127. To integrate the equation of the second order,

$$Rr + Ss + Tt = V,$$

where R, S, T, V are functions of x, y, z, p, q . The following process first given by Monge, may be frequently applied.

By means of the relations

$$dp = rdx + sdy, \quad dq = sdx + tdy,$$

eliminating two of the three coefficients, r and t , from the proposed, we get

$$Rdpdy + Tdqdx - Vdxdy = s\{R(dy)^2 + T(dx)^2 - Sdxdy\},$$

which is satisfied by

$$\left. \begin{aligned} Rdpdy + Tdqdx - Vdxdy &= 0 \\ R(dy)^2 + T(dx)^2 - Sdxdy &= 0 \end{aligned} \right\} \quad (1).$$

Let $M = a$, $N = b$, be two relations between x, y, z, p, q , and the arbitrary constants a, b , which satisfy these equations; then $M = \phi(N)$ satisfies the proposed equation. This will be shewn by proving that it can reproduce the proposed equation.

$$\text{Let } dy = m dx; \quad \therefore Rm^2 - Sm + T = 0.$$

For each root of this equation, we have

$$\left. \begin{aligned} dy - m dx &= 0, \quad Rmdp + Tdq - Vmdx = 0; \\ \therefore dy &= m dx, \\ dq &= \frac{Vm}{T} dx - \frac{Rm}{T} dp, \\ dz &= p dx + q dy. \end{aligned} \right\} \quad (2)$$

Hence $M = a$ gives on differentiation

$$\begin{aligned} 0 &= \frac{dM}{dx} \cdot dx + \frac{dM}{dy} \cdot m dx + \frac{dM}{dz} (p dx + q m dx) \\ &\quad + \frac{dM}{dp} \cdot dp + \frac{dM}{dq} \cdot \left(\frac{Vm}{T} dx - \frac{Rm}{T} dp \right), \end{aligned}$$

wherein all the known relations (2) having been introduced, dx and dp must be independent,

$$\begin{aligned} \therefore 0 &= \frac{dM}{dx} + m \frac{dM}{dy} + \frac{dM}{dz} \cdot (p + m q) + \frac{Vm}{T} \frac{dM}{dq}, \\ 0 &= \frac{dM}{dp} - \frac{Rm}{T} \frac{dM}{dq}; \\ \therefore \frac{dM}{dx} &= - \left\{ m \frac{dM}{dy} + (p + m q) \frac{dM}{dz} + \frac{Vm}{T} \frac{dM}{dq} \right\}. \\ \frac{dM}{dp} &= \frac{Rm}{T} \frac{dM}{dq}. \end{aligned}$$

$$\text{So } \frac{dN}{dx} = - \left\{ m \frac{dN}{dy} + (p + mq) \frac{dN}{dz} + \frac{Vm}{T} \frac{dN}{dq} \right\},$$

$$\frac{dN}{dp} = \frac{Rm}{T} \frac{dN}{dq}.$$

By differentiating the assumed equation $M = \phi(N)$ we have

$$dM = \phi'(N) \cdot dN.$$

Now

$$\begin{aligned} dM &= - \left\{ m \frac{dM}{dy} + (p + mq) \frac{dM}{dz} + \frac{Vm}{T} \frac{dM}{dq} \right\} dx + \frac{dM}{dy} dy \\ &\quad + \frac{dM}{dz} (pdx + qdy) + \frac{Rm}{T} \frac{dM}{dq} \cdot dp + \frac{dM}{dq} \cdot dq \\ &= \left(\frac{dM}{dy} + q \frac{dM}{dz} \right) (dy - m dx) + \frac{1}{T} \frac{dM}{dq} (Rmdp + Tdq - Vmdx), \end{aligned}$$

and a similar value exists for dN ;

$$\begin{aligned} \therefore \left(\frac{dM}{dy} + q \frac{dM}{dz} \right) (dy - m dx) + \frac{1}{T} \frac{dM}{dq} (Rmdp + Tdq - Vmdx) \\ = \phi'(N) \left\{ \left(\frac{dN}{dy} + q \frac{dN}{dz} \right) (dy - m dx) \right. \\ \left. + \frac{1}{T} \frac{dN}{dq} (Rmdp + Tdq - Vmdx) \right\}, \end{aligned}$$

which may be put under the form

$$Rmdp + Tdq - Vmdx = \omega (dy - m dx),$$

$$\text{or } Rm (rdx + sdy) + T(sdx + tdy) - Vmdx = \omega (dy - m dx),$$

where dx and dy are independent ;

$$\therefore Rmr + Ts - Vm = -\omega m,$$

$$Rms + Tt = \omega ;$$

$$\therefore Rr + Ss + Tt = V - \frac{s}{m} (Rm^2 - Sm + T) = V.$$

Hence, $M = \phi(N)$ satisfies the proposed equation.

According as the roots of

$$Rm^2 - Sm + T = 0,$$

are unequal or equal, we are thus supplied with a total or partial differential equation for the determination of z .

As the reducing equations (1) may contain

$$x, y, z, p, q,$$

and as these together with $dz = p dx + q dy$ will generally lead to an equation containing three variables, which will not always admit of a single primitive (Art. 98), it may happen that the first integral of the proposed equation cannot be determined; but we must not thence conclude that the proposed equation does not admit of being solved.

128. Hence, to integrate the equation of the second order

$$Rr + Ss + Tt = V,$$

the process is to obtain a value of m from the equation

$$Rm^2 - Sm + T = 0,$$

to substitute it in the system

$$dy - m dx = 0, \quad Rm dp + T dq = Vm dx, \quad (1)$$

to satisfy these conjointly or separately, by two relations between x, y, z, p, q ,

$$M = a, \quad N = b,$$

then to put $M = \phi(N)$ which will be a first integral of the proposed, and to integrate this equation of the first order. But this determination of the second integral from the first will often be attended with great difficulty, on account of its involving an arbitrary function; and therefore when possible it is often more convenient to find from the second value of m another first integral of the form $M' = \psi(N')$, and between these to eliminate p or q , so as to obtain an equation involving only one differential coefficient, and which is therefore easily integrated.

129. If R, S, T be constant, and V a function of x and y only, then the values of m will be numerical, m and n suppose; and the integrals of equations (1) will be

$$y - mx = a, \quad Rmp + Tq = m \int dx V + b,$$

where, previous to integration, $mx + a$ is substituted for y in V , and after integration the value of a , viz. $y - mx$, is restored: consequently calling this value V_1 , since $Rmn = T$, we have a first integral of the proposed, viz.

$$Rp + Rnq = V_1 + \phi'(y - mx).$$

Next to integrate this equation of the first order, we have the reducing equations

$$\begin{aligned} dy - ndx &= 0, \\ Rdz - \{V_1 + \phi'(y - mx)\} dx &= 0; \\ \therefore y - nx &= a, \\ Rz = \int dx V_1 + \int dx \phi'(y - mx) + \beta, \end{aligned}$$

$nx + a$ being substituted for y before the integration is performed, and afterwards the value of a , $y - nx$, restored; this gives $\int dx V_1 = V_2$, suppose,

$$\text{and } \int dx \phi'(y - mx) = \int dx \phi' \{(n - m)x + a\} = \frac{1}{n - m} \phi(y - mx);$$

hence, including the constant multiplier under the sign of the function, we have the complete integral required,

$$Rz = V_2 + \phi(y - mx) + \psi(y - nx);$$

which agrees with the result obtained at once by separation of symbols from the proposed when put under the form

$$R \left(\frac{d}{dx} + m \frac{d}{dy} \right) \left(\frac{d}{dx} + n \frac{d}{dy} \right) z = f(x, y).$$

$$\text{Ex. 1. } \frac{d^2 z}{dx^2} - c^2 \frac{d^2 z}{dy^2} = 0, \text{ or } r - c^2 t = 0.$$

The two systems of reducing equations here are

$$\left. \begin{aligned} dy + cdx &= 0 \\ dp + cdq &= 0 \end{aligned} \right\}, \quad \left. \begin{aligned} dy - cdx &= 0 \\ dp - cdq &= 0 \end{aligned} \right\};$$

from the former we get

$$y + cx = a, \quad p + cq = b; \quad \therefore p + cq = 2c\phi'(y + cx);$$

similarly the latter gives $p - cq = -2c\psi'(y - cx)$;

hence, eliminating q , we find

$$\begin{aligned} p = \frac{dz}{dx} &= c\phi'(y + cx) - c\psi'(y - cx), \\ \therefore z &= \phi(y + cx) + \psi(y - cx). \end{aligned}$$

Ex. 2. $r - t + \frac{4p}{x+y} = 0.$

Here the auxiliary equations are

$$\begin{aligned} dy - dx &= 0, & dp - dq + \frac{4p}{x+y} dx &= 0, \\ dy + dx &= 0, & dp + dq - \frac{4p}{x+y} dx &= 0 \quad (1). \end{aligned}$$

From the former we find

$$y - x = a, \text{ and therefore } dp - dq + \frac{4pdx}{2y-a} = 0,$$

$$\text{or } (dp - dq)(2y - a) + (p - q)2dy + 2dz = 0,$$

$$\text{since } dz = pdx + qdy, \text{ and } y - x = a;$$

$$\therefore (2y - a)(p - q) + 2z = b = f(y - x),$$

$$\text{or } p - q + \frac{2z}{x+y} = \frac{f(y-x)}{x+y}.$$

But from the first of equations (1) we have $y + x = a'$;

$$\therefore \frac{dz}{dy} - \frac{dz}{dx} - \frac{2z}{a'} = -\frac{f(y-x)}{a'};$$

which, being a linear equation, may be integrated, and we get (Art. 113)

$$z = -e^{\frac{2y}{x+y}} \int dy e^{-\frac{2y}{x+y}} \frac{f(y-x)}{a'} + e^{\frac{2y}{x+y}} \phi(x+y);$$

where, after the integration, a' is to be replaced by $x + y$.

Ex. 3. $r - \frac{2p}{q}s + \frac{p^2}{q^2}t = 0,$

$$\text{or } rdx dy - \frac{2p}{q}sdx dy + \frac{p^2}{q^2}tdy dx = 0,$$

or, eliminating r and t ,

$$dpdy - s(dy)^2 - \frac{2p}{q}sdx dy + \frac{p^2}{q^2}\{dqdx - s(dx)^2\} = 0;$$

this resolves itself into

$$(dy)^2 + 2\frac{p}{q}dydx + \frac{p^2}{q^2}(dx)^2 = 0, \quad dpdy + \frac{p^2}{q^2}dqdx = 0;$$

whence arise the auxiliary equations

$$dy = -\frac{p}{q} dx, \quad dp - \frac{p}{q} dq = 0;$$

$$\text{but } dz = p dx + q dy = 0, \quad \therefore z = a,$$

$$\text{and } \frac{q dp - p dq}{q^2} = 0 \text{ gives } \frac{p}{q} = b;$$

$$\therefore p - qf(z) = 0, \text{ the first integral.}$$

To integrate this equation of the first order, we have

$$dz = 0, \quad \text{or } z = \alpha,$$

$$dy + f(z) dx = 0, \quad \text{or } y + xf(\alpha) = \beta;$$

$$\therefore y + xf(z) = \phi(z);$$

this is the equation to the surface generated by a straight line, subjected to pass through three given fixed curves.

$$\text{Ex. 4. } (1 + pq + q^2) r + (q^2 - p^2) s - (1 + pq + p^2) t = 0,$$

$$\text{or } (1 + q\alpha) r + (q - p) as - (1 + p\alpha) t = 0,$$

putting $p + q = \alpha$. The values of m are to be obtained from the equation

$$(1 + q\alpha) m^2 - (q - p) \alpha m - 1 - p\alpha = 0,$$

which gives

$$m = -\frac{1 + p\alpha}{1 + q\alpha}, \quad m = 1;$$

and the two systems of auxiliary equations answering to these values of m , are

$$dy - dx = 0, \quad (1 + q\alpha) dp - (1 + p\alpha) dq = 0,$$

$$(1 + q\alpha) dy + (1 + p\alpha) dx = 0, \quad dp + dq = 0. \quad (1)$$

The latter gives $p + q = b$, or $\alpha = b$,

$$x + y + bz = a; \quad \therefore x + y + (p + q)z = \phi(p + q).$$

The former system gives $y - x = a_1$; and if we assume $p - q = \beta$, the second equation of this system becomes

$$(\alpha^2 + 2) d\beta - \alpha\beta d\alpha = 0, \quad \text{whence } \beta = b_1 \cdot (\alpha^2 + 2)^{\frac{1}{2}};$$

$$\therefore p - q = \{(p + q)^2 + 2\}^{\frac{1}{2}} \psi(y - x).$$

Next to integrate this equation of the first order, since

$$p = \frac{1}{2}(\alpha + \beta), \quad q = \frac{1}{2}(\alpha - \beta),$$

we have

$$\begin{aligned} dz &= \frac{1}{2}\alpha(dx + dy) + \frac{1}{2}\beta(dx - dy) \\ &= \frac{1}{2}\alpha(dx + dy) + \frac{1}{2}(dx - dy)(\alpha^2 + 2)^{\frac{1}{2}}\psi(y - x), \end{aligned}$$

this is integrable if we suppose α to be constant, and gives

$$z + \phi(\alpha) = \frac{1}{2}\alpha \cdot (x + y) + (\alpha^2 + 2)^{\frac{1}{2}}\psi_1(y - x);$$

which, combined with

$$\phi'(\alpha) = \frac{1}{2}(x + y) + \frac{\alpha}{\sqrt{\alpha^2 + 2}} \cdot \psi_1(y - x),$$

represents the integral of the proposed equation.

Integration by a Series.

When other methods fail, partial differential equations, like equations between two variables, can be sometimes integrated by a series; and we shall now give an example taken from Euler of the process.

130. To integrate by a series the equation

$$bz + a(x + y)(p + q) + (x + y)^2 s = 0.$$

Let $z = A(x + y)^m \phi(x) + B(x + y)^{m+1} \phi'(x) + C(x + y)^{m+2} \phi''(x) + \&c.$;

and as the integral must be symmetrical with respect to x and y , we must, after the determination of m , B , C , &c., replace $\phi(x)$, $\phi'(x)$, &c. by $\phi(x) + \psi(y)$, $\phi'(x) + \psi'(y)$, &c. Obtaining the values of p , q , s , and substituting them along with the assumed value of z in the proposed equation, we find

$$\begin{aligned} 0 &= \{b + 2ma + m(m - 1)\} A(x + y)^m \phi(x) \\ &\quad + \{bB + 2(m + 1)aB + aA + (m + 1)mB + mA\} (x + y)^{m+1} \phi'(x) \\ &\quad + \{bC + 2(m + 2)aC + aB + (m + 2)(m + 1)C + (m + 1)B\} (x + y)^{m+2} \phi''(x) \\ &\quad + \&c. \end{aligned}$$

which gives for the determination of m and B the relations

$$b + 2ma + m(m-1) = 0,$$

$$b + 2(m+1)a + (m+1)m = -(a+m) \frac{A}{B} \quad (1);$$

or, subtracting the former from the latter,

$$2a + 2m = -(a+m) \frac{A}{B}, \quad \therefore B = -\frac{1}{2}A.$$

For the determination of C we have

$$b + 2(m+2)a + (m+2)(m+1) = -(a+m+1) \frac{B}{C} = \frac{1}{2}(a+m+1) \frac{A}{C};$$

or, subtracting equation (1) from this,

$$2a + 2m + 2 = \frac{1}{2}(a+m+1) \frac{A}{C} - 2(a+m);$$

$$\therefore C = \frac{a+m+1}{2 \cdot 2 \cdot (2a+2m+1)} A = \frac{(i-1)A}{2 \cdot 2(2i-1)} \text{ putting } a+m = -i;$$

$$\text{similarly } D = \frac{-(i-2)A}{2^2 \cdot 2 \cdot 3(2i-1)}, \quad E = \frac{(i-2)(i-3)A}{2^3 \cdot 2 \cdot 3 \cdot 4(2i-1)(2i-3)}, \text{ \&c.}$$

The series will terminate whenever $a+m+n = n-i$ is a whole number, or when $b = (a+i)(a-i-1)$. Thus let $m = -a$, and therefore $b = a(a-1)$; it is evident, from the formation of the series, that $B = C = \&c. = 0$;

$$\therefore z = (x+y)^{-a} \{\phi(x) + \psi(y)\}.$$

Let $m = -a-2$, and therefore $b = (a+2)(a-3)$, then

$$z = \frac{\phi(x) + \psi(y)}{(x+y)^{a+2}} - \frac{1}{2} \frac{\phi'(x) + \psi'(y)}{(x+y)^{a+1}} + \frac{1}{12} \frac{\phi''(x) + \psi''(y)}{(x+y)^a}.$$

If $b = 0$, the equation becomes

$$a(p+q) + (x+y)s = 0;$$

and we must have $a = -i$ or $= i+1$, and therefore $m = 0$, or $m = -2i-1$; in the former case

$$z = \phi(x) + \psi(y) - \frac{1}{2}(x+y)\{\phi'(x) + \psi'(y)\} \\ + \frac{1}{2} \frac{(i-1)(x+y)^2}{2(2i-1)} \{\phi''(x) + \psi''(y)\} - \&c.;$$

in the latter, we have $(x+y)^{2+i}z$ equal to the same series.

To this form may likewise be reduced the equation (analogous to that of Riccati and integrable in the same cases)

$$(2m-1)^2 \frac{d^2 z}{dx^2} = c^2 x^{\frac{4m}{1-2m}} \frac{d^2 z}{dy^2},$$

by introducing new independent variables,

$$u = \frac{1}{2} x^{\frac{1}{1-2m}} + \frac{y}{2c}, \quad v = \frac{1}{2} x^{\frac{1}{1-2m}} - \frac{y}{2c};$$

when it becomes $(u+v) \frac{d^2 z}{du dv} + m \frac{dz}{du} + m \frac{dz}{dv} = 0$; and hence z

may be found in a series.

But when m is a positive integer, the solution may be readily obtained by assuming $x = t^{\pm 2m}$; then

$$(2m \pm 1)^2 \frac{d^2 z}{dx^2} = c^2 x^{\frac{-4m}{2m \pm 1}} \frac{d^2 z}{dy^2}$$

is transformed into

$$\frac{dz^2}{dt^2} + \frac{2m}{t} \frac{dz}{dt} = c^2 \frac{d^2 z}{dy^2}.$$

Hence, for the upper sign, we get (Ex. 5. Art. 120),

$$zt = \left(t^3 \frac{d}{dt} \right)^m \frac{\phi(y+ct) + \psi(y-ct)}{t^{2m-1}};$$

where, after performance of the operations indicated, we must replace t by $x^{\frac{1}{1+2m}}$; and for the lower sign the solution is (Art. 65)

$$zt^{2m} = \frac{d}{dt} \left\{ \frac{1}{t} \left(t^3 \frac{d}{dt} \right)^m \frac{\phi(y+ct) + \psi(y-ct)}{t^{2m-1}} \right\};$$

where, after performing the operations indicated, we must replace t by $x^{\frac{1}{1-2m}}$. The latter result is obtained from observing that, if u be a solution of $\frac{d^2 y}{dx^2} - \frac{2m}{x} \frac{dy}{dx} - c^2 y = 0$, then $x^{-2m} \frac{du}{dx}$ is a solution of $\frac{d^2 y}{dx^2} + \frac{2m}{x} \frac{dy}{dx} - c^2 y = 0$.

Simultaneous Equations.

131. A system of partial differential equations which involve two dependent variables z and u , provided they are linear and have constant coefficients, may be treated in the same manner as simultaneous differential equations of the same description containing only one independent variable. For the two symbols of differentiation $\frac{d}{dx}$ and $\frac{d}{dy}$, since they do not affect each other, and obey in their combinations the same laws as ordinary algebraical quantities, may be treated as two independent constants.

$$\begin{aligned}\text{Ex. 1.} \quad \frac{dz}{dx} + a \frac{dz}{dy} + bz + c \frac{du}{dx} &= 0, \\ \frac{du}{dx} + a \frac{du}{dy} + bu + c' \frac{dz}{dx} &= 0.\end{aligned}$$

Separating the symbols, and for convenience writing d and d' instead of $\frac{d}{dx}$ and $\frac{d}{dy}$, these become

$$(d + ad' + b)z + cdu = 0, \quad (d + ad' + b)u + c'dz = 0.$$

Now substitute in the former the value of u obtained from the latter, regarding d and d' as ordinary constants, and we get

$$\{(d + ad' + b)^2 - cc'd^2\}z = 0;$$

or, if we denote $1 - \sqrt{cc'}$ by $\frac{1}{m}$ and $1 + \sqrt{cc'}$ by $\frac{1}{n}$,

$$\{d + m(ad' + b)\} \{d + n(ad' + b)\} z = 0,$$

the integral of which is

$$z = e^{-mbx} \phi(y - amx) + e^{-nbx} \psi(y - anx),$$

where m and n have the values above-written; and from this u can be found.

$$\begin{aligned}\text{Ex. 2.} \quad \frac{d^2z}{dx dy} + a \frac{du}{dy} &= 0, \quad \frac{d^2u}{dx dy} + b \frac{dz}{dx} = 0; \\ \therefore \frac{d^3z}{dx^2 dy} + a \frac{d^2u}{dx dy} &= 0, \quad \text{or} \quad \frac{d^3z}{dx^2 dy} - ab \frac{dz}{dx} = 0; \\ \therefore z &= f(y) + x\phi(y) + e^{aby} \psi(x).\end{aligned}$$

132. If $U = f(x, y, z, p, q) = 0$ be a partial differential equation of the first order, the singular solution if there be one, as in the case of two variables, will be found by eliminating p and q between the three equations

$$U = 0, \quad \frac{dU}{dp} = 0, \quad \frac{dU}{dq} = 0.$$

Ex. $(z - px - qy)^2 - a^2(1 + p^2 + q^2) = 0$, which expresses that the surface represented by the complete integral has the perpendicular from the origin on the tangent plane of a constant length. Here

$$\frac{dU}{dp} = -(z - px - qy)x - a^2p = 0,$$

$$\frac{dU}{dq} = -(z - px - qy)y - a^2q = 0;$$

eliminating p and q we find

$$x^2 + y^2 + z^2 = a^2.$$

133. We shall terminate this part of the subject by the following geometrical problems; the first of them being the well-known solution by Monge of the Problem—to find a surface at every point of which the radii of curvature are equal and of the same sign. The conditions for this are expressed by the equations:

$$\text{Ex. 1. } \frac{p}{1+p^2} \frac{dp}{dx} = \frac{1}{q} \frac{dq}{dx}; \quad \frac{q}{1+q^2} \frac{dq}{dx} = \frac{1}{p} \frac{dp}{dy}.$$

Integrating these, and replacing the arbitrary constant by Y a function of y in the former, and by X a function of x in the latter, we find

$$1 + p^2 = q^2 Y, \quad 1 + q^2 = p^2 X;$$

$$\therefore p = \sqrt{\frac{Y+1}{XY-1}}, \quad q = \sqrt{\frac{X+1}{XY-1}} \dots\dots\dots (1).$$

But as the object is to find z a function of x and y which shall satisfy the two proposed equations, the quantities p and q must by their nature satisfy the condition $\frac{dp}{dy} = \frac{dq}{dx}$, which becomes

$$\frac{1}{(1+X)^{\frac{1}{2}}} \frac{dX}{dx} = \frac{1}{(1+Y)^{\frac{1}{2}}} \frac{dY}{dy},$$

and is of the form $\phi(x) = \psi(y)$, whatever be the functions X and Y ; it cannot therefore subsist for all values of x and y which are variables independent of one another, unless each member is reduced to the same arbitrary constant, $\frac{1}{2C}$ suppose; we then have

$$\frac{2C}{(1+X)^{\frac{1}{2}}} \frac{dX}{dx} = 1, \quad \frac{2C}{(1+Y)^{\frac{1}{2}}} \frac{dY}{dy} = 1,$$

which give by integration, a and b being two new arbitrary constants,

$$\frac{C}{\sqrt{1+X}} = a - x, \quad \frac{C}{\sqrt{1+Y}} = b - y.$$

Hence the quantities X and Y become known, and then p and q may be expressed in terms of x and y from equations (1), and substituting these values of p and q in $dz = p dx + q dy$ and integrating, we get

$$(x-a)^2 + (y-b)^2 + (z-c)^2 = C^2.$$

Ex. 2. To find the surface which cuts at right angles the surfaces represented by the equation $xy + xz + yz = a^2$, when a assumes all values.

Here $p = -\frac{y+z}{x+y}$, $q = -\frac{x+z}{x+y}$, and the equation of condition is

$$1 + p \frac{dz}{dx} + q \frac{dz}{dy} = 0;$$

$$\therefore (y+z) \frac{dz}{dx} + (x+z) \frac{dz}{dy} = x+y;$$

the integral of which is

$$(x+y+z) \cdot (y-z)^2 = \phi \{ (x+y+z) \cdot (x-z)^2 \}.$$

Ex. 3. To find the surface which cuts at right angles all spheres that touch a plane in a given point; taking the given point for the origin, and the plane for that of xy , the equation is

$$2rz - z^2 = x^2 + y^2;$$

$$\therefore p = \frac{x}{r-z}, \quad q = \frac{y}{r-z};$$

and the equation of condition is

$$x \frac{dz}{dx} + y \frac{dz}{dy} = z - r,$$

$$\text{or } 2xz \frac{dz}{dx} + 2yz \frac{dz}{dy} = z^2 - x^2 - y^2.$$

$$\text{Hence } ydx - xdy = 0, \quad \text{or } \frac{x}{y} = a,$$

$$2yzdz - (z^2 - x^2 - y^2) dy = 0,$$

$$\text{or } \frac{2zdz}{y} - \frac{z^2 dy}{y^2} + (1 + a^2) dy = 0;$$

$$\therefore \frac{z^2}{y} + (1 + a^2) y = b; \quad \therefore z^2 + x^2 + y^2 = y \phi \left(\frac{x}{y} \right).$$

If the required surface is to be of the second order, we must have

$$\phi \left(\frac{x}{y} \right) = b \frac{x}{y} + c, \quad \therefore z^2 + x^2 + y^2 = bx + cy,$$

which represents spheres passing through the given point, and having their centres in the given plane.








By integrating the auxiliary equations

$$xdy - ydx = 0, \quad dx(r - z) + xdz = 0,$$

$$\text{we get} \quad x(r - z) = \phi \left(\frac{y}{x} \right).$$

And if we wish to determine the arbitrary function so that when $y = ax^2$, z shall equal $r + bx$, then

$$x(z - r) = \frac{b}{a^2} \left(\frac{y}{x} \right)^2.$$



A TREATISE

ON THE

CALCULUS OF FINITE DIFFERENCES.

SECTION I.

DIRECT METHOD OF DIFFERENCES.

Definitions and Principles.

ART. 1. The Difference of a Function of one or more variables is the result obtained by subtracting from one another the two values of the function that arise from giving to the variables contained in it, different assigned values.

Thus if u_x denote any function of a variable x , and u_{x+h} the same function of $x + h$, and if the value of u_x be subtracted from that of u_{x+h} , the result is called the *Difference* of u_x ; and the quantity h is called the *increment* of the principal variable x . In the Differential Calculus it is the first term only of the series, arranged according to ascending powers of h , expressing $u_{x+h} - u_x$, or rather the coefficient of h in that term, with which we are principally concerned, and which we usually write $h \frac{du_x}{dx}$. But in the Calculus of Finite Differences, it is the whole of that series which forms the object of our investigations, and it is usually written Δu_x , so that

$$\Delta u_x = u_{x+h} - u_x.$$

2. In deducing the Difference of any proposed function u_x , the increment of the principal variable x is always supposed to be known and finite; its value however is not commonly taken

to be an undetermined quantity h , but to be unity ; both for the sake of simplicity, and because that is the value which the increment must necessarily have, in order to pass on to the succeeding term, when u_x is regarded as the general term of a Series ; and it is in that light that it is by far the most frequently regarded in Finite Differences ; so that the operation which the symbol Δ prefixed to u_x implies is, for the most part,

$$\Delta u_x = u_{x+1} - u_x.$$

There are, however, in this Subject, several important theorems which it is advantageous to investigate on the hypothesis of an indeterminate increment h for the principal variable, instead of unity ; as the process is the same on either supposition, and the result one of greater generality. And in any case if it should be desirable to introduce the same hypothesis, the expression $f(x)$ must be prepared by first writing hz instead of x ; then when z becomes $z + 1$, hz will become $h(z + 1)$ or $hz + h$, that is, the corresponding increment of x will be h ; and if the requisite analytical operations be now performed upon $f(hz)$, on the usual supposition of $\Delta z = 1$, we shall get the result expressed in terms of x , and adapted to the supposition of an indeterminate increment for x , by restoring $\frac{x}{h}$ in the place of z .

3. By a Series is meant a regular progression of terms increasing or decreasing in magnitude according to a certain law ; hence, when that law is given, and also the place of any term in the series, the magnitude of the term may be found, and thus the successive terms of the series may be produced in order. The place of any term in a series is assigned by giving the number of terms by which it is removed from some one which is considered as fixed. This number is called the *Index* of the term to which it belongs. Thus in the series

$$0, 1, 8, 27, 64, \dots x^3 \dots$$

taking the first term as the point of departure, we have the corresponding series of indices

$$0, 1, 2, 3, 4, \dots x, \dots$$

If the series be continued backwards, the indices must be considered as negative; thus the backward continuation of the above series gives the terms

$$\dots -x^3, \dots -27, -8, -1,$$

with the corresponding indices

$$-x, \dots -3, -2, -1.$$

4. Since the magnitude of every term is determined solely by its index and by the law of the series, it follows that any term is a certain function of its index, the form of which does not alter in passing from one term to another, but remains the same throughout the whole series. Thus in the above series every term is the cube of its index.

This function analytically expressed is called the general term of the series; (in the above series the general term is x^3 ;) and it is evident that all the terms of the series will be produced from it in order, by substituting successively for the index x , the progression of natural numbers

$$\dots -2, -1, 0, 1, 2, \dots$$

The general term of a series is usually denoted by u_x , where u_x is a certain function of x determined by the nature of the series. Thus, u_x denoting the general term, the series will be

$$\dots u_{-2}, u_{-1}, u_0, u_1, u_2, \dots u_{x-1}, u_x, u_{x+1}, \dots$$

Any group of consecutive terms u_x, u_{x+1}, u_{x+2} , &c. are called successive values of the function u_x .

5. The excess of any term u_{x+1} above that which immediately precedes it, or the function $u_{x+1} - u_x$ is called, as has been stated, the Difference of the function u_x , and is denoted by Δu_x . (In certain cases, which will however be expressly mentioned, we shall take Δu_x to mean $u_{x+h} - u_x$.) Hence the characteristic or symbol of operation Δ , prefixed to a given function of x , denotes the series of operations of changing x into $x+1$, and of subtracting from the altered value of the function the proposed value.

It is obvious that $u_{x+1} - u_x$ is itself in general a certain function of x , the nature of which is entirely dependent on that of the original function u_x from which it is derived, and is susceptible of a difference.

The difference, consequently, of the function Δu_x (which must be considered as having Δu for its characteristic, in the same manner as u_x has u) is

$$\Delta (\Delta u_x) = \Delta u_{x+1} - \Delta u_x,$$

which is usually written $\Delta^2 u_x$.

In like manner

$$\Delta (\Delta^2 u_x) = \Delta^3 u_x = \Delta^2 u_{x+1} - \Delta^2 u_x,$$

$$\Delta (\Delta^3 u_x) = \Delta^4 u_x = \Delta^3 u_{x+1} - \Delta^3 u_x,$$

.....

$$\Delta^n u_x = \Delta^{n-1} u_{x+1} - \Delta^{n-1} u_x.$$

6. Hence if in any function of x we change x into $x+1$, and from the result subtract the proposed function, we obtain the first difference of the proposed function; and the second, third, &c. differences are formed, each from the preceding, by a similar operation. To determine these differences of given functions, and to investigate the relations which hold between differences of any orders and the functions from which they are derived, is the object of the direct method of Finite Differences.

We shall now proceed to give instances of finding the differences of various functions, according to the above definition.

Differences of Explicit Functions.

7. To find the difference of $au_x + c$, u_x being any function of x , and a and c quantities independent of x .

$$\Delta (au_x + c) = au_{x+1} + c - (au_x + c) = a(u_{x+1} - u_x) = a\Delta u_x.$$

Hence, making $a = 0$, $\Delta c = 0$.

8. To find the difference of the sum of any number of functions of x .

$$\begin{aligned}\Delta(u_x + v_x + w_x) &= u_{x+1} + v_{x+1} + w_{x+1} - (u_x + v_x + w_x) \\ &= u_{x+1} - u_x + v_{x+1} - v_x + w_{x+1} - w_x \\ &= \Delta u_x + \Delta v_x + \Delta w_x.\end{aligned}$$

If some of the functions be preceded by negative signs, we shall find a similar result, viz.

$$\Delta(u_x - v_x - w_x) = \Delta u_x - \Delta v_x - \Delta w_x.$$

9. To find the difference of the product of two functions.

$$\begin{aligned}\Delta(u_x v_x) &= u_{x+1} v_{x+1} - u_x v_x = (u_x + \Delta u_x)(v_x + \Delta v_x) - u_x v_x \\ &= u_x \Delta v_x + v_x \Delta u_x + \Delta u_x \cdot \Delta v_x = u_x \Delta v_x + v_{x+1} \Delta u_x.\end{aligned}$$

10. To find the difference of the quotient of two functions.

$$\begin{aligned}\Delta\left(\frac{u_x}{v_x}\right) &= \frac{u_{x+1}}{v_{x+1}} - \frac{u_x}{v_x} = \frac{(u_x + \Delta u_x)v_x - (v_x + \Delta v_x)u_x}{v_{x+1}v_x} \\ &= \frac{v_x \Delta u_x - u_x \Delta v_x}{v_{x+1}v_x}.\end{aligned}$$

11. To find the difference of the continued product of any number of successive values of a function.

$$\begin{aligned}\Delta(u_x u_{x+1} \dots u_{x+n}) &= u_{x+1} u_{x+2} \dots u_{x+n+1} - u_x u_{x+1} \dots u_{x+n} \\ &= u_{x+1} u_{x+2} \dots u_{x+n} (u_{x+n+1} - u_x).\end{aligned}$$

Hence in the particular case where $u_x = a + bx$,

$$\text{since } u_{x+n+1} = a + b(x+n+1) = u_x + (n+1)b,$$

$$\Delta(u_x u_{x+1} \dots u_{x+n}) = u_{x+1} u_{x+2} \dots u_{x+n} \cdot (n+1)b.$$

12. To find the difference of a fraction whose numerator and denominator are the continued products of any number of successive values of two functions u_x and v_x respectively.

$$\Delta\left(\frac{u_x u_{x+1} \dots u_{x+n}}{v_x v_{x+1} \dots v_{x+m}}\right) = \frac{u_{x+1} u_{x+2} \dots u_{x+n}}{v_x v_{x+1} \dots v_{x+m+1}} (v_x u_{x+n+1} - u_x v_{x+m+1}).$$

Hence
$$\Delta \frac{1}{v_x v_{x+1} \dots v_{x+m}} = - \frac{v_{x+m+1} - v_x}{v_x v_{x+1} \dots v_{x+m+1}},$$

and in the particular case where $v_x = a + bx$, and therefore

$$v_{x+m+1} = a + b(x + m + 1) = v_x + (m + 1)b,$$

$$\Delta \frac{1}{v_x v_{x+1} \dots v_{x+m}} = - \frac{(m + 1)b}{v_x v_{x+1} \dots v_{x+m+1}}.$$

13. To find the differences of any rational integral function, and to shew that the n^{th} difference of a rational integral function of the n^{th} degree, is constant.

Let $u_x = Ax^n + Bx^{n-1} + \dots + Ix^2 + Kx + L$, be a rational integral function; then its first difference is

$$\begin{aligned} \Delta u_x &= A\{(x+1)^n - x^n\} + B\{(x+1)^{n-1} - x^{n-1}\} + \dots + I(2x+1) + K \\ &= nAx^{n-1} + B_1x^{n-2} + \dots + I_1x + K_1, \end{aligned}$$

which is a rational integral function, one degree lower than the original function. In like manner, for the difference of this, or the second difference of u_x , we have

$$\Delta^2 u_x = n(n-1)Ax^{n-2} + B_2x^{n-3} + \dots + I_2,$$

and so on; and for the n^{th} difference, we have

$$\Delta^n u_x = n(n-1) \dots 3 \cdot 2 \cdot 1 \cdot A.$$

Hence the n^{th} difference is constant, and the differences of all orders superior to the n^{th} vanish.

Also $\Delta^n(x^n) = 1 \cdot 2 \cdot 3 \dots n$.

14. To find the differences of a^x .

$$\Delta a^x = a^{x+1} - a^x = a^x(a-1),$$

$$\Delta^2 a^x = (a-1)\Delta a^x = a^x(a-1)^2,$$

$$\Delta^n a^x = a^x(a-1)^n.$$

Also $\Delta a^{v_x} = a^{v_x + \Delta v_x} - a^{v_x} = a^{v_x}(a^{\Delta v_x} - 1).$

15. To find the difference of $\log v_x$.

$$\Delta (\log v_x) = \log v_{x+1} - \log v_x = \log \frac{v_{x+1}}{v_x} = \log \left(1 + \frac{\Delta v_x}{v_x} \right).$$

16. To find the differences of $\sin v_x$ and $\cos v_x$.

$$\begin{aligned} \Delta \sin v_x &= \sin (v_x + \Delta v_x) - \sin v_x = 2 \sin \frac{\Delta v_x}{2} \cos \left(v_x + \frac{\Delta v_x}{2} \right) \\ &= 2 \sin \frac{\Delta v_x}{2} \sin \left\{ v_x + \frac{1}{2} (\pi + \Delta v_x) \right\}. \end{aligned}$$

$$\begin{aligned} \Delta \cos v_x &= \cos (v_x + \Delta v_x) - \cos v_x = -2 \sin \frac{\Delta v_x}{2} \sin \left(v_x + \frac{\Delta v_x}{2} \right) \\ &= 2 \sin \frac{\Delta v_x}{2} \cos \left\{ v_x + \frac{1}{2} (\pi + \Delta v_x) \right\}. \end{aligned}$$

Hence making $v_x = x\theta + \alpha$ so that $\Delta v_x = \theta$, we get

$$\Delta \sin (x\theta + \alpha) = 2 \sin \frac{1}{2} \theta \sin \left\{ x\theta + \alpha + \frac{1}{2} (\pi + \theta) \right\},$$

which shews that the difference of the sine of an angle $x\theta + \alpha$ is found by adding $\frac{1}{2}(\pi + \theta)$ to the angle, and multiplying by $2 \sin \frac{1}{2} \theta$.

Hence, repeating the operation n times, we get

$$\Delta^n \sin (x\theta + \alpha) = (2 \sin \frac{1}{2} \theta)^n \sin \left\{ x\theta + \alpha + \frac{1}{2} n (\pi + \theta) \right\}.$$

Similarly,

$$\Delta^n \cos (x\theta + \alpha) = (2 \sin \frac{1}{2} \theta)^n \cos \left\{ x\theta + \alpha + \frac{1}{2} n (\pi + \theta) \right\}.$$

Also $\Delta^n \sin \pi x = 2^n \sin \pi (x + n),$

$$\Delta^n \cos \pi x = 2^n \cos \pi (x + n).$$

16*. To find the differences of $\tan v_x$ and $\tan^{-1} v_x$.

$$\begin{aligned} \Delta \tan v_x &= \tan v_{x+1} - \tan v_x = \frac{\sin v_{x+1} \cos v_x - \cos v_{x+1} \sin v_x}{\cos v_{x+1} \cos v_x} \\ &= \frac{\sin (v_{x+1} - v_x)}{\cos v_{x+1} \cos v_x} = \frac{\sin \Delta v_x}{\cos v_{x+1} \cos v_x}. \end{aligned}$$

$$\text{Hence} \quad \Delta \tan x\theta = \frac{\sin \theta}{\cos (x+1) \theta \cos x\theta}.$$

$$\begin{aligned} \text{Also} \quad \Delta \tan^{-1} v_x &= \tan^{-1} v_{x+1} - \tan^{-1} v_x = \tan^{-1} \frac{v_{x+1} - v_x}{1 + v_{x+1} v_x} \\ &= \tan^{-1} \frac{\Delta v_x}{1 + v_{x+1} v_x}. \end{aligned}$$

$$\text{Hence} \quad \Delta \tan^{-1} x\theta = \tan^{-1} \frac{\theta}{1 + (x+1) x\theta^2}.$$

Relations between the Successive Values and the Differences of a Function.

17. The successive values of u_x any function of x are, as has been stated, the values which arise from substituting $x+1$, $x+2$, &c., or more generally $x+h$, $x+2h$, &c., for x in u_x . These values are usually written u_{x+h} , u_{x+2h} , u_{x+3h} , &c.; but it is further requisite that the operation of forming these values should be denoted by a prefixed symbol, in order that the notation for the successive values of u_x may be analogous to that for its successive differences, and in order that we may be able to avail ourselves of the advantages which, in these investigations, the Method of Separation of Symbols offers. Suppose therefore D to be a symbol of operation implying the change of x into $x+h$ in any function of x to which it is prefixed, so that $Du_x = u_{x+h}$; then

$$D(Du_x) = Du_{x+h} = u_{x+2h},$$

which may be written $D^2u_x = u_{x+2h}$; similarly $D^3u_x = u_{x+3h}$, and generally $D^n u_x = u_{x+nh}$.

Hence since $\Delta u_x = u_{x+h} - u_x$, we have $\Delta u_x = Du_x - u_x$, which, if we separate the symbol of operation D from that of quantity u_x , may be written $\Delta u_x = (D-1)u_x$, and expresses that the operation indicated by Δ is equivalent to that indicated by $D-1$. Also since $u_{x+1} = u_x + \Delta u_x$, we have $Du_x = u_x + \Delta u_x$, which, if we separate the symbols, may be written

$$Du_x = (1 + \Delta) u_x,$$

and expresses that the operation indicated by D is equivalent to that denoted by $1 + \Delta$.

18. To express $\Delta^n u_x$ by u_x and its n successive values,

$$u_{x+h}, \quad u_{x+2h}, \quad \dots \quad u_{x+nh}.$$

We here take h instead of unity for the increment of the principal variable, as the investigation is precisely the same on either supposition.

$$\Delta u_x = u_{x+h} - u_x,$$

$$\Delta^2 u_x = u_{x+2h} - u_{x+h} - (u_{x+h} - u_x) = u_{x+2h} - 2u_{x+h} + u_x,$$

$$\begin{aligned} \Delta^3 u_x &= u_{x+3h} - 2u_{x+2h} + u_{x+h} - (u_{x+2h} - 2u_{x+h} + u_x), \\ &= u_{x+3h} - 3u_{x+2h} + 3u_{x+h} - u_x. \end{aligned}$$

Now suppose this law of the coefficients, which as far as we have gone is the same as that of an expanded binomial whose index is the order of the difference, to hold for the n^{th} difference, so that

$$\Delta^n u_x =$$

$$u_{x+nh} - p_1 u_{x+(n-1)h} + p_2 u_{x+(n-2)h} - \dots \mp p_1 u_{x+h} \pm u_x,$$

$$\text{then } \Delta^{n+1} u_x =$$

$$u_{x+(n+1)h} - p_1 u_{x+nh} + p_2 u_{x+(n-1)h} - \dots \mp p_1 u_{x+2h} \pm u_{x+h}$$

$$- (u_{x+nh} - p_1 u_{x+(n-1)h} + \dots \pm p_2 u_{x+2h} \mp p_1 u_{x+h} \pm u_x)$$

$$= u_{x+(n+1)h} - (1 + p_1) u_{x+nh} + (p_1 + p_2) u_{x+(n-1)h} - \dots$$

$$\pm (1 + p_1) u_{x+h} \mp u_x,$$

which is the same alteration with regard to the coefficients as occurs in passing from $(z-1)^n$ to $(z-1)^{n+1}$. If therefore, for any value of n supposed a positive integer, the coefficients of the expansions of $\Delta^n u_x$ and $(z-1)^n$ are the same, they will always be the same; but these coefficients are identical, as we have seen, when $n = 1, 2, 3$; therefore they are always the same;

$$\therefore \Delta^n u_x = u_{x+n} - nu_{x+(n-1)h} + \frac{n(n-1)}{1 \cdot 2} u_{x+(n-2)h} - \dots$$

$$\mp nu_{x+h} \pm u_x;$$

or, supposing $h = 1$,

$$\Delta^n u_x = u_{x+n} - nu_{x+n-1} + \frac{n(n-1)}{1 \cdot 2} u_{x+n-2} - \dots \mp nu_{x+1} \pm u_x.$$

19. If for the descending values u_{x+n} , u_{x+n-1} , &c. we substitute their equivalents (Art. 17), the second member becomes

$$D^n u_x - nD^{n-1} u_x + \frac{n(n-1)}{1 \cdot 2} D^{n-2} u_x - \&c. \dots \pm u_x,$$

or, separating the symbols of operation from those of quantity,

$$(D^n - nD^{n-1} + \frac{n(n-1)}{1 \cdot 2} D^{n-2} - \&c. \dots \pm 1) u_x,$$

$$\therefore \Delta^n u_x = (D-1)^n u_x,$$

each term of the development of $(D-1)^n$ being understood to be prefixed to u_x . This formula results immediately from the definitions of Δ and D (Art. 17); for as the operation denoted by Δ is equivalent to that denoted by $D-1$, if these operations be performed n times upon the same function u_x , we must have

$$\Delta^n u_x = (D-1)^n u_x.$$

20. If in the series just investigated, we assign a particular value to u_x , we shall readily obtain an expression for its n^{th} difference. Thus let $u_x = x^m$,

$$\therefore \Delta^n (x^m) = (x+n)^m - n(x+n-1)^m + \frac{n(n-1)}{1 \cdot 2} (x+n-2)^m - \&c.,$$

in which equation, if $n > m$, since the former member vanishes (Art. 13), the second member is zero for every value of x ; and if $m = n$, so that $\Delta^n x^n = 1 \cdot 2 \cdot 3 \dots n$, we get

$$1 \cdot 2 \cdot 3 \dots n = (x+n)^n - n(x+n-1)^n + \frac{n(n-1)}{1 \cdot 2} (x+n-2)^n - \&c.;$$

and making $x = 0$, since the equation holds for all values of x ,

$$1 \cdot 2 \cdot 3 \dots n = n^n - n(n-1)^n + \frac{n(n-1)}{1 \cdot 2} (n-2)^n - \&c.$$

If $x = 0$, and $\Delta^n 0^m$ denote the particular value of $\Delta^n x^m$ when $x = 0$, we have

$$\Delta^n 0^m = n^m - n(n-1)^m + \frac{n(n-1)}{1 \cdot 2} (n-2)^m - \&c.$$

Of the numbers comprised in the form $\Delta^n 0^m$, which are called the Differences of zero, we shall make considerable use in future investigations; whenever $n > m$ the value is zero, in other cases it may be computed by the above formula; thus,

$$\Delta^0 0^m = 1,$$

$$\Delta^2 0^2 = 2, \quad \Delta^2 0^3 = 6, \quad \Delta^2 0^4 = 14, \dots$$

$$\Delta^3 0^3 = 6, \quad \Delta^3 0^4 = 36, \quad \Delta^3 0^5 = 150, \dots$$

21. Reversing the order of the series in Art. 18, we find

$$(-1)^n \Delta^n u_x = u_x - n u_{x+1} + \frac{n(n-1)}{1 \cdot 2} u_{x+2} - \dots \pm u_{x+n}.$$

Also putting $u_x = x^m$, and then supposing $m = n$, $x = 1$, we find successively,

$$(-1)^n \Delta^n x^m = x^m - n(x+1)^m + \frac{n(n-1)}{1 \cdot 2} (x+2)^m - \dots \pm (x+n)^m,$$

$$(-1)^n 1 \cdot 2 \cdot 3 \dots n = x^n - n(x+1)^n + \frac{n(n-1)}{1 \cdot 2} (x+2)^n - \dots \pm (x+n)^n,$$

$$(-1)^n 1 \cdot 2 \cdot 3 \dots n = 1^n - n \cdot 2^n + \frac{n(n-1)}{1 \cdot 2} \cdot 3^n - \dots \pm (n+1)^n.$$

22. Hence it may be proved that $1 \cdot 2 \cdot 3 \dots (p-1) + 1$ is divisible by p , if p be a prime number. (Wilson's Theorem)

Let $u_x = x^n - 1$;

$$\begin{aligned} \therefore \Delta^n (x^n - 1) &= 1 \cdot 2 \cdot 3 \dots n = (x+n)^n - 1 - n \{(x+n-1)^n - 1\} \\ &\quad + \frac{n(n-1)}{1 \cdot 2} \{(x+n-2)^n - 1\} - \&c. \end{aligned}$$

Let $x = 1$, and $n + 1 = p$, then $x + n = p$, and

$$1 \cdot 2 \cdot 3 \dots (p-1) + 1 = p^n - (p-1) \{ (p-1)^{p-1} - 1 \} \\ + \frac{(p-1)(p-2)}{1 \cdot 2} \{ (p-2)^{p-1} - 1 \} - \&c.$$

Now by Fermat's Theorem every term of the second member is divisible by p when p is a prime number; consequently

$$1 \cdot 2 \cdot 3 \dots (p-1) + 1 \text{ is divisible by } p.$$

23. To express u_{x+nh} by u_x and its first n differences.

We take h instead of unity for the increment of the principal variable, the investigation being precisely the same on either supposition.

$$u_{x+h} = u_x + \Delta u_x,$$

$$u_{x+2h} = u_x + \Delta u_x + \Delta(u_x + \Delta u_x) = u_x + 2\Delta u_x + \Delta^2 u_x,$$

$$u_{x+3h} = u_x + 2\Delta u_x + \Delta^2 u_x + \Delta(u_x + 2\Delta u_x + \Delta^2 u_x) \\ = u_x + 3\Delta u_x + 3\Delta^2 u_x + \Delta^3 u_x.$$

Now suppose this law of coefficients, which as far as we have gone is the same as that of an expanded binomial whose index is the number of increments which the principal variable has received, to hold for n increments, so that

$$u_{x+nh} = u_x + p_1 \Delta u_x + p_2 \Delta^2 u_x + \dots + p_{n-1} \Delta^{n-1} u_x + \Delta^n u_x,$$

$$\text{then } u_{x+(n+1)h} = u_x + p_1 \Delta u_x + p_2 \Delta^2 u_x + \dots + p_{n-1} \Delta^{n-1} u_x + \Delta^n u_x \\ + \Delta(u_x + p_1 \Delta u_x + \dots + p_{n-1} \Delta^{n-1} u_x + \Delta^n u_x) \\ = u_x + (1 + p_1) \Delta u_x + (p_1 + p_2) \Delta^2 u_x + \dots \\ + (p_2 + p_1) \Delta^{n-1} u_x + (p_1 + 1) \Delta^n u_x + \Delta^{n+1} u_x,$$

which is the same alteration with regard to the coefficients as occurs in passing from $(1+z)^n$ to $(1+z)^{n+1}$.

Hence if the coefficients of the expansions of u_{x+nh} and $(1+z)^n$ are the same for any value of n supposed a positive integer, they

will always be the same; but they are identical as we have seen when $n = 1, 2, 3$; therefore they are always the same;

$$\therefore u_{x+n\Delta} = u_x + n\Delta u_x + \frac{n(n-1)}{1 \cdot 2} \Delta^2 u_x + \dots + n\Delta^{n-1} u_x + \Delta^n u_x,$$

or, supposing $h = 1$,

$$u_{x+n} = u_x + n\Delta u_x + \frac{n(n-1)}{1 \cdot 2} \Delta^2 u_x + \dots + n\Delta^{n-1} u_x + \Delta^n u_x.$$

24. This result, if we separate the symbols, may be written

$$u_{x+n} = (1 + n\Delta + \frac{n(n-1)}{1 \cdot 2} \Delta^2 + \dots + n\Delta^{n-1} + \Delta^n) u_x,$$

$$\text{or } u_{x+n} = (1 + \Delta)^n u_x,$$

each term of the development of $(1 + \Delta)^n$ being understood to be prefixed to u_x . The same formula follows immediately from the definitions of the symbols D and Δ ; for as D is equivalent to $1 + \Delta$, if the operations denoted by D and $1 + \Delta$ be performed n times upon the same function u_x , we must have

$$D^n u_x = (1 + \Delta)^n u_x; \text{ but } D^n u_x = u_{x+n},$$

$$\therefore u_{x+n} = (1 + \Delta)^n u_x.$$

24*. Let $u_x = x^m$, then $u_{x+n} = (x+n)^m$;

$$\therefore (x+n)^m = (1 + \Delta)^n x^m,$$

$$\text{and making } x = 0, \quad n^m = (1 + \Delta)^n 0^m,$$

each term of the development of $(1 + \Delta)^n$ being prefixed, as said above, to x^m and 0^m , respectively. By the latter formula any power of a number is expressed by the numbers comprised in the form $\Delta^n 0^m$, i. e. by the differences of zero.

25. To deduce Taylor's Theorem from the formula

$$u_{x+n\Delta} = u_x + n\Delta u_x + \frac{n(n-1)}{1 \cdot 2} \Delta^2 u_x + \&c.$$

Let $n\Delta = t$, and let h be infinitely diminished whilst t remains finite; therefore n is infinitely increased; and since

h is indefinitely diminished, we have, regarding the differential coefficient as the limit of the ratio of the simultaneous increments of the function and the variable,

$$\frac{\Delta u_x}{h} = \frac{du_x}{dx},$$

$$\frac{\Delta^2 u_x}{h^2} = \frac{d}{dx} \left(\frac{\Delta u_x}{h} \right) = \frac{d^2 u_x}{dx^2}; \text{ \&c.}$$

Hence, preparing the formula as follows,

$$u_{x+nh} = u_x + nh \frac{\Delta u_x}{h} + \frac{nh(nh-h)}{1 \cdot 2} \frac{\Delta^2 u_x}{h^2} + \text{\&c.},$$

and taking the limit of both sides by supposing h to be infinitely diminished and n infinitely increased, their product always remaining equal to a finite magnitude t , we get

$$u_{x+t} = u_x + t \frac{du_x}{dx} + \frac{t^2}{1 \cdot 2} \frac{d^2 u_x}{dx^2} + \text{\&c.}$$

The Differential Calculus is a particular case of that of Finite Differences; and the above investigation is introduced to show how, from results in Finite Differences obtained with an indeterminate increment for the principal variable, we may pass to the corresponding results in the Differential Calculus.

The theorems of Arts. 18 and 23 have been proved by an inductive process; they may also be established by the theory of Generating Functions, the principles of which we shall now proceed to explain; as it is a theory which, for its generality and power, especially merits our attention.

Generating Functions.

26. Let $\phi(t)$ be a function of t , susceptible of the development

$$\begin{aligned} \phi(t) = & \dots + u_{-1}t^{-1} + u_0 + u_1t + \dots \\ & + u_{x-1}t^{x-1} + u_x t^x + u_{x+1}t^{x+1} + \dots; \end{aligned}$$

then u_x may evidently represent any function of x whatever, if we regard this equation as the definition of $\phi(t)$. The

function $\phi(t)$ consequently by its development generates the coefficients $u_0, u_1, \dots u_x$ annexed to their proper powers of t , and is therefore called the Generating Function of u_x , and is denoted by Gu_x , so that

$$\phi(t) = Gu_x.$$

Thus since

$$\log(1-t)^{-1} = t + \frac{1}{2}t^2 + \frac{1}{3}t^3 + \dots + \frac{1}{x}t^x + \dots$$

$$\log(1-t)^{-1} = G \frac{1}{x}.$$

Similarly, since

$$t(1-t)^{-2} = t + 2t^2 + 3t^3 + \dots + xt^x + \dots$$

$$t(1-t)^{-2} = Gx.$$

27. To determine the generating functions of u_{x+n} and $\Delta^n u_x$ from that of u_x .

Let $\phi(t) = Gu_x$,

then $\phi(t) = \dots + u_n t^n + u_{n+1} t^{n+1} + \dots + u_{x+n} t^{x+n} + \dots (1);$

$$\therefore t^n \phi(t) = \dots + u_n + u_{n+1} t + \dots + u_{x+n} t^x + \dots$$

$$\therefore t^n \phi(t) = Gu_{x+n}, \text{ or } t^n Gu_x = Gu_{x+n}.$$

Again, $t^n \phi(t) = \dots + u_{-n} + u_{-n+1} t + \dots + u_{x-n} t^x + \dots$

$$\therefore t^n \phi(t) = Gu_{x-n}, \text{ or } t^n Gu_x = Gu_{x-n}.$$

Hence it follows that the generating function of

$$\Delta u_x, \text{ or } u_{x+1} - u_x, \text{ is } \left(\frac{1}{t} - 1\right) \phi(t),$$

for this function being developed will produce the difference of two series whose general terms are respectively $u_{x+1} t^x$ and $u_x t^x$;

$$\therefore G(\Delta u_x) = \left(\frac{1}{t} - 1\right) Gu_x.$$

Similarly,

$$G(\Delta^2 u_x) = \left(\frac{1}{t} - 1\right) G(\Delta u_x) = \left(\frac{1}{t} - 1\right)^2 Gu_x,$$

$$\text{and } G(\Delta^n u_x) = \left(\frac{1}{t} - 1\right)^n Gu_x.$$

Also

$$G(\Delta^n u_{x-n}) = \left(\frac{1}{t} - 1\right)^n G u_{x-n} = \left(\frac{1}{t} - 1\right)^n t^n G u_x = (1-t)^n G u_x.$$

28. If instead of multiplying $\phi(t)$ by a power of t , we multiply it by another function of t , $\psi(t)$ similar to $\phi(t)$, that is, capable of being developed in a series of integral powers of t positive or negative, so that

$$\psi(t) = \dots + p_{-1}t^{-1} + p_0 + p_1t + p_2t^2 + \dots + p_xt^x + \dots (2),$$

then in the product we find for the coefficient of t^x ,

$$\dots + p_{-1}u_{x+1} + p_0u_x + p_1u_{x-1} + p_2u_{x-2} + \dots$$

which may be replaced (Art. 17) by

$$\dots + p_{-1}Du_x + p_0u_x + p_1D^{-1}u_x + p_2D^{-2}u_x + \dots;$$

and this, separating the symbols, may be written

$$(\dots + p_{-1}D + p_0 + p_1D^{-1} + p_2D^{-2} + \dots) u_x,$$

or $\psi\left(\frac{1}{D}\right)u_x$, (each term of the development of $\psi\left(\frac{1}{D}\right)$ being understood to be prefixed to u_x); which shews that $\psi(t) \times \phi(t)$ is the generating function of $\psi\left(\frac{1}{D}\right)u_x$; in other words,

$$\psi(t) \times G u_x = G \left\{ \psi\left(\frac{1}{D}\right) u_x \right\}.$$

And if in (2) we replace t by t^{-1} it may be shewn in exactly the same way, that $\psi\left(\frac{1}{t}\right) \times \phi(t)$ is the generating function of $\psi(D)u_x$, that is

$$\psi\left(\frac{1}{t}\right) \times G u_x = G \left\{ \psi(D) u_x \right\}.$$

Suppose for example, $\psi(t)$ to assume successively the forms t^n , $(t^{-1} - 1)^n$; then, as before, we get

$$t^n \times G u_x = G (D^n u_x) = G (u_{x+n}),$$

$$(t^{-1} - 1)^n \times G u_x = G \{(D - 1)^n u_x\} = G (\Delta^n u_x).$$

29. To investigate the expression for $\Delta^n u_x$ in terms of u_{x+n} , u_{x+n-1} , &c., by Generating Functions.

$$\begin{aligned}
 G(\Delta^n u_x) &= \left(\frac{1}{t} - 1\right)^n G u_x \\
 &= t^n G u_x - n t^{n+1} G u_x + \frac{n(n-1)}{1 \cdot 2} t^{n+2} G u_x - \&c. \\
 &= G u_{x+n} - n G u_{x+n-1} + \frac{n(n-1)}{1 \cdot 2} G u_{x+n-2} - \&c. \\
 &= G(u_{x+n} - n u_{x+n-1} + \frac{n(n-1)}{1 \cdot 2} u_{x+n-2} - \&c.) \\
 \therefore \Delta^n u_x &= u_{x+n} - n u_{x+n-1} + \frac{n(n-1)}{1 \cdot 2} u_{x+n-2} - \&c.
 \end{aligned}$$

for, the generating functions of both being the same, the coefficients of t^x in the developments of those functions must be identical, however those developments have been effected.

30. To investigate the expression for u_{x+n} in terms of u_x and its first n differences, by Generating Functions.

$$\begin{aligned}
 G u_{x+n} &= t^n G u_x \\
 &= \left\{1 + \left(\frac{1}{t} - 1\right)\right\}^n G u_x \\
 &= G u_x + n \left(\frac{1}{t} - 1\right) G u_x + \frac{n(n-1)}{1 \cdot 2} \left(\frac{1}{t} - 1\right)^2 G u_x + \&c. \\
 &= G u_x + n G(\Delta u_x) + \frac{n(n-1)}{1 \cdot 2} G(\Delta^2 u_x) + \&c. \\
 &= G(u_x + n \Delta u_x + \frac{n(n-1)}{1 \cdot 2} \Delta^2 u_x + \&c.); \\
 \therefore u_{x+n} &= u_x + n \Delta u_x + \frac{n(n-1)}{1 \cdot 2} \Delta^2 u_x + \&c.
 \end{aligned}$$

31. It is obvious that by transforming the expressions

$$\left(\frac{1}{t} - 1\right)^n G u_x, \text{ and } t^n G u_x,$$

in different ways, we may obtain various other expressions for $\Delta^n u_x$ and u_{x+n} besides the above.

Thus to express $\Delta^n u_x$ in terms of $\Delta^n u_{x-n}$, $\Delta^{n+1} u_{x-n-1}$, &c., we have

$$\begin{aligned} G(\Delta^n u_x) &= \left(\frac{1}{t} - 1\right)^n G u_x = \frac{(1-t)^n G u_x}{\{1 - (1-t)\}^n} \\ &= (1-t)^n G u_x + n(1-t)^{n+1} G u_x \\ &\quad + \frac{n(n+1)}{1 \cdot 2} (1-t)^{n+2} G u_x + \&c. \\ &= G(\Delta^n u_{x-n}) + n G(\Delta^{n+1} u_{x-n-1}) + \frac{n(n+1)}{1 \cdot 2} G(\Delta^{n+2} u_{x-n-2}) + \&c. \\ &= G\left\{\Delta^n u_{x-n} + n\Delta^{n+1} u_{x-n-1} + \frac{n(n+1)}{1 \cdot 2} \Delta^{n+2} u_{x-n-2} + \&c.\right\}; \\ \therefore \Delta^n u_x &= \Delta^n u_{x-n} + n\Delta^{n+1} u_{x-n-1} + \frac{n(n+1)}{1 \cdot 2} \Delta^{n+2} u_{x-n-2} + \&c. \end{aligned}$$

32. Again, to express u_{x+n} in terms of u_x , Δu_{x-r} , $\Delta^2 u_{x-2r}$, &c., we must transform t^n into a series of powers of $t^r \left(\frac{1}{t} - 1\right)$, that is, we must develop t^n in powers of x from the equation

$$\frac{1}{t} = 1 + x \left(\frac{1}{t}\right)^r,$$

which may be done by Lagrange's Theorem; and we find (see Herschel's Examples)

$$\begin{aligned} u_{x+n} &= u_x + n\Delta u_{x-r} + \frac{n(n+2r-1)}{1 \cdot 2} \Delta^2 u_{x-2r} \\ &\quad + \frac{n(n+3r-1)(n+3r-2)}{1 \cdot 2 \cdot 3} \Delta^3 u_{x-3r} + \&c. \end{aligned}$$

This method is obviously not confined to the function $u_{x+1} - u_x$; it is equally applicable to any other combination of the successive values u_x , u_{x+1} , u_{x+2} , &c., of the first degree. If we

take Δu_x to mean $au_{x+2} + bu_{x+1} + cu_x$, then the generating functions of Δu_x and $\Delta^n u_x$ will evidently be

$$\left(\frac{a}{\ell^2} + \frac{b}{\ell} + c\right) Gu_x, \quad \left(\frac{a}{\ell^2} + \frac{b}{\ell} + c\right)^n Gu_x,$$

and the expression for $\Delta^n u_x$ in terms of u_x and its successive values, might be obtained as in the preceding case.

Separation of the Symbols of Operation from those of Quantity.

33. We have seen (Arts. 18 and 23) in the formulæ

$$D^n u_x = (1 + \Delta)^n u_x, \quad \Delta^n u_x = (D - 1)^n u_x,$$

instances of the method which consists in separating the symbols of operation from those of quantity; the use of which is not confined to simple cases like those just noticed, but may be extended with remarkable effect to a great variety of investigations connected with this subject.

By symbols of operation are meant certain characteristic letters placed before any functions, to denote that certain operations have been performed on them: thus D placed before a function of any variable x , denotes the operation of changing in it x into $x + h$; and Δ placed before the same function, implies that two different values of the variable have been substituted in it and the results subtracted from one another. By symbols of quantity are meant the subjects of the operations just mentioned; that is, letters taken to represent numbers, or algebraical expressions; and it must be remarked that these latter may be also regarded as symbols of operation. For if a be a number, then a denotes that the unit employed in the investigation, whatever it be, is to be added to itself n times; and a^2 or $a \cdot a$ denotes that the same operation has been performed on a that was performed on unity; whenever therefore the two kinds of symbols occur together in the same formula, as in $(Da - 1)^n$, a must be regarded as a symbol of operation.

34.* The expressions $(1 + \Delta)^n$, $(D - 1)^n$ must be taken as abbreviated forms for their developments; and when prefixed to the function u_x , each term of these developments is understood to

be applied separately to that function. And, in general, if $F(\Delta)$ be a function of Δ capable of being developed in a series of powers of Δ such as

$$A\Delta^{\alpha} + B\Delta^{\beta} + \&c.; \text{ then for } A\Delta^{\alpha}u_x + B\Delta^{\beta}u_x + \&c.$$

the expression $F(\Delta) u_x$ is used as an abbreviation; and the same notation is applicable to other characteristic symbols, such as

$$\frac{d}{dx}, \int dx, \Sigma, \delta_x,$$

that in their combinations are subject to the same laws as algebraical quantities.

Also if we replace $\Delta^{\alpha}u_x$, $\Delta^{\beta}u_x$, &c. by their equivalents, we get

$$\begin{aligned} F(\Delta) u_x &= A(D-1)^{\alpha}u_x + B(D-1)^{\beta}u_x + \&c. \\ &= \{A(D-1)^{\alpha} + B(D-1)^{\beta} + \&c.\} u_x = F(D-1) u_x, \end{aligned}$$

which shews that in any formula the symbols Δ and $D-1$ may be interchanged.

Hence also the successive performance of two or more series of operations represented by $F(\Delta)$, $F'(\Delta)$, upon the same function u_x , is equivalent to the performance of that series of operations denoted by their product. The method of separation of symbols is in every case capable of a strict inductive proof, and does not rest merely upon accidental analogies; and it deserves great notice on account of the facility with which it enables us to conduct many intricate processes. But as the generalizations which it offers, may present some difficulties to the student, we shall continue, as we have done hitherto, to obtain several of the principal results by an elementary process; before investigating them by the method of separation of symbols, or pointing out how they arise from that method.

We shall now give instances of the application of the method of separation of symbols, to obtain several important results.

34. To find the n^{th} difference of the product of two functions.

We have seen that $\Delta(u_x v_x) = \Delta u_x v_{x+1} + u_x \Delta v_x$

$$= \Delta u_x D v_x + u_x \Delta v_x = (\Delta D' + \Delta') u_x v_x,$$

separating the symbols, and supposing in the second member Δ to affect u_x only, and Δ' and D' to affect v_x only,

then $\Delta^2(u_x v_x) = (\Delta D' + \Delta') \Delta (u_x v_x) = (\Delta D' + \Delta')^2 u_x v_x$;
and, generally, $\Delta^n(u_x v_x) = (\Delta D' + \Delta')^n u_x v_x$

$$= (\Delta^n D'^n + n \Delta^{n-1} \Delta' D'^{n-1} + \frac{n(n-1)}{1 \cdot 2} \Delta^{n-2} \Delta'^2 D'^{n-2} + \&c.) u_x v_x$$

$$= \Delta^n u_x \cdot v_{x+n} + n \Delta^{n-1} u_x \cdot \Delta v_{x+n-1} + \frac{n(n-1)}{1 \cdot 2} \Delta^{n-2} u_x \Delta^2 v_{x+n-2} + \&c.,$$

which may be also proved inductively by shewing, as in Art. 13, that the coefficients of the developments of $\Delta^n(u_x v_x)$ and $(1+z)^n$, which are identical when $n=1$, undergo the same changes in passing from n to $n+1$.

If the series be reversed or, which is the same thing, if we develop the formula $\Delta^n(u_x v_x) = (\Delta' + \Delta D')^n u_x v_x$, we get

$$\Delta^n(u_x v_x) = u_x \Delta^n v_x + n \Delta u_x \Delta^{n-1} v_{x+1} + \frac{n(n-1)}{1 \cdot 2} \Delta^2 u_x \Delta^{n-2} v_{x+2} + \&c.$$

Since $\Delta(u_x v_x) = u_{x+1} v_{x+1} - u_x v_x = D u_x D v_x - u_x v_x = (DD' - 1) u_x v_x$ where D affects u_x only, and D' affects v_x only, we obtain another development of $\Delta^n(u_x v_x)$, viz.

$$\Delta^n(u_x v_x) = (DD' - 1)^n u_x v_x$$

$$= (D^n D'^n - n D^{n-1} D'^{n-1} + \frac{n(n-1)}{1 \cdot 2} D^{n-2} D'^{n-2} - \&c.) u_x v_x$$

$$= u_{x+n} v_{x+n} - n u_{x+n-1} v_{x+n-1} + \frac{n(n-1)}{1 \cdot 2} u_{x+n-2} v_{x+n-2} - \&c.$$

In the formula $\Delta^n(u_x v_x) = (DD' - 1)^n u_x v_x$ suppose $v_x = a^x$, then $D'a^x = a^{x+1}$, so that the operation upon a^x denoted by D' is multiplying it by a ,

$$\therefore \Delta^n(u_x a^x) = (Da - 1)^n u_x a^x = a^x (Da - 1)^n u_x.$$

Also $u_{x+n} \Delta^n v_x$
 $= (\Delta' D)^n v_x u_x = (\Delta' D + \Delta - \Delta)^n v_x u_x$
 $= \{(\Delta' D + \Delta)^n - n(\Delta' D + \Delta)^{n-1} \Delta + \&c.\} v_x u_x$
 $= \Delta^n(v_x u_x) - n \Delta^{n-1}(v_x \Delta u_x) + \frac{1}{2} n(n-1) \Delta^{n-2}(v_x \Delta^2 u_x) - \dots \pm v_x \Delta^n u_x,$
 a formula by which $u_{x+n} \Delta^n v_x$ is expressed by a series of differences with constant coefficients.

In the above instance we use an accent not to imply that the operations denoted by Δ , D , are altered at all, but merely that Δ' , D' , affect v_x only, whilst Δ , D affect u_x only. The above result is sometimes written,

$$\Delta^n (u_x v_x) = (DD' - 1)^n u_x v_x = \{(1 + \Delta) (1 + \Delta') - 1\}^n u_x v_x,$$

replacing D , D' by their equivalents; or, if there are more functions w_x , z_x , &c. and we use Δ'' , Δ''' , &c. to imply that in the second member these symbols only affect w_x , z_x , &c. respectively; and similarly for D , D' , D'' , &c.; we have

$$\begin{aligned} \Delta^n (u_x v_x w_x z_x \dots) &= (DD'D'' \dots - 1)^n u_x v_x w_x \dots \\ &= \{(1 + \Delta) (1 + \Delta') (1 + \Delta'') (1 + \Delta''') \dots - 1\}^n u_x v_x w_x z_x \dots \end{aligned}$$

35. To shew that $\Delta^n u_x = (e^{\frac{d}{dx}} - 1)^n u_x$, in which the symbols of operation are separated from those of quantity.

By Taylor's Theorem, we have

$$u_{x+n} = u_x + n \frac{du_x}{dx} + \frac{n^2}{1 \cdot 2} \frac{d^2 u_x}{dx^2} + \&c.,$$

and separating the symbols of operation from those of quantity, we get

$$u_{x+n} = \left\{ 1 + n \frac{d}{dx} + \frac{n^2}{1 \cdot 2} \left(\frac{d}{dx} \right)^2 + \frac{n^3}{1 \cdot 2 \cdot 3} \left(\frac{d}{dx} \right)^3 + \&c. \right\} u_x = e^{n \frac{d}{dx}} u_x.$$

Similarly, $u_{x+n-1} = e^{(n-1) \frac{d}{dx}} u_x$, &c. $u_{x+1} = e^{\frac{d}{dx}} u_x$;

$$\therefore \Delta^n u_x = e^{n \frac{d}{dx}} u_x - n e^{(n-1) \frac{d}{dx}} u_x + \frac{n(n-1)}{1 \cdot 2} e^{(n-2) \frac{d}{dx}} u_x - \&c.;$$

or, again separating the symbols of operation from those of quantity,

$$\Delta^n u_x = \left\{ e^{n \frac{d}{dx}} - n e^{(n-1) \frac{d}{dx}} + \frac{n(n-1)}{1 \cdot 2} e^{(n-2) \frac{d}{dx}} - \&c. \right\} u_x = \left(e^{\frac{d}{dx}} - 1 \right)^n u_x,$$

a celebrated theorem first given by Lagrange.

36. To find a general expression for the n^{th} difference of a function in terms of its differential coefficients.

The development of the second member of the equation

$$\Delta^n u_x = \left(e^{\frac{d}{dx}} - 1 \right)^n u_x$$

will consist of a series of terms of the form

$$\{A_0 + A_1 \frac{d}{dx} + \dots + A_m \left(\frac{d}{dx} \right)^m + \dots\} u_x,$$

$$\text{or } A_0 u_x + A_1 \frac{du_x}{dx} + \dots + A_m \frac{d^m u_x}{dx^m} + \dots,$$

and A_m is evidently the coefficient of t^m in the expansion of $(e^t - 1)^n$.

$$\text{Now } (e^t - 1)^n = e^{nt} - ne^{(n-1)t} + \frac{n(n-1)}{1 \cdot 2} e^{(n-2)t} - \&c.,$$

and the coefficients of t^m in the developments of e^{nt} , $e^{(n-1)t}$, &c. are respectively

$$\frac{n^m}{\underline{m}}, \quad \frac{(n-1)^m}{\underline{m}}, \quad \&c.;$$

$$\therefore A_m = \frac{1}{\underline{m}} \{n^m - n(n-1)^m + \frac{n(n-1)}{1 \cdot 2} (n-2)^m - \&c.\} = \frac{\Delta^n 0^m}{\underline{m}},$$

from Art. 20.

Now so long as $m < n$, this vanishes; and when $n = m$, $\Delta^n 0^m = \underline{n}$;

$$\therefore \Delta^n u_x = \frac{d^n u_x}{dx^n} + \frac{\Delta^n 0^{n+1}}{\underline{n+1}} \frac{d^{n+1} u_x}{dx^{n+1}} + \frac{\Delta^n 0^{n+2}}{\underline{n+2}} \frac{d^{n+2} u_x}{dx^{n+2}} + \&c.$$

Ex. Let $u_x = x^m$, then

$$\begin{aligned} \Delta^n (x^m) &= m(m-1) \dots (m-n+1) x^{m-n} \\ &+ \frac{\Delta^n 0^{n+1}}{\underline{n+1}} \cdot m(m-1) \dots (m-n) x^{m-n-1} + \&c. + \Delta^n 0^n. \end{aligned}$$

37. If in Lagrange's Theorem for $\Delta^n u_x$, $n = 1$, we have $\Delta u_x = (e^{\frac{d}{dx}} - 1) u_x$, or $\Delta = e^{\frac{d}{dx}} - 1$, the meaning of which is, that the operation denoted by Δ is equivalent to the series of opera-

tions denoted by $e^{\frac{d}{dx}} - 1$. And generally, the series of operations denoted by $f(\Delta)$ is equivalent to that denoted by $f(e^{\frac{d}{dx}} - 1)$. For let $f(\Delta)$ be developed in a series of the form

$$f(\Delta) = A\Delta^a + B\Delta^b + C\Delta^c + \&c.,$$

then $f(\Delta) u_x = A\Delta^a u_x + B\Delta^b u_x + C\Delta^c u_x + \&c.$

$$= A(e^{\frac{d}{dx}} - 1)^a u_x + B(e^{\frac{d}{dx}} - 1)^b u_x + C(e^{\frac{d}{dx}} - 1)^c u_x + \&c.$$

or, separating the symbols of operation from those of quantity,

$$\begin{aligned} f(\Delta) u_x &= \{A(e^{\frac{d}{dx}} - 1)^a + B(e^{\frac{d}{dx}} - 1)^b + C(e^{\frac{d}{dx}} - 1)^c + \&c.\} u_x \\ &= f(e^{\frac{d}{dx}} - 1) u_x. \end{aligned}$$

Thus, suppose $f(\Delta) = (1 + \Delta)^n$,

$$\text{then } f(e^{\frac{d}{dx}} - 1) = (1 + e^{\frac{d}{dx}} - 1)^n = e^{n\frac{d}{dx}};$$

therefore, annexing a function u_x for the symbols to operate upon,

$$(1 + \Delta)^n u_x = e^{n\frac{d}{dx}} u_x = u_{x+n},$$

as already proved.

Similarly, making $n = 1$ in the formula $u_{x+n} = e^{n\frac{d}{dx}} u_x$ (Art. 35), we find $u_{x+1} = e^{\frac{d}{dx}} u_x$, or $Du_x = e^{\frac{d}{dx}} u_x$; therefore $D = e^{\frac{d}{dx}}$, the meaning of which is that the operation denoted by D is equivalent to the series of operations denoted by $e^{\frac{d}{dx}}$; and, generally, the series of operations denoted by $f(D)$ is equivalent to that denoted by $f(e^{\frac{d}{dx}})$, which is expressed by the formula

$$f(D) u_x = f(e^{\frac{d}{dx}}) u_x.$$

38. To express the n^{th} differential coefficient of any function by its differences.

Suppose $f(D) = (\log D)^n$, then $f(e^{\frac{d}{dx}}) = (\log e^{\frac{d}{dx}})^n = \left(\frac{d}{dx}\right)^n$;

$$\therefore \frac{d^n u_x}{dx^n} = (\log D)^n u_x = \{\log(1 + \Delta)\}^n u_x.$$

39. To find the general term of the expansion of $f(e^t)$ in a series ascending by powers of t .

Writing $f(e^t)$ in the form $f\{1 + (e^t - 1)\}$, and expanding by Taylor's theorem, we find

$$f(e^t) = f(1) + f'(1)(e^t - 1) + \frac{1}{1 \cdot 2} f''(1)(e^t - 1)^2 + \dots \\ + \frac{1}{n!} f^{(n)}(1)(e^t - 1)^n + \dots$$

Then taking, as in Art. 36, the coefficient of t^m in each term of the second member, and observing that in $f(1)$ it may be represented by $\frac{f(1) \cdot 0^m}{[m]}$, this quantity being $f(1)$ when $m = 0$, and zero in all other cases; and that in

$$\frac{1}{n!} f^{(n)}(1)(e^t - 1)^n \text{ it is } \frac{1}{n!} f^{(n)}(1) \frac{\Delta^n 0^m}{[m]},$$

we have for the coefficient of t^m in the expansion of $f(e^t)$, the value

$$A_m = \frac{1}{[m]} \{f(1) 0^m + f'(1) \Delta 0^m + \frac{1}{1 \cdot 2} f''(1) \Delta^2 0^m + \dots\} \\ = \frac{1}{[m]} f(1 + \Delta) 0^m,$$

a remarkable theorem first given by Herschel; for the applications of which, see his Collection of Examples. Hence in the development of e^{e^t} the coefficient of t^m is

$$\frac{1}{[m]} e^{1+\Delta} 0^m = \frac{e}{[m]} \left(0^m + \Delta 0^m + \frac{1}{1 \cdot 2} \Delta^2 0^m + \dots + \frac{\Delta^m 0^m}{[m]} \right);$$

and in the series for $\frac{1}{e^t + 1}$ the coefficient of t^m is

$$\frac{1}{[m]} \frac{1}{2 + \Delta} 0^m = \frac{1}{[m]} \left(\frac{0^m}{2} - \frac{\Delta 0^m}{2^2} + \frac{\Delta^2 0^m}{2^3} - \dots \pm \frac{\Delta^m 0^m}{2^{m+1}} \right);$$

the advantage of using the differences of zero being that any series of them necessarily terminates at $\Delta^m 0^m$.

SECTION II.

INVERSE METHOD OF DIFFERENCES.

Integration of Explicit Functions.

40. THE Inverse Method of Differences has for its object to determine the primitive function from its given difference; or from given relations between it and its differences. We shall begin with the simplest case,

$$\Delta u_x = f(x),$$

in which it is required to determine a function whose difference is given explicitly in terms of the principal variable.

41. Since Δu_x is the difference of $u_x + C$, as well as of u_x , it will be necessary, in passing from the given difference Δu_x to the primitive function, to annex an arbitrary constant C , in order to give the result all the generality of which it is capable. Also C may be a function of x as well as an arbitrary constant, provided its value remains unaltered whilst x changes to $x + 1$. For if C_x denote such a function of x that $C_{x+1} = C_x$, or $\Delta C_x = 0$, we shall have

$$\Delta(u_x + C_x) = \Delta u_x.$$

It is evident that $C_x = \phi(2\lambda\pi x)$ has the property in question, ϕ denoting any trigonometrical function, sine, cosine, &c., and λ any integer. We shall see further on the importance of this remark.

42. The symbol Σ is used to denote the operation by which we pass from the difference Δu_x to the primitive function; so that

$$\Sigma(\Delta u_x) = u_x + \text{constant};$$

hence Σ and Δ denote operations the reverse of each other.

Also, as the same function admits of successive differences, so a function may be integrated any number of times; the second integral of u_x , or $\Sigma(\Sigma u_x)$, is written $\Sigma^2 u_x$, and the n^{th} integral $\Sigma^n u_x$.

If in the formula $\Sigma(\Delta u_x) = u_x$, we suppose the symbols of operation to be separated from those of quantity, we find $\Sigma \Delta u_x = u_x$; but on the same supposition $\Delta^{-1} \Delta u_x = \Delta^0 u_x = u_x$; so that Σ produces exactly the same effect as Δ^{-1} . Similarly Σ^n may be shewn to be identical with Δ^{-n} ; so that in any formula a negative power of Δ may be always replaced by the same positive power of Σ , and *vice versa*.

We now proceed to deduce the integrals of various expressions; chiefly, by reversing the processes given in Section I. for finding the differences of functions.

43. It is evident that $\Sigma(u_x + v_x + w_x) = \Sigma u_x + \Sigma v_x + \Sigma w_x$; for if we take the difference of both sides, we get the same result, viz. $u_x + v_x + w_x$. And in the same manner it appears that $\Sigma(au_x) = a\Sigma u_x$, and $\Sigma 0 = C$.

44. To find the integral of any rational integral function.

Since the difference of a rational integral function is a function of the same kind one dimension lower, it follows that the integral of a function of that description is a similar function one dimension higher; hence, to find the integral of

$$p_0 x^n + p_1 x^{n-1} + p_2 x^{n-2} + \dots + p_{n-1} x + p_n,$$

we may assume it equal to

$$ax^{n+1} + bx^n + \dots + kx + l;$$

then upon taking the difference of each side, and equating the coefficients of like powers of x , there will arise $n+1$ simple equations to determine the $n+1$ quantities a, b, c, \dots, k ; the last term l will remain indeterminate, being in fact the arbitrary constant which must be added to make the integral complete.

Ex. To find $\Sigma(x^4 + 1)$.

$$\text{Assume } \Sigma(x^4 + 1) = ax^5 + bx^4 + cx^3 + dx^2 + ex;$$

$$\begin{aligned} \therefore x^4 + 1 &= a(5x^4 + 10x^3 + 10x^2 + 5x + 1) + b(4x^3 + 6x^2 + 4x + 1) \\ &\quad + c(3x^2 + 3x + 1) + d(2x + 1) + e; \end{aligned}$$

$$\therefore 1 = 5a, \quad 0 = 10a + 4b, \quad 0 = 10a + 6b + 3c, \quad 0 = 5a + 4b + 3c + 2d,$$

$$1 = a + b + c + d + e.$$

$$\therefore a = \frac{1}{5}, \quad b = -\frac{1}{2}, \quad c = \frac{1}{3}, \quad d = 0, \quad e = \frac{29}{30},$$

$$\therefore \Sigma (x^4 + 1) = \frac{x^5}{5} - \frac{x^4}{2} + \frac{x^3}{3} + \frac{29}{30}x + C.$$

45. To find the integral of the product of consecutive terms of an arithmetic progression, we must annex one more factor at the beginning, and divide by the number of factors so increased and by the common difference.

For let $u_x = a + bx$, then we have seen (Art. 11) that

$$\Delta u_x u_{x+1} \dots u_{x+n} = u_{x+1} u_{x+2} \dots u_{x+n} \cdot (n+1) b,$$

therefore, taking the integrals of both sides, and writing $x-1$ for x , we get

$$\Sigma u_x u_{x+1} \dots u_{x+n-1} = \frac{u_{x-1} u_x \dots u_{x+n-1}}{(n+1) b} + C,$$

which proves the rule stated above.

$$\begin{aligned} \text{Ex. } \Sigma \left(2x + \frac{1}{2}\right) \left(2x + \frac{5}{2}\right) \left(2x + \frac{9}{2}\right) \\ = \frac{1}{8} \left(2x - \frac{3}{2}\right) \left(2x + \frac{1}{2}\right) \left(2x + \frac{5}{2}\right) \left(2x + \frac{9}{2}\right) + C. \end{aligned}$$

Each factor of an expression capable of being integrated by this rule, must be derivable from the preceding factor by changing x into $x+1$.

If one or more factors be deficient in a factorial of this kind, it may be resolved into others which are complete, as in the following instance;

$$\begin{aligned} (2x+1)(2x+5)(2x+7) &= (2x+3-2)(2x+5)(2x+7) \\ &= (2x+3)(2x+5)(2x+7) - 2(2x+5)(2x+7). \end{aligned}$$

46. A rational integral function may often be resolved into factorials of the above form, and in this way its integral more conveniently found, than by the method of Art. 44.

$$\text{Ex. 1. } x^3 + x^2 = x^2 (x + 1) = (x - 1 + 1) x (x + 1)$$

$$= (x - 1) x (x + 1) + x (x + 1);$$

$$\therefore \Sigma (x^3 + x^2) = \frac{(x - 2) (x - 1) x (x + 1)}{4} + \frac{(x - 1) x (x + 1)}{3} + C.$$

And in general, any quantity of the form

$$ax^n + bx^{n-1} + cx^{n-2} + \&c.$$

may be resolved into factorials, by the method of indeterminate coefficients; thus, if we assume

$$ax^2 + bx + c = A (x + 1) (x + 2) + B (x + 1) + C,$$

making $x = -1$, we get $a - b + c = C$;

$$\therefore a (x^2 - 1) + b (x + 1) = A (x + 1) (x + 2) + B (x + 1),$$

$$\text{or } a (x - 1) + b = A (x + 2) + B;$$

$$\text{make } x = -2, \therefore -3a + b = B;$$

$$\therefore a (x - 1) + 3a = A (x + 2), \therefore A = a.$$

In practice, however, it is generally easier to resolve a function by inspection, as in Ex. 1, than by this method, which is theoretically certain.

47. To find the integral of a fraction whose denominator is the product of consecutive terms of an arithmetic progression, and numerator constant, we must efface the last factor, divide by the number of factors remaining and by the common difference, and prefix a negative sign.

For let $u_x = a + bx$, then we have seen (Art. 12) that

$$\Delta \frac{c}{u_x u_{x+1} \dots u_{x+n-1}} = - \frac{nbc}{u_x u_{x+1} \dots u_{x+n}};$$

therefore taking the integral of both sides,

$$\Sigma \frac{c}{u_x u_{x+1} \dots u_{x+n}} = - \frac{c}{n b u_x u_{x+1} \dots u_{x+n-1}} + C,$$

which proves the rule just stated.

48. If the proposed fraction, instead of having its numerator constant, be

$$\frac{Ax^{n-2} + Bx^{n-3} + \dots + Kx + L}{u_x u_{x+1} \dots u_{x+n-1}},$$

(the degree of the numerator being at least lower by two units than that of the denominator,) we must reduce the numerator to a series of terms each of which is the product of consecutive factors reckoning from the beginning of the denominator; that is, assume

$$Ax^{n-2} + Bx^{n-3} + \dots + Kx + L = A' + B'u_x + C'u_x u_{x+1} + \dots \\ + K'u_x u_{x+1} \dots u_{x+n-3},$$

then, developing the second member, and equating coefficients of like powers of x , we obtain $n-1$ equations for determining A' , B' , C' , ... K' ; and the fraction resolves itself into the following, each of which is integrable,

$$\frac{A'}{u_x \dots u_{x+n-1}} + \frac{B'}{u_{x+1} \dots u_{x+n-1}} + \&c. + \frac{K'}{u_{x+n-2} \cdot u_{x+n-1}}.$$

If the degree of the numerator were the same as that of the denominator, or only lower by one unit than that of the denominator, we should arrive at a term $\frac{L'}{u_{x+n-1}}$, of which we are able to find the integral, only approximately.

Hence also, if any of the factors of the denominator of the fraction in Art. 47 be wanting, they may be supplied by introducing them into the numerator and denominator at the same time; and then the resulting fraction may be treated as in the present Article.

$$\begin{aligned}\text{Ex. 1. } \frac{1}{x^2-4} &= \frac{(x-1)x(x+1)}{(x-2)\dots(x+2)} \\ &= \frac{1}{(x+1)(x+2)} + \frac{3}{x(x+1)(x+2)} + \frac{6}{(x-1)x(x+1)(x+2)} \\ &\quad + \frac{6}{(x-2)(x-1)x(x+1)(x+2)},\end{aligned}$$

which is got by assuming

$$(x-1)x(x+1) = a(x-2)(x-1)x + b(x-2)(x-1) + c(x-2) + d,$$

and making $x = 2, 1, 0$, successively; taking care to reject the factor common to both sides, after each substitution.

$$\text{Ex. 2. } \Sigma \frac{1}{4x^2-9} = -\frac{1}{6} \frac{12x^2-12x-1}{(2x-3)(4x^2-1)}.$$

49. The fraction in the preceding Art. may be also integrated when the denominator is the product of any number of factors $x, x+mh, x+n, x+rh$, &c., m, n, r , &c., denoting whole numbers, and h the increment of x .

For by taking the difference of both members, we perceive the truth of the result (which, although expressed in a series the number of whose terms is variable, is often useful)

$$\Sigma \frac{M}{x(x+mh)} = -\frac{M}{mh} \left\{ \frac{1}{x} + \frac{1}{x+h} + \dots + \frac{1}{x+(m-1)h} \right\} \dots (1).$$

$$\text{But } \frac{x+a}{x(x+mh)(x+nh)} = \frac{A}{x(x+mh)} + \frac{B}{x(x+nh)} \dots (2),$$

$$\text{where } A+B=1, (An+Bm)h=a;$$

so that this fraction is integrable by formula (1). Next multiply both sides of (2) by $\frac{x+b}{x+rh}$; then the second member of the result can be resolved by (2) into fractions having a constant numerator, and the product of two simple factors for denominator, and is therefore integrable by (1); and so on to any number of factors, the dimension of the numerator being always less by at least two units than that of the denominator.

50. To find the integrals of a^x , and $\log v_x$.

We have seen (Art. 14) that $\Delta a^x = (a-1) a^x$;

$$\therefore \Sigma a^x = \frac{a^x}{a-1} + C. \quad \text{Also } \Sigma^n a^x = \frac{a^x}{(a-1)^n},$$

suppressing the part introduced by the constants, which would be a rational integral function of the $(n-1)^{\text{th}}$ degree, and might be represented by $\Sigma^n 0$, since $\Sigma 0 = C$, $\Sigma^2 0 = Cx + C'$, &c. The formula seems to fail when $a=1$, as it gives infinity for the value of Σa^x in that case, instead of x ; but if we give the constant C the form $C' - \frac{1}{a-1}$, then

$$\Sigma a^x = \frac{a^x - 1}{a - 1} + C';$$

now let $a = 1 + h$, where h is very small, then

$$\Sigma a^x = \frac{1 + hx + \frac{1}{2}x(x-1)h^2 + \&c. - 1}{h} = x + \frac{1}{2}x(x-1)h + \&c.;$$

therefore, when $a = 1$, $\Sigma a^x = x + C'$.

Next to find the integral of $\log v_x$.

If $u_x = \log (v_1 v_2 v_3 \dots v_{x-1})$, then

$$\Delta u_x = \log (v_1 v_2 \dots v_x) - \log (v_1 v_2 \dots v_{x-1}) = \log v_x;$$

$$\therefore \Sigma \log v_x = u_x + \log C = \log (C \cdot v_1 v_2 \dots v_{x-1}) = \log CP_{v_{x-1}},$$

using Pv_x to denote the product of all the successive values of the function v_x , from some fixed term v_1 (or more generally v_n , n being independent of x) to v_x inclusive.

51. To find the integrals of $\cos x\theta$, $\sin x\theta$.

$$\text{Since } \Delta \cos x\theta = -2 \sin \frac{1}{2}\theta \sin (x + \frac{1}{2})\theta,$$

$$\therefore \Delta \cos (x - \frac{1}{2})\theta = -2 \sin \frac{1}{2}\theta \sin x\theta;$$

$$\therefore \Sigma \sin x\theta = -\frac{\cos (x - \frac{1}{2})\theta}{2 \sin \frac{1}{2}\theta} + C.$$

Also since $\Delta^n \sin(x\theta + \alpha) = (2 \sin \frac{1}{2}\theta)^n \sin\{x\theta + \alpha + \frac{1}{2}n(\pi + \theta)\}$, (Art. 16), integrating both sides n times, and replacing α by $\alpha - \frac{1}{2}n(\pi + \theta)$, we get

$$\Sigma^n \sin(x\theta + \alpha) = \frac{\sin\{x\theta + \alpha - \frac{1}{2}n(\pi + \theta)\}}{(2 \sin \frac{1}{2}\theta)^n};$$

the same result as if in the value of $\Delta^n \sin(x\theta + \alpha)$ we had changed the sign of n ; as we should expect, Δ^{-n} being equivalent to Σ^n .

$$\text{Again, } \Delta \sin x\theta = 2 \sin \frac{1}{2}\theta \cos(x + \frac{1}{2})\theta,$$

$$\therefore \Delta \sin(x - \frac{1}{2})\theta = 2 \sin \frac{1}{2}\theta \cos x\theta;$$

$$\therefore \Sigma \cos x\theta = \frac{\sin(x - \frac{1}{2})\theta}{2 \sin \frac{1}{2}\theta} + C;$$

$$\text{and } \Sigma^n \cos(x\theta + \alpha) = \frac{\cos\{x\theta + \alpha - \frac{1}{2}n(\pi + \theta)\}}{(2 \sin \frac{1}{2}\theta)^n},$$

changing the sign of n in the value of $\Delta^n \cos(x\theta + \alpha)$ in Art. 16.

52. The preceding expressions may also be integrated by substituting for them their exponential values; as in the following instance.

$$\begin{aligned} \Sigma a^x \cos x\theta &= \frac{1}{2} \Sigma (a^x e^{x\theta\sqrt{-1}} + a^x e^{-x\theta\sqrt{-1}}) \\ &= \frac{1}{2} \frac{a^x e^{x\theta\sqrt{-1}}}{ae^{\theta\sqrt{-1}} - 1} + \frac{1}{2} \frac{a^x e^{-x\theta\sqrt{-1}}}{ae^{-\theta\sqrt{-1}} - 1} \\ &= \frac{a^x}{2} \frac{ae^{(x-1)\theta\sqrt{-1}} + ae^{-(x-1)\theta\sqrt{-1}} - e^{x\theta\sqrt{-1}} - e^{-x\theta\sqrt{-1}}}{a^2 - a(e^{\theta\sqrt{-1}} + e^{-\theta\sqrt{-1}}) + 1} \\ &= a^x \frac{a \cos(x-1)\theta - \cos x\theta}{a^2 - 2a \cos \theta + 1} + C. \end{aligned}$$

Hence, putting $a^2 - 2a \cos \theta + 1 = c$, and denoting

$$a^x \cos x\theta \text{ by } u_x, \quad \text{we have } c\Sigma u_x = a^2 u_{x-1} - u_x,$$

$$c^2 \Sigma^2 u_x = a^4 u_{x-2} - 2a^2 u_{x-1} + u_x;$$

and generally

$$c^n \Sigma^n u_x = a^{2n} u_{x-n} - na^{2n-2} u_{x-n+1} + \frac{n(n-1)}{1 \cdot 2} a^{2n-4} u_{x-n+2} - \&c. \pm u_x.$$

This result may be obtained immediately by separation of symbols; for we have

$$\Sigma u_x = \frac{1}{c} (a^2 u_{x-1} - u_x) = \frac{1}{c} (a^2 D^{-1} - 1) u_x;$$

$$\text{consequently } \Sigma^n u_x = \frac{1}{c^n} (a^2 D^{-1} - 1)^n u_x;$$

and this when developed produces the preceding result.

Exactly the same formulæ hold for $u_x = a^x \sin x\theta$.

In the same manner the integrals of $a^x (\sin x\theta)^m$, $a^x (\cos x\theta)^m$ may be obtained, when the powers of the sine and cosine of $x\theta$ have been replaced by the sines and cosines of multiples of $x\theta$.

$$53. \text{ To find the integral of } \frac{1}{\cos x\theta \cos (x+1)\theta}.$$

$$\text{Since } \Delta \tan x\theta = \frac{\sin \theta}{\cos x\theta \cos (x+1)\theta}, \quad (\text{Art. 16}^*.)$$

$$\text{we have } \Sigma \frac{1}{\cos x\theta \cos (x+1)\theta} = \frac{\tan x\theta}{\sin \theta} + C.$$

$$54. \text{ To find the integral of } \tan^{-1} \frac{1}{p + qx + rx^2}.$$

$$\text{Since } \Delta \tan^{-1} (a + bx) = \tan^{-1} \frac{b}{1 + (a+bx)(a+bx+b)}, \quad (\text{Art. 16}^*.)$$

$$\text{we may assume } \Sigma \tan^{-1} \frac{1}{p + qx + rx^2} = \tan^{-1} (a + bx),$$

and take the difference of both sides; then if the proposed function is capable of being integrated, the indeterminate coefficients a and b will become known. Also by differencing $n^x \tan^{-1} (\theta n^{-x})$ we may find an expression of which it is the integral, when $n = 2, 3, \&c.$

55. The integral of $a^x u_x u_{x+1} \dots u_{x+n-1}$, where $u_x = pa^x + q$, may be determined by assuming it equal to the same expression (only with another factor at the beginning instead of a^x) multiplied by an indeterminate coefficient, and taking the difference of both sides; for

$$\begin{aligned} \Delta u_{x-1} u_x \dots u_{x+n-1} &= u_x u_{x+1} \dots u_{x+n-1} (u_{x+n} - u_{x-1}) \\ &= a^x u_x u_{x+1} \dots u_{x+n-1} \cdot p (a^n - a^{-1}); \end{aligned}$$

$$\therefore \Sigma a^x u_x \dots u_{x+n-1} = \frac{a}{p(a^{n+1}-1)} \cdot u_{x-1} u_x \dots u_{x+n-1}.$$

Similarly, to find $\Sigma \frac{a^x}{u_x u_{x+1} \dots u_{x+n}}$, for the assumption we must efface the last factor in the denominator, and write instead of a^x an indeterminate coefficient; for

$$\Delta \frac{1}{u_x u_{x+1} \dots u_{x+n-1}} = - \frac{u_{x+n} - u_x}{u_x u_{x+1} \dots u_{x+n}} = - \frac{p a^x (a^n - 1)}{u_x u_{x+1} \dots u_{x+n}};$$

$$\therefore \Sigma \frac{a^x}{u_x u_{x+1} \dots u_{x+n}} = - \frac{1}{p(a^n - 1)} \cdot \frac{1}{u_x u_{x+1} \dots u_{x+n-1}};$$

56. In like manner, if $u_x = a + bx$, expressions of the forms

$$\frac{(p + qx) t^x}{u_x u_{x+1} \dots u_{x+n-1}}, \quad \frac{(p + qx + rx^2) t^x}{u_x u_{x+1} \dots u_{x+n-1}},$$

can sometimes be integrated, by assuming their integrals equal to expressions of the same form, except that the last factor in the denominator is effaced, and the polynomial in the numerator is replaced by another one dimension lower with indeterminate coefficients. It is of course only when a certain equation of condition between the quantities a, b, p, q, t is satisfied, that this method succeeds.

Ex. 1. Let $\Sigma \frac{x+1}{(2x-1)(2x+1)3^x} = \frac{A(\frac{1}{3})^x}{2x-1};$

$$\therefore \frac{x+1}{(2x-1)(2x+1)3^x} = A(\frac{1}{3})^{x+1} \left(\frac{1}{2x+1} - \frac{3}{2x-1} \right)$$

$$= A(\frac{1}{3})^{x+1} \frac{-4(x+1)}{(2x-1)(2x+1)};$$

$$\therefore A = -\frac{3}{4}, \text{ and } \Sigma u_x = C - \frac{3}{4} \frac{1}{(2x-1)3^x}.$$

Ex. 2. Let $\Sigma \frac{x^2 + 6x + 12}{x(x+1)(x+2)2^x} = \frac{A+Bx}{x(x+1)2^x},$

then $A = -6, \quad B = -2, \text{ and } \Sigma u_x = C - \frac{x+3}{x(x+1)2^{x-1}}.$

57. Since $\Delta(u_x v_x) = u_x \Delta v_x + v_{x+1} \Delta u_x$, we have

$$\Sigma(u_x \Delta v_x) = u_x v_x - \Sigma(v_{x+1} \Delta u_x),$$

the formula for integration by parts, corresponding to the formula

$$\int dx u \frac{dv}{dx} = uv - \int dx v \frac{du}{dx}.$$

Change v_x into Σv_x , and consequently Δv_x into v_x ,

$$\text{then } \Sigma(u_x v_x) = u_x \Sigma v_x - \Sigma(\Delta u_x \Sigma v_{x+1});$$

hence, by successive substitutions, we get

$$\Sigma(\Delta u_x \Sigma v_{x+1}) = \Delta u_x \Sigma^2 v_{x+1} - \Sigma(\Delta^2 u_x \Sigma^2 v_{x+2}),$$

$$\Sigma(\Delta^2 u_x \Sigma^2 v_{x+2}) = \Delta^2 u_x \Sigma^3 v_{x+2} - \Sigma(\Delta^3 u_x \Sigma^3 v_{x+3}),$$

.....

$$\Sigma(\Delta^n u_x \Sigma^n v_{x+n}) = \Delta^n u_x \Sigma^{n+1} v_{x+n} - \Sigma(\Delta^{n+1} u_x \Sigma^{n+1} v_{x+n+1});$$

$$\therefore \Sigma(u_x v_x) = u_x \Sigma v_x - \Delta u_x \Sigma^2 v_{x+1} + \Delta^2 u_x \Sigma^3 v_{x+2} \\ - \&c. \pm \Delta^n u_x \Sigma^{n+1} v_{x+n} \mp \Sigma(\Delta^{n+1} u_x \Sigma^{n+1} v_{x+n+1}).$$

58. The above formula, replacing in the second member Σ by its equivalent Δ^{-1} , may be written

$$\Sigma(u_x v_x) = u_x \Delta^{-1} v_x - \Delta u_x \cdot \Delta^{-2} (D v_x) + \Delta^2 u_x \cdot \Delta^{-3} (D^2 v_x) - \&c. \\ = (\Delta' + \Delta D')^{-1} u_x v_x,$$

if we restrict Δ to affect u_x only and Δ' and D' to affect v_x only; hence $\Sigma^2(u_x v_x) = (\Delta' + \Delta D')^{-2} u_x v_x$; and generally

$$\Sigma^n(u_x v_x) = (\Delta' + \Delta D')^{-n} u_x v_x \dots\dots\dots (1) \\ = (\Delta'^{-n} - n \Delta'^{-n-1} \Delta D' + \frac{n(n+1)}{1 \cdot 2} \Delta'^{-n-2} \Delta^2 D'^2 - \&c.) u_x v_x,$$

$$\text{or } \Sigma^n(u_x v_x) = u_x \Sigma^n v_x - n \Delta u_x \Sigma^{n+1} v_{x+1} \\ + \frac{n(n+1)}{1 \cdot 2} \Delta^2 u_x \Sigma^{n+2} v_{x+2} - \&c. \dots (2),$$

which may be also proved inductively, by shewing that the coefficients of the developments of $\Sigma^n(u_x v_x)$ and $(1+z)^{-n}$, which

are identical when $n=1$, undergo the same changes in passing from n to $n-1$. This appears from differencing both sides of equation (2). Both the formula (1) for $\Sigma^n(u_x v_x)$, and its development (2), result immediately, as we should expect, from the expressions for $\Delta^n(u_x v_x)$ in Art. 34, by changing the sign of n .

If in the formula $\Sigma^n(u_x v_x) = (\Delta' + \Delta D')^{-n} u_x v_x$ where Δ affects u_x only, and Δ' , D' affect v_x only, we suppose $v_x = a^x$; then

$$\Delta' v_x = (a-1) a^x, \quad D' v_x = a \cdot a^x;$$

and the formula becomes

$$\Sigma^n(u_x a^x) = (a-1 + \Delta a)^{-n} u_x a^x = a^x (Da-1)^{-n} u_x,$$

which agrees with the result in Art. 34 when n has its sign changed.

If in this last formula we change a into $\frac{1}{a}$, we get

$$(D-a)^{-n} u_x = a^{x-n} \Sigma^n(u_x a^{-x}),$$

$$\text{and } (D-a)^{-1} u_x = a^{x-1} \Sigma(u_x a^{-x}).$$

59. The above formula for $\Sigma^n(u_x v_x)$ always enables us to find the integrals of functions made up of two factors, one of which leads to zero, as the value of one of its successive differences, and the other admits of successive integrations. Suppose, for example, that u_x is a rational integral function of the n^{th} degree, and that $v_x = a^x$; then

$$\Sigma v_x = \frac{a^x}{a-1}, \quad \Sigma^2 v_{x+1} = \frac{a^{x+1}}{(a-1)^2}, \quad \&c.; \quad \Delta^n u_x = \text{const}, \quad \Delta^{n+1} u_x = 0;$$

$$\therefore \Sigma(u_x a^x) = \frac{u_x a^x}{a-1} - \frac{\Delta u_x a^{x+1}}{(a-1)^2} + \frac{\Delta^2 u_x a^{x+2}}{(a-1)^3} - \dots \pm \frac{\Delta^n u_x a^{x+n}}{(a-1)^{n+1}} + C.$$

Again, suppose u_x to be a rational integral function of the n^{th} degree, and $v_x = \cos x\theta$; then taking the value of $\Sigma^n v_x$ from Art. 51,

$$\begin{aligned} \Sigma(u_x \cos x\theta) &= \frac{u_x \cos \{x\theta - \frac{1}{2}(\pi + \theta)\}}{2 \sin \frac{1}{2}\theta} \\ &- \Delta u_x \frac{\cos \{(x+1)\theta - (\pi + \theta)\}}{(2 \sin \frac{1}{2}\theta)^2} + \Delta^2 u_x \frac{\cos \{(x+2)\theta - \frac{3}{2}(\pi + \theta)\}}{(2 \sin \frac{1}{2}\theta)^3} - \&c., \end{aligned}$$

the series terminating with $\Delta^n u_x$. Similarly, if $v_x = a^x \cos x\theta$; or if $v_x = a^x \cos^n x\theta \cdot \sin^n x\theta$, since the product $\cos^n x\theta \cdot \sin^n x\theta$ may be replaced by simple dimensions of sines and cosines of multiples of $x\theta$; and it will be noticed that the fraction of Art. 48 may be brought under this case.

60. Since the performance of the operation Σ upon any series of terms $A\Delta^m u_x + B\Delta^n u_x + \dots$, reduces it to

$$A\Delta^{m-1} u_x + B\Delta^{n-1} u_x + \dots;$$

it appears that prefixing Σ to $(A\Delta^m + B\Delta^n + \dots) u_x$ has the same effect as prefixing Δ^{-1} ; in other words, Σ is equivalent to Δ^{-1} . And in like manner, since integrating $\Delta^m u_x$ n times, reduces it to $\Delta^{m-n} u_x$, Σ^n must be equivalent to Δ^{-n} .

The same reasoning is applicable to the symbols $(\int dx)^n, \left(\frac{d}{dx}\right)^n$; whenever therefore, in separating the symbols of operation from those of quantity, as in the expression $F(\Delta)u_x, f\left(\frac{d}{dx}\right)u_x$, terms containing negative powers of Δ and $\frac{d}{dx}$ occur, they must be understood to be replaced by the corresponding positive powers of Σ and $\int dx$. This being premised, we proceed to investigate a general series for Σu_x ; preparatory to which the following propositions must be proved.

61. To determine the generating functions of

$$\Sigma^n u_x, \quad \frac{d^n u_x}{dx^n}, \quad \int^n dx^n u_x,$$

from that of u_x .

By virtue of the relations

$$G(\Delta u_x) = \left(\frac{1}{t} - 1\right) G u_x,$$

$$G(\Delta^n u_x) = \left(\frac{1}{t} - 1\right)^n G u_x, \dots\dots\dots (1)$$

$$\text{we have } \left(\frac{1}{t} - 1\right) G(\Sigma u^x) = G(\Delta \Sigma u_x) = G u_x;$$

$$\therefore G(\Sigma u_x) = \left(\frac{1}{t} - 1\right)^{-1} G u_x,$$

$$\text{and } G(\Sigma^n u_x) = \left(\frac{1}{t} - 1\right)^{-n} G u_x,$$

the same result as if we had changed the sign of n in (1) and replaced Δ^{-n} by Σ^n .

62. Again, since

$$G u_x = \dots + u_x t^x + \dots + u_{x+h} t^{x+h} + \dots;$$

$$\therefore \dots + (u_{x+h} - u_x) t^x + \dots = \left(\frac{1}{t^h} - 1\right) G u_x$$

$$= \left\{1 - h \log t + \frac{h^2}{1 \cdot 2} (\log t)^2 - \&c. - 1\right\} G u_x;$$

therefore, dividing both sides by h and then making $h = 0$, which we are at liberty to do since h is indeterminate,

$$\dots + \left\{ \frac{u_{x+h} - u_x}{h} \right\}_{h=0} t^x + \dots = -\log t G u_x = \log \frac{1}{t} \cdot G u_x,$$

$$\text{or } \dots + \frac{du_x}{dx} \cdot t^x + \dots = \log \frac{1}{t} \cdot G u_x;$$

$$\therefore G\left(\frac{du_x}{dx}\right) = \log \frac{1}{t} \cdot G u_x,$$

$$\text{and } G\left(\frac{d^n u_x}{dx^n}\right) = \left(\log \frac{1}{t}\right)^n G u_x \dots \dots \dots (1).$$

$$\text{Again, } \log \frac{1}{t} G \left(\int dx u_x \right) = G \left(\frac{d}{dx} \int dx u_x \right) = G u_x;$$

$$\therefore G \left(\int dx u_x \right) = \left(\log \frac{1}{t} \right)^{-1} G u_x, \text{ and } G \left(\int^n dx^n u_x \right) = \left(\log \frac{1}{t} \right)^{-n} G u_x;$$

the same result as if we had changed the sign of n in (1) and replaced $\left(\frac{d}{dx}\right)^{-n}$ by $(\int dx)^n$.

These values for $G\left(\frac{d^n u_x}{dx^n}\right)$, $G(\int^n dx^n u_x)$, may be immediately obtained from Art. 28; for if in the formula

$$G\{\psi(D) u_x\} = \psi\left(\frac{1}{t}\right) \cdot G u_x,$$

we suppose $\psi\left(\frac{1}{t}\right) = \left(\log \frac{1}{t}\right)^n$, then

$$\psi(D) = (\log D)^n = (\log e^{\frac{d}{dx}})^n = \left(\frac{d}{dx}\right)^n,$$

$$\therefore G\left(\frac{d^n u_x}{dx^n}\right) = \left(\log \frac{1}{t}\right)^n \cdot Gu_x;$$

and changing the sign of n , and replacing $\left(\frac{d}{dx}\right)^n$ by $(\int dx)^n$,

$$G(\int^n dx^n u_x) = \left(\log \frac{1}{t}\right)^{-n} \cdot Gu_x.$$

63. To investigate a general series for Σu_x , involving only $\int dx u_x$, u_x , and the differential coefficients of u_x .

$$\begin{aligned} G(\Sigma u_x) &= \left(\frac{1}{t} - 1\right)^{-1} Gu_x = (e^{\log \frac{1}{t}} - 1)^{-1} Gu_x \\ &= \left(\log \frac{1}{t}\right)^{-1} \cdot \frac{\log \frac{1}{t}}{e^{\log \frac{1}{t}} - 1} \cdot Gu_x \\ &= \left(\log \frac{1}{t}\right)^{-1} \left\{1 - \frac{1}{2} \log \frac{1}{t} + \frac{B_1}{1 \cdot 2} \left(\log \frac{1}{t}\right)^2 - \&c. \right. \\ &\quad \left. + (-1)^{n+1} \frac{B_{2n-1}}{[2n]} \left(\log \frac{1}{t}\right)^{2n} + \dots \right\} Gu_x, \end{aligned}$$

assuming, as will be proved in the next Art., that $\frac{v}{e^v - 1}$ can be expanded in a series of the same form as that within brackets, and denoting by

$$\frac{B_1}{1 \cdot 2}, \quad \frac{-B_3}{1 \cdot 2 \cdot 3 \cdot 4}, \quad \&c., \quad (-1)^{n+1} \frac{B_{2n-1}}{[2n]},$$

the coefficients of $v^2, v^4, \dots v^{2n}$ in that expansion. Hence

$$G(\Sigma u_x) = G(\int dx u_x) - \frac{1}{2} Gu_x + \frac{B_1}{1 \cdot 2} G\left(\frac{d u_x}{dx}\right) - \&c.$$

$$= G \left\{ f dx u_x - \frac{1}{2} u_x + \frac{B_1}{1 \cdot 2} \frac{du_x}{dx} - \frac{B_3}{[4]} \frac{d^3 u_x}{dx^3} + \&c. \right\};$$

$$\begin{aligned} \therefore \Sigma u_x &= f dx u_x - \frac{1}{2} u_x + \frac{B_1}{1 \cdot 2} \frac{du_x}{dx} - \frac{B_3}{[4]} \frac{d^3 u_x}{dx^3} + \dots \\ &\quad + (-1)^{n+1} \frac{B_{2n-1}}{[2n]} \frac{d^{2n-1} u_x}{dx^{2n-1}} + \dots \end{aligned}$$

$$\begin{aligned} \text{Ex. 1. } \Sigma x^m &= \frac{x^{m+1}}{m+1} - \frac{1}{2} x^m + \frac{1}{2} B_1 m x^{m-1} \\ &\quad - \frac{1}{4} B_3 \frac{m(m-1)(m-2)}{1 \cdot 2 \cdot 3} x^{m-3} + \&c. + C. \end{aligned}$$

$$\text{Hence } \Sigma x^4 = \frac{x^5}{5} - \frac{x^4}{2} + \frac{x^3}{3} - \frac{x}{30} + C;$$

$$\Sigma x^5 = \frac{x^6}{6} - \frac{x^5}{2} + \frac{5x^4}{12} - \frac{x^2}{12} + C.$$

$$\begin{aligned} \text{Ex. 2. } \Sigma \frac{1}{a+bx} &= \frac{1}{b} \log(a+bx) - \frac{1}{2(a+bx)} \\ &\quad - \frac{B_1 b}{2(a+bx)^2} + \frac{B_3 b^3}{4(a+bx)^4} - \&c. + C. \end{aligned}$$

Similarly $\Sigma^n u_x$ may be expressed in terms of the integrals and differential coefficients of u_x by making n negative in the formula of Art. 35, and expanding $e^{\frac{d}{dx}}$, which gives

$$\Sigma^n u_x = \left(\frac{d}{dx} + \frac{1}{1 \cdot 2} \frac{d^2}{dx^2} + \frac{1}{1 \cdot 2 \cdot 3} \frac{d^3}{dx^3} + \dots \right)^{-n} u_x.$$

Numbers of Bernoulli.

64. The numbers B_1, B_3, \dots which are required in the general value of Σu_x in the preceding Art., are called the Numbers of Bernoulli, and are of great importance in the theory of Series. They are defined by the equation

$$\frac{t}{e^t - 1} = 1 - \frac{1}{2} t + \frac{B_1}{1 \cdot 2} t^2 - \frac{B_3}{[4]} t^4 + \dots + (-1)^{n+1} \frac{B_{2n-1}}{[2n]} t^{2n} \pm \&c.,$$

and their values may be computed in the following manner.

$$\text{If } \phi(t) = \frac{t}{e^t - 1}, \text{ then } \phi(-t) = \frac{-t}{e^{-t} - 1} = \frac{te^t}{e^t - 1},$$

$$\therefore \phi(t) - \phi(-t) = -t;$$

which shews that the only term in the development of $\phi(t)$ which involves an odd power of t is $-\frac{1}{2}t$; for if any higher odd power entered, it would occur in $\phi(t) - \phi(-t)$ with its coefficient doubled; we may assume therefore for $\phi(t)$ the form of development given above, viz.

$$\frac{t}{e^t - 1} = 1 - \frac{1}{2}t + \frac{B_1}{1 \cdot 2}t^2 - \frac{B_3}{4}t^4 + \dots + (-1)^{n+1} \frac{B_{2n-1}}{2n} t^{2n} \pm \&c. \quad (1).$$

$$\begin{aligned} \text{Now } \frac{t}{e^t - 1} &= \frac{\log \{1 + (e^t - 1)\}}{e^t - 1} = 1 - \frac{1}{2}(e^t - 1) + \frac{1}{3}(e^t - 1)^2 - \dots \\ &\quad + \frac{1}{2n+1}(e^t - 1)^{2n} - \&c., \end{aligned}$$

and if A_{2n} be the coefficient of t^{2n} in this development of the second member,

$$\begin{aligned} A_{2n} &= -\frac{1}{2} \frac{1^{2n}}{2n} + \frac{1}{3} \cdot \frac{2^{2n} - 2 \cdot 1^{2n}}{2n} - \&c. \\ &\quad + \frac{1}{2n+1} \cdot \frac{(2n)^{2n} - 2n(2n-1)^{2n} + \dots}{2n} - \&c. \\ &= \frac{1}{2n} \left\{ -\frac{1}{2} \Delta 0^{2n} + \frac{1}{3} \cdot \Delta^2 0^{2n} - \&c. + \frac{1}{2n+1} \Delta^{2n} 0^{2n} \right\}; \end{aligned}$$

all the terms after $\Delta^{2n} 0^{2n}$ vanishing, since $\Delta^m 0^{2n}$ is zero when $m > 2n$. Hence

$$B_{2n-1} = (-1)^{n+1} \left\{ -\frac{1}{2} \Delta 0^{2n} + \frac{1}{3} \Delta^2 0^{2n} - \&c. \dots + \frac{1}{2n+1} \Delta^{2n} 0^{2n} \right\}.$$

By this formula B_1, B_3, B_5 , &c. may be readily computed, supposing the numbers comprised in the form $\Delta^m 0^{2n}$, or the differences of zero to be known; we find

$$B_1 = \frac{1}{6}, \quad B_3 = \frac{1}{30}, \quad B_5 = \frac{1}{42}, \quad B_7 = \frac{1}{30}, \quad B_9 = \frac{5}{66}, \quad \&c.$$

values that may be easily verified by applying Maclaurin's Theorem to obtain the development (1).

65. Also, since

$$\Delta^n 0^{2n} = m^{2n} - m(m-1)^{2n} + \frac{m(m-1)}{1 \cdot 2} (m-2)^{2n} - \&c. \quad (\text{Art. 20})$$

we may, if we please, eliminate the differences of zero from the expression for B_{2n-1} ; and we find

$$(-1)^{n+1} B_{2n-1} = -\frac{1}{2} + \frac{1}{3} (2^{2n} - 2) - \frac{1}{4} (3^{2n} - 3 \cdot 2^{2n} + 3) + \&c. + \frac{\lfloor 2n}{2n+1}.$$

Besides serving to express the general value of Σu_x , the numbers of Bernoulli have various other uses, of which we shall now give one or two of the most remarkable. Any function that can be put under the form $t \div (e^t - 1)$ may be expanded in terms involving those numbers.

66. To find the general terms of the expansions of $\cot \theta$ and $\tan \theta$ in powers of θ .

$$\cot \theta = \sqrt{-1} \left(1 + \frac{2}{e^{2\theta\sqrt{-1}} - 1} \right) = \sqrt{-1} + \frac{1}{\theta} \cdot \frac{2\theta\sqrt{-1}}{e^{2\theta\sqrt{-1}} - 1}.$$

Now the general term of the expansion of

$$\frac{2\theta\sqrt{-1}}{e^{2\theta\sqrt{-1}} - 1} \text{ is } (-1)^{n+1} \frac{B_{2n-1}}{\lfloor 2n} (2\theta\sqrt{-1})^{2n}, \text{ or } -\frac{2^{2n} B_{2n-1}}{\lfloor 2n} \theta^{2n};$$

\therefore the general term of the expansion of $\cot \theta$ is $-\frac{2^{2n} B_{2n-1}}{\lfloor 2n} \theta^{2n-1}$,

$$\text{and } \cot \theta = \frac{1}{\theta} - \frac{2^2}{1 \cdot 2} B_1 \theta - \frac{2^4 B_3}{1 \cdot 2 \cdot 3 \cdot 4} \theta^3 - \&c.$$

Also since $\tan \theta = \cot \theta - 2 \cot 2\theta$,

the general term of the series for $\tan \theta$ is

$$-\frac{2^{2n} B_{2n-1} \theta^{2n-1}}{\lfloor 2n} + 2 \cdot \frac{2^{2n} B_{2n-1} (2\theta)^{2n-1}}{\lfloor 2n} = \frac{2^{2n} (2^{2n} - 1) B_{2n-1}}{\lfloor 2n} \theta^{2n-1};$$

$$\therefore \tan \theta = \frac{4 \cdot 3}{1 \cdot 2} B_1 \theta + \frac{16 \cdot 15}{1 \cdot 2 \cdot 3 \cdot 4} B_3 \theta^3 + \frac{2^6 (2^6 - 1)}{1 \cdot 2 \dots 6} B_5 \theta^5 + \&c.$$

Hence, by differentiating the above expressions for $\cot \theta$ and $\tan \theta$, we may deduce the general terms of the expansions of

$\cot^2 \theta$ and $\tan^2 \theta$; and by integrating them, the general terms of the expansions of $\log \sin \theta$, $\log \cos \theta$.

67. To find S the sum of the infinite series

$$\frac{1}{1^{2n}} + \frac{1}{2^{2n}} + \frac{1}{3^{2n}} + \dots$$

$$\text{Since } \sin \theta = \theta \left(1 - \frac{\theta^2}{\pi^2}\right) \left(1 - \frac{\theta^2}{2^2 \pi^2}\right) \left(1 - \frac{\theta^2}{3^2 \pi^2}\right) \dots$$

changing θ into $\pi\theta$, we get

$$\sin \pi\theta = \pi\theta \left(1 - \frac{\theta^2}{1^2}\right) \left(1 - \frac{\theta^2}{2^2}\right) \left(1 - \frac{\theta^2}{3^2}\right) \dots$$

therefore, differentiating the logarithm of each member,

$$\begin{aligned} \pi \cot \pi\theta &= \frac{1}{\theta} - \frac{2\theta}{1^2} \left(1 - \frac{\theta^2}{1^2}\right)^{-1} - \frac{2\theta}{2^2} \left(1 - \frac{\theta^2}{2^2}\right)^{-1} - \frac{2\theta}{3^2} \left(1 - \frac{\theta^2}{3^2}\right)^{-1} - \&c. \\ &= \frac{1}{\theta} - 2\theta \left(\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots\right) - 2\theta^3 \left(\frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \dots\right) - \&c. \\ &\quad - 2\theta^{2n-1} \left(\frac{1}{1^{2n}} + \frac{1}{2^{2n}} + \frac{1}{3^{2n}} + \dots\right) - \dots \end{aligned}$$

But the coefficient of θ^{2n-1} in the expansion of $\pi \cot \pi\theta$ is

$$\begin{aligned} &= -\frac{2^{2n} B_{2n-1} \pi^{2n}}{2n}, \\ \therefore S &= \frac{2^{2n-1} B_{2n-1} \pi^{2n}}{1 \cdot 2 \cdot 3 \dots 2n}. \end{aligned}$$

$$\text{Hence } \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots = \frac{\pi^2}{6}, \quad \frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \dots = \frac{\pi^4}{90}.$$

Calling S_{2n} the sum of this infinite series, we have

$$\frac{S_{2n}}{S_{2n+2}} = \frac{(2n+1)(2n+2)}{4\pi^2} \frac{B_{2n-1}}{B_{2n+1}}.$$

Now suppose n very great, then $\frac{B_{2n+1}}{B_{2n-1}} = \frac{n^2}{\pi^2}$, which proves the divergency of the series formed by the numbers of Bernoulli; these numbers increase very rapidly, beginning with B_{19} .

68. To find S the sum of the infinite series

$$\frac{1}{1^{2n}} + \frac{1}{3^{2n}} + \frac{1}{5^{2n}} + \dots$$

$$\text{Since } \cos \theta = \left(1 - \frac{4\theta^2}{\pi^2}\right) \left(1 - \frac{4\theta^2}{3^2\pi^2}\right) \left(1 - \frac{4\theta^2}{5^2\pi^2}\right) \dots$$

$$\text{changing } \theta \text{ into } \frac{\pi\theta}{2}, \text{ we get } \cos \frac{\pi\theta}{2} = \left(1 - \frac{\theta^2}{1^2}\right) \left(1 - \frac{\theta^2}{3^2}\right) \left(1 - \frac{\theta^2}{5^2}\right) \dots$$

therefore, differentiating the logarithm of each member,

$$\frac{\pi}{2} \tan \frac{\pi\theta}{2} = \frac{2\theta}{2} \left(1 - \frac{\theta^2}{1^2}\right)^{-1} + \frac{2\theta}{3^2} \left(1 - \frac{\theta^2}{3^2}\right)^{-1} + \frac{2\theta}{5^2} \left(1 - \frac{\theta^2}{5^2}\right)^{-1} + \&c.;$$

and equating the coefficients of θ^{2n-1} in each member, which are respectively,

$$\left(\frac{\pi}{2}\right)^{2n} B_{2n-1} \frac{2^{2n} (2^{2n} - 1)}{2n}, \text{ and } 2S,$$

$$\text{we get } S = \frac{1}{2} B_{2n-1} \frac{\pi^{2n} (2^{2n} - 1)}{1 \cdot 2 \cdot 3 \dots 2n}.$$

$$\text{Hence } \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \&c. = \frac{1}{2} \cdot \frac{1}{6} \cdot \frac{3\pi^2}{1 \cdot 2} = \frac{\pi^2}{8}.$$

$$69. \text{ Also } \frac{1}{1^{2n}} - \frac{1}{2^{2n}} + \frac{1}{3^{2n}} - \frac{1}{4^{2n}} + \dots$$

$$= \frac{1}{1^{2n}} + \frac{1}{3^{2n}} + \frac{1}{5^{2n}} + \dots - \frac{1}{2^{2n}} \left(\frac{1}{1^{2n}} + \frac{1}{2^{2n}} + \frac{1}{3^{2n}} + \dots \right)$$

$$= \frac{1}{2} \frac{2^{2n} - 1}{2n} \pi^{2n} B_{2n-1} - \frac{1}{2^{2n}} \frac{2^{2n-1} \pi^{2n}}{2n} B_{2n-1}$$

$$= \frac{(2^{2n-1} - 1) \pi^{2n} B_{2n-1}}{2n}.$$

Since $\log (1 + e^{-z}) = e^{-z} - \frac{1}{2} e^{-2z} + \frac{1}{3} e^{-3z} - \&c.$,

integrating this $2n - 1$ times between the limits $z = 0$, $z = \infty$, we find for result the series just summed;

$$\therefore (\int dz_0^\infty)^{2n-1} \log (1 + e^{-z}) = \frac{(2^{2n-1} - 1) \pi^{2n} B_{2n-1}}{|2n|}.$$

70. To find an approximate value of

$$\Gamma(x+1) = 1.2.3 \dots x, \text{ when } x \text{ is very large.}$$

Making $u_x = \log x$ in the formula

$$\Sigma u_x = \int dx u_x - \frac{1}{2} u_x + \frac{B_1}{1.2} \frac{du_x}{dx} - \&c.$$

we find, (Art. 50) adding $\log x$ to both sides,

$$\begin{aligned} \log \{1.2.3 \dots (x-1)x\} &= C + x \log x - x - \frac{1}{2} \log x \\ &\quad + \frac{B_1}{1.2} \frac{1}{x} - \frac{B_3}{3.4} \frac{1}{x^3} + \dots + \log x \\ &= C + (x + \frac{1}{2}) \log x - x + \log (1+h), \end{aligned}$$

$$\text{putting } \log (1+h) = \frac{B_1}{1.2x} - \frac{B_3}{3.4x^3} + \frac{B_5}{5.6x^5} - \&c.,$$

so that h is a quantity continually approaching to zero as x increases.

Now to determine C , suppose x very large so that h may be neglected, and change x into $2x$, then

$$\begin{aligned} \log (1.2.3 \dots 2x) &= C + (2x + \frac{1}{2}) \log 2x - 2x \\ &= C + (2x + \frac{1}{2}) (\log x + \log 2) - 2x, \end{aligned}$$

$$\text{and } \log (2.4.6 \dots 2x) = \log (2^x.1.2.3 \dots x)$$

$$= x \log 2 + C + (x + \frac{1}{2}) \log x - x;$$

$$\therefore \log \{1.3.5 \dots (2x-1)\} = x \log x + (x + \frac{1}{2}) \log 2 - x;$$

$$\begin{aligned}\therefore \log \frac{2 \cdot 4 \cdot 6 \dots 2x}{1 \cdot 3 \cdot 5 \dots (2x-1)} &= C + \frac{1}{2} \log x - \frac{1}{2} \log 2 \\ &= C + \frac{1}{2} \log 2x - \log 2;\end{aligned}$$

$$\begin{aligned}\therefore 2C - 2 \log 2 &= 2 \log \frac{2 \cdot 4 \cdot 6 \dots (2x)}{1 \cdot 3 \cdot 5 \dots (2x-1)} - \log 2x \\ &= \log \frac{2 \cdot 2 \cdot 4 \cdot 4 \cdot 6 \cdot 6 \dots (2x-2) \cdot 2x}{1 \cdot 3 \cdot 3 \cdot 5 \cdot 5 \cdot 7 \dots (2x-1) (2x-1)} \\ &= \log \frac{\pi}{2}, \text{ by Wallis's Theorem,}\end{aligned}$$

since x is indefinitely large;

$$\therefore C = \frac{1}{2} \log 2\pi;$$

$$\begin{aligned}\therefore \log (1 \cdot 2 \cdot 3 \dots x) &= \frac{1}{2} \log 2\pi + (x + \frac{1}{2}) \log x - x + \log (1+h) \\ &= \log \sqrt{2\pi x} + \log \left(\frac{x}{e}\right)^x + \log (1+h); \\ \therefore 1 \cdot 2 \cdot 3 \dots x &= \sqrt{2\pi x} \cdot \left(\frac{x}{e}\right)^x \cdot (1+h),\end{aligned}$$

where h is to be calculated from the series

$$\log (1+h) = \frac{B_1}{1 \cdot 2} \cdot \frac{1}{x} - \frac{B_3}{3 \cdot 4} \cdot \frac{1}{x^3} + \frac{B_5}{5 \cdot 6} \cdot \frac{1}{x^5} - \&c.,$$

and in general it will suffice to take the first term only, which gives

$$1 \cdot 2 \cdot 3 \dots x = \sqrt{2\pi x} x^x e^{-x + \frac{1}{12x}};$$

or, more accurately, $1 \cdot 2 \cdot 3 \dots x = \sqrt{2\pi x} x^x e^{-x + \frac{\mu}{12x}}$, where μ denotes a quantity lying between 0 and 1.

OBS. The preceding series, even for large values of x , becomes divergent after a certain number of terms; this will happen after n terms if

$$\begin{aligned}\frac{B_{2n+1}}{(2n+1)(2n+2)} \cdot \frac{1}{x^{2n+1}} &> \frac{B_{2n-1}}{(2n-1)2n} \cdot \frac{1}{x^{2n-1}}, \\ \text{or } \frac{B_{2n+1}}{B_{2n-1}} &> \frac{(2n+1)(2n+2)}{(2n-1)2n} x^2,\end{aligned}$$

but the first member of this inequality never exceeds

$$\frac{(2n+1)(2n+2)}{4\pi^2}, \text{ (Art. 67.)}$$

$$\therefore (2n-1)2n > 4\pi^2 x^2, \text{ or } n > \pi x.$$

It can, however, be proved that an approximate value of the series will be obtained by taking the aggregate of the convergent terms only, as will be seen in the next Article.

71. An approximate value of Σu_x will be obtained by taking the aggregate of the converging terms only, in the series for Σu_x , involving $\int dx u_x$, u_x , and the differential coefficients of u_x ; and the error will be less than the last of the convergent, or the first of the divergent terms.

We have by Art. 63, (omitting the index of u_x in the second member,)

$$\begin{aligned} \Sigma u_x = C + \int dx u - \frac{1}{2} u + \frac{B_1}{1 \cdot 2} \frac{du}{dx} - \frac{B_2}{4} \frac{d^2 u}{dx^2} + \dots \\ + (-1)^{n-1} \frac{B_{2n-1}}{2n} \frac{d^{2n-1} u}{dx^{2n-1}} + (-1)^n 2R_n, \quad (1) \end{aligned}$$

$$\text{where } 2R_n = \frac{B_{2n+1}}{2n+2} \frac{d^{2n+1} u}{dx^{2n+1}} - \frac{B_{2n+3}}{2n+4} \frac{d^{2n+3} u}{dx^{2n+3}} + \&c.;$$

or since in general, by Art. 67,

$$\begin{aligned} \frac{B_{2n-1}}{2n} = 2 \left\{ \frac{1}{(2\pi)^{2n}} + \frac{1}{(4\pi)^{2n}} + \frac{1}{(6\pi)^{2n}} + \&c. \right\}; \\ R_n = \left\{ \frac{1}{(2\pi)^{2n+2}} + \frac{1}{(4\pi)^{2n+2}} + \frac{1}{(6\pi)^{2n+2}} + \dots \right\} \frac{d^{2n+1} u}{dx^{2n+1}} \\ - \left\{ \frac{1}{(2\pi)^{2n+4}} + \frac{1}{(4\pi)^{2n+4}} + \frac{1}{(6\pi)^{2n+4}} + \dots \right\} \frac{d^{2n+3} u}{dx^{2n+3}} \\ + \left\{ \frac{1}{(2\pi)^{2n+6}} + \frac{1}{(4\pi)^{2n+6}} + \frac{1}{(6\pi)^{2n+6}} + \dots \right\} \frac{d^{2n+5} u}{dx^{2n+5}} \\ - \&c.; \end{aligned}$$

or, adding the terms vertically, and calling the resulting series $v_1, v_2, \&c.$,

$$R_n = v_1 + v_2 + v_3 + \dots + v_m + \dots$$

$$\text{where } v_m = \frac{1}{(2m\pi)^{2n+2}} \frac{d^{2n+1}u}{dx^{2n+1}} - \frac{1}{(2m\pi)^{2n+4}} \frac{d^{2n+3}u}{dx^{2n+3}}$$

$$+ \frac{1}{(2m\pi)^{2n+6}} \frac{d^{2n+5}u}{dx^{2n+5}} - \&c.;$$

$$\therefore \frac{d^2 v_m}{dx^2} = - (2m\pi)^2 \left\{ v_m - \frac{1}{(2m\pi)^{2n+2}} \frac{d^{2n+1}u}{dx^{2n+1}} \right\},$$

$$\text{or } \frac{d^2 v_m}{dx^2} + (2m\pi)^2 v_m = \frac{1}{(2m\pi)^{2n}} \frac{d^{2n+1}u}{dx^{2n+1}},$$

and integrating this by the method of parameters, as in Art. 72,

$$v_m = - \frac{1}{(2m\pi)^{2n+1}} \int dx \left(\sin 2m\pi x \cdot \frac{d^{2n+1}u}{dx^{2n+1}} \right),$$

since, x being an integer, the term multiplied by $\sin 2m\pi x$ disappears, and $\cos 2m\pi x = 1$; and the arbitrary constant is unnecessary, being already introduced in equation (1);

$$\therefore R_n = - \int dx \left\{ \frac{\sin 2x\pi}{(2\pi)^{2n+1}} + \frac{\sin 4x\pi}{(4\pi)^{2n+1}} + \frac{\sin 6x\pi}{(6\pi)^{2n+1}} + \&c. \right\} \frac{d^{2n+1}u}{dx^{2n+1}};$$

therefore, numerically, $2R_n$ is less than

$$- 2 \int dx \left\{ \frac{1}{(2\pi)^{2n}} + \frac{1}{(4\pi)^{2n}} + \frac{1}{(6\pi)^{2n}} + \dots \right\} \frac{d^{2n+1}u}{dx^{2n+1}} < - \frac{B_{2n-1}}{2n} \frac{d^{2n}u}{dx^{2n}};$$

which last quantity lies between $\frac{B_{2n-1}}{2n} \frac{d^{2n-1}u}{dx^{2n-1}}$ and $\frac{B_{2n+1}}{2n+2} \frac{d^{2n+1}u}{dx^{2n+1}}$,

if these be the last of the convergent and first of the divergent terms respectively of the series for Σu_x . Consequently the sum of all the divergent terms in the series for Σu_x is less than the last of the convergent or the first of the divergent terms.

SECTION III.

EQUATIONS OF DIFFERENCES.

72. WE now come to the case in which the relation between the principal variable and any function of it is to be determined by means of an equation between x , u_x , and one or more of the successive values, or differences, of u_x ; that is, from equations of the form

$$F(x, u_x, u_{x+1}, \dots u_{x+n}) = 0,$$

$$\text{or, } f(x, u_x, \Delta u_x, \dots \Delta^n u_x) = 0,$$

since, by the theorems of Arts. 18 and 23, these forms are convertible one into the other. An Equation of Differences is said to be of the n^{th} order when the successive value, or the difference, of the highest order which it involves is the n^{th} . An Equation of Differences of any order is said moreover to be of the first, second, &c. degree, when the successive value, or the Difference, which marks its order, is raised at most to the first, second, &c. power; or, when it involves a product of successive values or differences at most of two, three, &c. dimensions, it is said to be of the second, third, &c. degree.

73. The complete integral of an equation of differences of the n^{th} order will contain n arbitrary constants.

Let $\dots u_x, u_{x+h}, u_{x+2h}, \dots$ be a series of terms corresponding to the successive values $x, x+h, x+2h, \dots$; and let

$$F(x, u_x, a) = 0$$

be the equation by which the general term is determined as a function of x and a , or the equation of the series, a being an arbitrary constant. Since this equation must hold for all the succeeding terms, we shall have

$$F(x+h, u_{x+h}, a) = 0.$$

Eliminating a between these two equations, we get an equation between x , u_x , and u_{x+h} ; or, substituting $u_x + \Delta u_x$ for u_{x+h} , an equation between x , u_x , and Δu_x , which is the equation of differences of the first order whose primitive equation is

$$F(x, u_x, a) = 0.$$

In like manner if the equation of the general term contained two arbitrary constants a and b , as

$$F(x, u_x, a, b) = 0,$$

we might eliminate a and b by means of the two succeeding equations,

$$F(x+h, u_{x+h}, a, b) = 0, \quad F(x+2h, u_{x+2h}, a, b) = 0,$$

and thus get an equation between x , u_x , u_{x+h} , u_{x+2h} ; or, substituting

$$u_x + \Delta u_x \text{ for } u_{x+h}, \text{ and } u_x + 2\Delta u_x + \Delta^2 u_x \text{ for } u_{x+2h},$$

an equation between x , u_x , Δu_x , $\Delta^2 u_x$, without the constants a and b , which is an equation of differences of the second order, having for its complete primitive the equation

$$F(x, u_x, a, b) = 0.$$

Hence it appears that every equation of differences of the first order, or between two successive terms of a series, will introduce one arbitrary constant into the equation of the series; every equation of differences of the second order, or between three successive terms, will introduce two arbitrary constants into the equation of the series; and, generally, every equation of differences of the n^{th} order will introduce n arbitrary constants into the equation of the series.

Linear Equation of Differences of the First Order.

74. The general equation of the first order and degree is

$$u_{x+1} - A_x u_x = B_x,$$

A_x and B_x being functions of x . To integrate it, assume

$$u_x = v_x w_x, \quad \therefore v_{x+1}(w_x + \Delta w_x) - A_x v_x w_x = B_x;$$

and in order that this equation may resolve itself into two others each of which admits of being integrated, assume (as we are at liberty to do, having made only one supposition respecting v_x and w_x)

$$v_{x+1} w_x - A_x v_x w_x = 0,$$

$$\text{or, dividing by } v_x w_x, \quad \frac{v_{x+1}}{v_x} = A_x.$$

$$\text{But } \Delta \log v_x = \log \frac{v_{x+1}}{v_x}, \quad \therefore \Delta \log v_x = \log A_x,$$

$$\therefore \log v_x = \Sigma \log A_x = \log PA_{x-1}, \quad (\text{Art. 50.})$$

or $v_x = PA_{x-1}$, the constant being unnecessary.

The other part of the equation gives $v_{x+1} \Delta w_x = B_x$,

$$\therefore \Delta w_x = \frac{B_x}{PA_x}, \quad \text{or } w_x = \Sigma \left(\frac{B_x}{PA_x} \right) + C,$$

$$\text{and } u_x = PA_{x-1} \left\{ \Sigma \left(\frac{B_x}{PA_x} \right) + C \right\}$$

the complete integral, involving one arbitrary constant.

Taking the difference of the result

$$\frac{u_x}{PA_{x-1}} = \Sigma \left(\frac{B_x}{PA_x} \right) + C, \quad \text{we get } \frac{u_{x+1} - A_x u_x}{PA_x} = \frac{B_x}{PA_x},$$

which shews that $\frac{1}{PA_x}$ is a factor which makes each side of the proposed equation integrable; and it is generally the most convenient way of integrating the equation to multiply it by this factor.

$$\text{Ex. 1. } u_{x+1} - au_x = x^2.$$

$$\text{Here } A_x = a, \quad PA_x = a^x;$$

$$\begin{aligned}\therefore \frac{u_x}{a^{x-1}} &= \Sigma \left(\frac{x^2}{a^x} \right) = \Sigma (x^2 a^x), \text{ putting } \frac{1}{a} = \alpha, \\ &= \frac{x^2 a^x}{\alpha - 1} - \frac{(2x+1) \alpha^{x+1}}{(\alpha - 1)^2} + \frac{2 \alpha^{x+2}}{(\alpha - 1)^3} + C; \text{ (Art. 59.)} \\ \therefore u_x &= \frac{x^2}{1 - \alpha} - \frac{2x+1}{(1 - \alpha)^2} + \frac{2}{(1 - \alpha)^3} + C \alpha^{x-1}.\end{aligned}$$

$$\text{Ex. 2. } u_{x+1} = 2 \cdot \frac{2x+1}{x+2} u_x.$$

$$u_x = C \cdot 2^x \frac{1 \cdot 3 \cdot 5 \dots (2x-1)}{1 \cdot 2 \cdot 3 \dots (x+1)}.$$

$$\text{Ex. 3. } u_{x+1} - a u_x = \cos x \theta.$$

$$u_x = \frac{\cos (x-1) \theta - a \cos x \theta}{a^2 - 2a \cos \theta + 1} + C a^x.$$

Ex. 4. Two vessels which hold a and b gallons respectively are filled, the one with proof spirit, the other with water; c gallons are taken from each and poured into the other; and this is repeated such a number of times as to make their contents of the same strength; find the number of times

$$x = \frac{\log \frac{1}{2} \left(1 - \frac{a}{b} \right)}{\log \left(1 - \frac{c}{a} - \frac{c}{b} \right)}.$$

Indirect Integrals of Equations of Differences.

75. Since the equation of differences of the first order,

$$f(x, u_x, \Delta u_x) = 0,$$

is formed by eliminating the constant a between the equations

$$u_x = F(x, a), \quad u_{x+h} = F(x+h, a),$$

it follows that we shall arrive at the same equation of differences, whether a be constant, or be a function of x such as a_x , provided it satisfies the condition

$$F(x+h, a_{x+h}) - F(x+h, a_x) = 0 \quad (1).$$

Now this equation is satisfied by $a_{x+h}=a_x$, which gives $\Delta a_x=0$, and $a_x=a$, a constant, and leads to the ordinary or direct equation to the series $u_x=F(x, a)$.

Also the first member of (1) will be divisible by $a_{x+h}-a_x$, because a_x is a value of a_{x+h} , which satisfies equation (1); and if dimensions of a_{x+h} and a_x superior to the first are involved in it, the result of this division will be an equation involving a_{x+h} and a_x ; i.e. an equation of differences of the first order with respect to a_x , the solution of which will give one or more values of a_x in terms of x and arbitrary constants; and these being substituted for a_x in the equation $u_x=F(x, a_x)$, will furnish equations of series, which are primitive equations of

$$f(x, u_x, \Delta u_x) = 0,$$

and each involves an arbitrary constant.

If equation (1) does not involve higher dimensions of a_x and a_{x+h} than the first, they will disappear from the result when it is divided by $a_{x+h}-a_x$. In this case a_x will have only one value, viz. $a_x=a$, and there will be only one equation of a series corresponding to the proposed equation of differences. The mode in which the indirect solutions just treated of are obtained, is analogous to that in which the singular solutions of differential equations are obtained; but whereas the latter can contain no arbitrary constant, indirect solutions of equations of differences may contain as many arbitrary constants as the complete integral itself from which they are deduced.

Ex. $u_x = x\Delta u_x + F(\Delta u_x)$.

Taking the Difference, we find

$$\Delta u_x = \Delta u_x + (x+1) \Delta^2 u_x + \Delta F(\Delta u_x),$$

$$\text{or } 0 = (x+1) \Delta^2 u_x + \Delta F(\Delta u_x),$$

which is evidently satisfied by $\Delta u_x = a$, a constant;

$$\therefore u_x = ax + a';$$

and substituting in the proposed equation

$$ax + a' = ax + F(a), \quad \therefore a' = F(a);$$

$$\therefore u_x = xa + F(a),$$

the complete integral, containing one arbitrary constant.

For the indirect solutions we shall have

$$u_x = xa_x + F(a_x),$$

a_x being determined from the equation

$$(x+1)a_{x+1} + F(a_{x+1}) - (x+1)a_x - F(a_x) = 0.$$

Suppose, for instance, that $F(a_x) = a_x^2$,

$$\therefore (x+1)(a_{x+1} - a_x) + a_{x+1}^2 - a_x^2 = 0,$$

or, rejecting the factor $a_{x+1} - a_x$,

$$a_{x+1} + a_x = -(x+1);$$

$$\therefore \frac{a_x}{(-1)^{x-1}} = -\sum (x+1)(-1)^x = \frac{2x+1}{4}(-1)^x + C,$$

$$\text{or } a_x = -\frac{2x+1}{4} + C(-1)^x;$$

$$\therefore u_x = xa_x + a_x^2 = \frac{1-4x^2}{16} - \frac{1}{2}C(-1)^x + C^2.$$

Linear Equations of Differences of all Orders.

The linear equation of Differences of the n^{th} order is

$$u_{x+n} + p_1 u_{x+n-1} + p_2 u_{x+n-2} + \dots + p_n u_x = X,$$

all the coefficients being functions of x ; the first step towards its integration is to establish the following theorem.

76. If there be n particular values v_x, w_x, \dots, z_x , which, when substituted for u_x , satisfy the equation

$$u_{x+n} + p_1 u_{x+n-1} + p_2 u_{x+n-2} + \dots + p_n u_x = 0,$$

that has no term independent of u_x , its complete integral is

$$u_x = a_1 v_x + a_2 w_x + \dots + a_n z_x,$$

$a_1, a_2, \dots a_n$ being arbitrary constants.

For let this value be substituted in the expression

$$u_{x+n} + p_1 u_{x+n-1} + p_2 u_{x+n-2} + \dots + p_n u_x,$$

and it becomes, (collecting the terms multiplied by the factors $a_1, a_2, \dots a_n$)

$$a_1 (v_{x+n} + p_1 v_{x+n-1} + \dots + p_n v_x) + a_2 (w_{x+n} + p_1 w_{x+n-1} + \dots + p_n w_x) + \dots + a_n (z_{x+n} + p_1 z_{x+n-1} + \dots + p_n z_x).$$

Now since $v_x, w_x, \dots z_x$, satisfy the proposed equation, each of the quantities included within brackets is equal to zero, therefore the whole is identically zero; consequently the assumed value of u_x satisfies the proposed equation, and it contains n arbitrary constants, therefore it is the complete integral of that equation.

77. To integrate the equation of differences,

$$u_{x+n} + p_1 u_{x+n-1} + p_2 u_{x+n-2} + \dots + p_n u_x = q,$$

all the coefficients and q being constants.

Assume $u_x = v_x + k$; then by substitution we get

$$v_{x+n} + p_1 v_{x+n-1} + \dots + p_n v_x + k(1 + p_1 + \dots + p_n) - q = 0.$$

Let $k = \frac{q}{1 + p_1 + p_2 + \dots + p_n}$, then the equation becomes

$$v_{x+n} + p_1 v_{x+n-1} + p_2 v_{x+n-2} + \dots + p_n v_x = 0 \dots\dots\dots (1).$$

Let $v_x = a^x$, then $a^x (a^n + p_1 a^{n-1} + p_2 a^{n-2} + \dots + p_n)$ is the value of the first member; now this will vanish if a be any root of the equation (called the auxiliary equation),

$$f(a) = a^n + p_1 a^{n-1} + p_2 a^{n-2} + \dots + p_{n-1} a + p_n = 0.$$

Hence the n roots of this equation $a_1, a_2, a_3, \dots a_n$ will give n particular values of $v_x, a_1^x, a_2^x, a_3^x \dots a_n^x$ which satisfy equation (1); therefore its complete integral is

$$v_x = c_1 a_1^x + c_2 a_2^x + \dots + c_n a_n^x;$$

and the complete integral of the proposed equation is

$$u_x = c_1 a_1^x + c_2 a_2^x + \dots + c_n a_n^x + \frac{q}{f(1)}.$$

This shews that we may treat the proposed equation as if it had no term q independent of u_x , provided we divide that term by $f(1)$ the sum of the coefficients, and add the quotient to the value of u_x obtained on the supposition that $q = 0$.

If we replace c_1 by $c_0 - q \div f(1)$, and a_1 by $1 + h$ where h is small, so that very nearly $a_1^x = 1 + hx$, and $f(1) = -hf'(1)$ since $1 + h$ is a root of $f(a) = 0$, then reducing and making $h = 0$, we get the form which the solution takes when $f(1) = 0$, viz.

$$u_x = c_0 + \frac{qx}{f'(1)} + \&c.$$

78. If the auxiliary equation have equal roots, the above ceases to be the form of the complete solution; because, in that case, it does not involve the due number of arbitrary constants; and it must be modified as follows. Suppose two roots a_1, a_2 , to be very nearly equal to one another, so that $a_2 = a_1 + h$, h being a very small known quantity; then

$$\begin{aligned} c_1 a_1^x + c_2 a_2^x &= (c_1 + c_2) a_1^x + c_2 \{ x a_1^{x-1} h + \frac{x(x-1)}{1 \cdot 2} a_1^{x-2} h^2 + \&c. \} \\ &= C_1 a_1^x + C_2 \{ x a_1^{x-1} + \frac{x(x-1)}{1 \cdot 2} a_1^{x-2} h + \&c. \}, \end{aligned}$$

replacing the constants $c_1 + c_2$ by C_1 , and $\frac{h c_2}{a_1}$ by C_2 ;

now this equation continues true however small h be taken, and therefore when $h = 0$; in which case the second member becomes $(C_1 + C_2 x) a_1^x$;

$$\therefore u_x = (C_1 + C_2 x) a_1^x + c_3 a_3^x + \&c.$$

Similarly, if the auxiliary equation have r roots equal to a_1 , the complete solution will be

$$u_x = (c_0 + c_1x + c_2x^2 + \dots + c_{r-1}x^{r-1}) a_1^x + c_{r+1}a_1^{x+r+1} + \&c ;$$

of the correctness of which, we may be assured by the following reverse process. Assume $u_x = a^x w_x$, then

$$u_{x+n} = a^{x+n} \cdot w_{x+n} = a^x \cdot a^n (1 + \Delta)^n w_x = a^x (a + a\Delta)^n w_x,$$

and the first side of the proposed equation becomes, when divided by a^x ,

$$\begin{aligned} &= \{(a + a\Delta)^n + p_1 (a + a\Delta)^{n-1} + \&c. + p_n\} w_x \\ &= f(a + a\Delta) w_x = \{f(a) + f'(a) \cdot a\Delta + f''(a) \frac{a^2 \Delta^2}{1 \cdot 2} + \&c.\} w_x \\ &= f(a) w_x + \frac{a}{1} \cdot f'(a) \cdot \Delta w_x + \frac{a^2}{1 \cdot 2} f''(a) \cdot \Delta^2 w_x + \&c. + a^n \Delta^n w_x. \end{aligned}$$

Now suppose $a = a_1$, and $f(a) = 0$ to have r roots equal to a_1 ; this makes the terms as far as $f^{(r-1)}(a)$ vanish; and if

$$w_x = c_0 + c_1x + \dots + c_{r-1}x^{r-1}, \text{ then } \Delta^r w_x = 0, \Delta^{r+1} w_x = 0, \&c.,$$

and all the remaining terms vanish; and consequently the equation is satisfied by

$$u_x = (c_0 + c_1x + \dots + c_{r-1}x^{r-1}) a_1^x.$$

Hence we see that every root a_1 that occurs r times in the auxiliary equation, gives rise, in the complete integral, to a term of the form $f_{r-1}(x)a_1^x$, where $f_{r-1}(x)$ denotes a rational integral function of x of the $(r-1)^{\text{th}}$ degree involving r arbitrary constants. The root a_1 may be either real or imaginary.

79. Also if the auxiliary equation have a pair of imaginary roots,

$$m \pm n \sqrt{-1} = \rho (\cos \theta \pm \sqrt{-1} \sin \theta),$$

$$\text{putting } \rho = \sqrt{m^2 + n^2}, \tan \theta = \frac{n}{m},$$

the corresponding terms in the value of u_x will be

$$\begin{aligned} &C\rho^x (\cos \theta + \sqrt{-1} \sin \theta)^x + C'\rho^x (\cos \theta - \sqrt{-1} \sin \theta)^x \\ &= \rho^x (c_1 \cos x\theta + c_2 \sin x\theta), \end{aligned}$$

changing the arbitrary constants.

And if the same pair of imaginary roots occur r times in the auxiliary equation, the corresponding terms in the value of u_x will be

$$(a_0 + a_1 x + \dots + a_{r-1} x^{r-1}) \rho^x (\cos \theta + \sqrt{-1} \sin \theta)^x \\ + (b_0 + b_1 x + \dots + b_{r-1} x^{r-1}) \rho^x (\cos \theta - \sqrt{-1} \sin \theta)^x ;$$

or changing the arbitrary constants,

$$(c_0 + c_1 x + \dots + c_{r-1} x^{r-1}) \rho^x \cos x\theta + (c'_0 + c'_1 x + \dots + c'_{r-1} x^{r-1}) \rho^x \sin x\theta.$$

Ex. 1. $u_{x+4} - 2u_{x+3} - 13u_{x+2} + 14u_{x+1} + 24u_x = 0.$

The auxiliary equation is $(a+1)(a-2)(a+3)(a-4) = 0$;

$$\therefore u_x = c_1 (-1)^x + c_2 2^x + c_3 (-3)^x + c_4 4^x.$$

Ex. 2. $u_{x+3} - 5u_{x+2} + 8u_{x+1} - 4u_x = 0.$

The auxiliary equation is $(a-1)(a-2)^2 = 0$;

$$\therefore u_x = c_0 + (c + c'x) 2^x;$$

and if the second member be q ,

$$u_x = c_0 + qx + (c + c'x) 2^x.$$

Ex. 3. $u_{x+3} - u_x = q.$

The auxiliary equation is $a^3 - 1 = 0$, whose roots are 1, and $\cos \frac{2\pi}{3} \pm \sqrt{-1} \sin \frac{2\pi}{3}$;

$$\therefore u_x = c_0 + c \cos \frac{2\pi x}{3} + c' \sin \frac{2\pi x}{3} + \frac{1}{3} qx.$$

80.* The Differential Calculus being a particular case of that of Finite Differences, a strict analogy exists between the methods and results in the two Subjects, as the reader cannot fail to have observed; indeed whenever in the latter a result is obtained with an indeterminate increment for the principal variable, it is possible to pass to the corresponding result in the former, by a method similar to that pursued in Art. 25; and which we shall further illustrate by the following instance.

In the equation $u_{x+2h} - 2mu_{x+h} + (m^2 + n^2) u_x = 0$,

putting $u_x = a^x$, we find $a = (m \pm n\sqrt{-1})^{\frac{1}{2}}$;

$$\begin{aligned} \therefore u_x &= C_1 (m + n\sqrt{-1})^{\frac{x}{2}} + C_2 (m - n\sqrt{-1})^{\frac{x}{2}} \\ &= (m^2 + n^2)^{\frac{x}{2k}} \left\{ c_1 \cos \left(\frac{x}{h} \tan^{-1} \frac{n}{m} \right) + c_2 \sin \left(\frac{x}{h} \tan^{-1} \frac{n}{m} \right) \right\}. \end{aligned}$$

Now the proposed equation is

$$\Delta^2 u_x - 2(m-1)\Delta u_x + \{(m-1)^2 + n^2\} u_x = 0;$$

or, if we replace the known quantities $m-1$ and n by other known quantities $m_1 h$ and $n_1 h$, the equation and its solution take the forms

$$\begin{aligned} \frac{\Delta^2 u_x}{h^2} - 2m_1 \frac{\Delta u_x}{h} + (m_1^2 + n_1^2) u_x &= 0, \\ u_x &= \{(1 + m_1 h)^2 + n_1^2 h^2\}^{\frac{x}{2h}} \\ &\times \left\{ c_1 \cos \left(\frac{x}{h} \tan^{-1} \frac{n_1 h}{1 + m_1 h} \right) + c_2 \sin \left(\frac{x}{h} \tan^{-1} \frac{n_1 h}{1 + m_1 h} \right) \right\}. \end{aligned}$$

Now take the limits of these expressions when $h=0$, and we find the well-known results, since $(1 + 2m_1 h)^{\frac{1}{2h}}$ becomes e^{m_1} ,

$$\begin{aligned} \frac{d^2 u_x}{dx^2} - 2m_1 \frac{du_x}{dx} + (m_1^2 + n_1^2) u_x &= 0, \\ u_x &= e^{m_1 x} (c_1 \cos n_1 x + c_2 \sin n_1 x). \end{aligned}$$

80. Before proceeding to the general case of linear equations, we shall consider the case of a linear equation with all its coefficients constant, but having a function of x , X , for the term independent of u_x .

This equation can be shortly treated by the method of separation of symbols. For if we substitute $D^n u_x$ for u_{x+n} , it becomes

$$(D^n + p_1 D^{n-1} + p_2 D^{n-2} + \dots + p_n) u_x = X;$$

or, if $a_1, a_2, \dots a_n$ be the roots of the auxiliary equation,

$$\begin{aligned} & (D - a_1)(D - a_2) \dots (D - a_n) u_x = X; \\ \therefore u_x &= \frac{1}{(D - a_1)(D - a_2) \dots (D - a_n)} X \dots \dots \dots (1). \end{aligned}$$

Now resolve this function of D into partial fractions (see Integral Calculus, Art. 34), so that the values of $A_1, A_2, \&c.$, are all known in terms of $a_1, a_2, \&c.$ respectively; then

$$u_x = \left(\frac{A_1}{D - a_1} + \frac{A_2}{D - a_2} + \dots + \frac{A_n}{D - a_n} \right) X.$$

But we have seen Art. 58, that $(D - a)^{-1} X = a^{-1} \Sigma (X a^{-x}) + C a^{-x}$.

$$\begin{aligned} \therefore u_x &= A_1 a_1^{-1} \Sigma (X a_1^{-x}) + A_2 a_2^{-1} \Sigma (X a_2^{-x}) + \dots + A_n a_n^{-1} \Sigma (X a_n^{-x}) \\ &\quad + c_1 a_1^{-x} + c_2 a_2^{-x} + \dots + c_n a_n^{-x}; \end{aligned}$$

an arbitrary constant being introduced by each of the integrations, and the product of two or more constants being replaced by a single constant. The part of the above expression involving the arbitrary constants is called the complementary function; and it is the value of u_x which satisfies the proposed equation when $X = 0$.

81. If the auxiliary equation have r roots equal to a_1 , then from (1)

$$u_x = \frac{1}{(D - a_1)^r (D - a_{r+1}) \dots (D - a_n)} X;$$

and this function of D may be resolved into partial fractions, (see Integral Calculus, Art. 36) so that

$$\begin{aligned} u_x &= \left\{ \frac{A_1}{D - a_1} + \frac{A_2}{(D - a_1)^2} + \dots + \frac{A_r}{(D - a_1)^r} \right. \\ &\quad \left. + \frac{A_{r+1}}{D - a_{r+1}} + \dots + \frac{A_n}{D - a_n} \right\} X. \end{aligned}$$

But $(D - a)^{-n} X = a^{-n} \Sigma^n (a^{-x} X) + a^{-n} (C_0 + C_1 x + \dots + C_{n-1} x^{n-1})$, $C_0, C_1, \&c.$ being the constants introduced after each integration;

$$\begin{aligned} \therefore u_x &= A_1 a_1^{-1} \Sigma (a_1^{-x} X) + A_2 a_1^{-2} \Sigma^2 (a_1^{-x} X) + \dots + A_r a_1^{-r} \Sigma^r (a_1^{-x} X) \\ &\quad + A_{r+1} a_{r+1}^{-1} \Sigma (a_{r+1}^{-x} X) + \dots + A_n a_n^{-1} \Sigma (a_n^{-x} X) \\ &\quad + (c_0 + c_1 x + \dots + c_{r-1} x^{r-1}) a_1^{-x} + c_{r+1} a_{r+1}^{-x} + \dots + c_n a_n^{-x}, \end{aligned}$$

a single constant, as before, being substituted for the sum or product of several constants in forming the complementary function.

82. Again, suppose $\alpha_1 = \rho (\cos \theta + \sqrt{-1} \sin \theta)$ to be an imaginary root of the auxiliary equation; then since $\frac{A_1}{a_1}$ is a function of a_1 , it will be of the form $R (\cos \alpha + \sqrt{-1} \sin \alpha)$; and the term involving α_1 in the value of u_x will consequently be

$$R\rho^x \{ \cos (x\theta + \alpha) + \sqrt{-1} \sin (x\theta + \alpha) \} \Sigma \rho^{-x} X (\cos x\theta - \sqrt{-1} \sin x\theta) \dots (1).$$

Now the term in u_x involving the conjugate root to α_1 , will result from this, by changing the sign of $\sqrt{-1}$; and therefore the sum of the two terms introduced into the value of u_x by the pair of imaginary roots $\rho (\cos \theta \pm \sqrt{-1} \sin \theta)$ will equal twice the real part of expression (1); that is,

$$2R\rho^x \cos (x\theta + \alpha) (\Sigma \rho^{-x} X \cos x\theta + C) \\ + 2R\rho^x \sin (x\theta + \alpha) (\Sigma \rho^{-x} X \sin x\theta + C');$$

where

$$2R\rho^x C \cos (x\theta + \alpha), \quad 2R\rho^x C' \sin (x\theta + \alpha)$$

are the terms introduced into the complementary function; and, if we alter the constants, may be replaced by

$$\rho^x (c \cos x\theta + c' \sin x\theta).$$

83. Exactly in the same way, if the imaginary root

$$\alpha_1 = \rho (\cos \theta + \sqrt{-1} \sin \theta),$$

occur r times in the auxiliary equation it will produce in the value of u_x , r terms of the form $A_m a_1^{x-m} \Sigma^m (a_1^{-x} X)$; or, since $\frac{A_m}{a_1^m}$ is a function of a_1 and may be assumed

$$= R_m (\cos \alpha_m + \sqrt{-1} \sin \alpha_m),$$

of the form

$$R_m \rho^x \{ \cos (x\theta + \alpha_m) \\ + \sqrt{-1} \sin (x\theta + \alpha_m) \} \Sigma^m \rho^{-x} (\cos x\theta - \sqrt{-1} \sin x\theta) X \dots (2).$$

But the root conjugate to α_1 will produce a term exactly the

same as this, except with $-\sqrt{-1}$ instead of $+\sqrt{-1}$; consequently the sum of these terms will equal twice the real part of (2); that is,

$$2R_m \rho^x \{ \cos(x\theta + \alpha_m) \Sigma^m (\rho^{-x} \cos x\theta X) + \sin(x\theta + \alpha_m) \Sigma^m (\rho^{-x} \sin x\theta X) \},$$

and to get all the terms introduced by the pair of imaginary roots $\rho (\cos \theta \pm \sqrt{-1} \sin \theta)$ that occur r times in the auxiliary equation, m in the above formula must receive all values from 1 to r . Also we see that the part of the complementary function introduced by these roots, by substituting a single constant for the sum or product of other constants, will take the form

$$(c_0 + c_1 x + \dots + c_{r-1} x^{r-1}) \rho^x \cos x\theta + (c'_0 + c'_1 x + \dots + c'_{r-1} x^{r-1}) \rho^x \sin x\theta.$$

We shall now give an instance of each of the cases that have been examined. It may be observed, that for equations capable of being reduced to either of the forms

$$f(D) u_x = a^x, \quad \text{or} \quad f(D) u_x = p_0 x^m + p_1 x^{m-1} + \&c. + p_m,$$

the process may be greatly simplified. For in the former case, since $D a^x = a \cdot a^x$, the symbol D is equivalent to the factor a ,

and may have a substituted for it in the formula $u_x = \frac{1}{f(D)} a^x$; so

that the solution of the equation $f(D) u_x = a^x$ is $u_x = \frac{a^x}{f(a)}$ + com-

plementary function. And in the latter case, if $\frac{1}{f(D)} = \frac{1}{f(1+\Delta)}$ can be expanded in the form $A_0 + A_1 \Delta + A_2 \Delta^2 + \&c.$, then

$$u_x = (A_0 + A_1 \Delta + A_2 \Delta^2 + \dots + A_m \Delta^m) (p_0 x^m + p_1 x^{m-1} + \dots + p_m)$$

+ complementary function.

$$\text{Ex. 1. } u_{x+3} - 9u_{x+2} + 26u_{x+1} - 24u_x = 5^x,$$

$$\text{or } (D^3 - 9D^2 + 26D - 24) u_x = f(D) u_x = 5^x;$$

$$\therefore u_x = \frac{5^x}{(D-2)(D-3)(D-4)} = \left(\frac{1}{2} \frac{1}{D-2} - \frac{1}{D-3} + \frac{1}{2} \frac{1}{D-4} \right) 5^x$$

$$= \frac{1}{2} \cdot 2^{x-1} \Sigma \left(\frac{5}{2} \right)^x - 3^{x-1} \Sigma \left(\frac{5}{3} \right)^x + \frac{1}{2} \cdot 4^{x-1} \Sigma \left(\frac{5}{4} \right)^x$$

$$\begin{aligned}
&= 2^{x-2} \frac{2}{3} \left(\frac{5}{2}\right)^x - 3^{x-1} \frac{3}{2} \left(\frac{5}{3}\right)^x + \frac{1}{2} \cdot 4^{x-1} \frac{4}{3} \left(\frac{5}{4}\right)^x + \&c. \\
&= \frac{5^x}{6} + c_0 2^x + c_1 3^x + c_2 4^x,
\end{aligned}$$

a result we could have foreseen because $f(5) = 3 \cdot 2 \cdot 1 = 6$.

$$\text{Ex. 2. } a^2 u_{x+2} - 2a u_{x+1} + u_x = a^2 x,$$

$$\text{or } (aD - 1)^2 u_x = a^2 x;$$

$$\begin{aligned}
\therefore u_x &= \left(D - \frac{1}{a}\right)^{-2} x = \frac{1}{a^{x-2}} \Sigma^2 (x a^x) = \frac{1}{a^{x-2}} (x \Sigma^2 a^x - 2 \Sigma^3 a^{x+1}) \\
&= \frac{1}{a^{x-2}} \left\{ \frac{x a^x}{(a-1)^2} - \frac{2 a^{x+1}}{(a-1)^3} + Cx + C' \right\} \\
&= \frac{a^2 x}{(a-1)^2} - \frac{2 a^3}{(a-1)^3} + (c + c'x) \frac{1}{a^x}.
\end{aligned}$$

Otherwise,

$$\begin{aligned}
u_x &= \left(1 - \frac{1}{a} + \Delta\right)^{-2} x = \left\{ \left(1 - \frac{1}{a}\right)^{-2} - 2 \left(1 - \frac{1}{a}\right)^{-3} \Delta + \&c. \right\} x \\
&= \frac{a^2 x}{(a-1)^2} - \frac{2 a^3}{(a-1)^3} + (c + c'x) a^{-x}.
\end{aligned}$$

$$\text{Ex. 3. } u_{x+2} - 2a \cos \theta u_{x+1} + a^2 u_x = X,$$

$$\text{or } (D^2 - 2a \cos \theta D + a^2) u_x = X;$$

$$\therefore u_x = \frac{X}{(D - az)(D - az^{-1})} = \frac{1}{a(z - z^{-1})} \left(\frac{X}{D - az} - \frac{X}{D - az^{-1}} \right),$$

putting $2 \cos \theta = z + z^{-1}$, and consequently $2 \sqrt{-1} \sin \theta = z - z^{-1}$;

$$\therefore u_x = \frac{1}{a(z - z^{-1})} \{ (az)^{x-1} \Sigma X (az)^{-x} - (az^{-1})^{x-1} \Sigma X (a^{-1}z)^x \}.$$

Now the value of the former of these integrals is

$$\begin{aligned}
&\frac{a^{x-2}}{2 \sqrt{-1} \sin \theta} \{ \cos (x-1) \theta \\
&+ \sqrt{-1} \sin (x-1) \theta \} \Sigma X a^{-x} (\cos x \theta - \sqrt{-1} \sin x \theta);
\end{aligned}$$

hence, taking twice the possible part of this expression, we have

$$u_x = \frac{a^{x-2}}{\sin \theta} \{ \sin (x-1) \theta \Sigma (X a^{-x} \cos x \theta) - \cos (x-1) \theta \Sigma (X a^{-x} \sin x \theta) \},$$

the complementary function being

$$a^x (c \cos x \theta + c' \sin x \theta).$$

If we suppose $X = a^x$, then as we could have foreseen

$$u_x = \frac{a^{x-2}}{2(1 - \cos \theta)} + a^x (c \cos x \theta + c' \sin x \theta).$$

The case of $\theta = 0$ will be presently noticed.

Ex. 4. $u_{x+4} + p_1 u_{x+3} + p_2 u_{x+2} + p_3 u_{x+1} + p_4 u_x = X,$

where the auxiliary equation is supposed to have two pairs of imaginary roots of the form $a(\cos \theta \pm \sqrt{-1} \sin \theta)$, and consequently the proposed equation is of the form

$$(D^2 - 2a \cos \theta D + a^2)^2 u_x = (D - az)^2 (D - az^{-1})^2 u_x = X.$$

Now by the preceding example we have

$$\frac{2a \sqrt{-1} \sin \theta}{(D - az)(D - az^{-1})} = \frac{1}{D - az} - \frac{1}{D - az^{-1}};$$

$$\therefore \frac{-4a^2 \sin^2 \theta}{(D - az)^2 (D - az^{-1})^2} = \frac{1}{(D - az)^2}$$

$$+ \frac{\sqrt{-1}}{a \sin \theta} \left(\frac{1}{D - az} - \frac{1}{D - az^{-1}} \right) + \frac{1}{(D - az^{-1})^2};$$

$$\therefore -4a^2 \sin^2 \theta u_x = (az)^{x-2} \Sigma^2 (az)^{-x} X$$

$$+ \frac{\sqrt{-1}}{a \sin \theta} (az)^{x-1} \Sigma (az)^{-x} X + \text{terms in } az^{-1}$$

$$= a^{x-2} \{ \cos (x-2) \theta + \sqrt{-1} \sin (x-2) \theta \}$$

$$\times \Sigma^2 a^{-x} X (\cos x \theta - \sqrt{-1} \sin x \theta)$$

$$+ \frac{a^{x-2} \sqrt{-1}}{\sin \theta} \{ \cos (x-1) \theta + \sqrt{-1} \sin (x-1) \theta \}$$

$$\times \Sigma a^{-x} X (\cos x \theta - \sqrt{-1} \sin x \theta) + \&c.,$$

and taking in these terms only the possible part, and doubling it, we have $-2 \sin^3 \theta a^{-x+4} u_x$

$$\begin{aligned} &= \sin \theta \cos (x-2) \theta \Sigma^2 (a^{-x} X \cos x\theta) \\ &+ \sin \theta \sin (x-2) \theta \Sigma^2 (a^{-x} X \sin x\theta) \\ &+ \cos (x-1) \theta \Sigma (a^{-x} X \sin x\theta) - \sin (x-1) \theta \Sigma (a^{-x} X \cos x\theta), \end{aligned}$$

the complementary function being

$$(b + b'x) a^x \sin x\theta + (c + c'x) a^x \cos x\theta.$$

If $X = a^x$, the result will be, as we are aware,

$$u_x = \frac{a^{x-4}}{4(1 - \cos \theta)^2} + \text{complementary function.}$$

84. To integrate the linear equation of differences of the n^{th} order, the coefficients being functions of x ,

$$u_{x+n} + p_1 u_{x+n-1} + p_2 u_{x+n-2} + \dots + p_n u_x = X,$$

on the supposition that it can be solved when $X = 0$.

If $z_x, {}^1z_x, \dots {}^{n-1}z_x$ be n particular values of v_x in the equation

$$v_{x+n} + p_1 v_{x+n-1} + p_2 v_{x+n-2} + \dots + p_n v_x = 0 \dots \dots \dots (1),$$

with which the proposed coincides when its second member is zero, we have (Art. 77)

$$v_x = c_0 z_x + c_1 {}^1z_x + c_2 {}^2z_x + \dots + c_{n-1} {}^{n-1}z_x.$$

If we now divide both sides by z_x and take the difference, we shall eliminate c_0 ; next dividing both sides by the coefficient of c_1 in the new result, which suppose y_x , and taking the difference, we shall eliminate c_1 ; again dividing by w_x the coefficient of c_2 and taking the difference, we shall eliminate c_2 ; and proceeding in this manner till all the constants are eliminated, our final result will be of the form (each Δ affecting the whole of the expression that follows it)

$$\Delta \frac{1}{t_x} \Delta \frac{1}{w_x} \dots \Delta \frac{1}{y_x} \Delta \left(\frac{v_x}{z_x} \right),$$

in which expression the coefficient of v_{x+n} is evidently

$$\frac{1}{t_{x+1} w_{x+2} \dots y_{x+n-1} z_{x+n}};$$

therefore, dividing by this coefficient, we get the expression

$$t_{x+1} w_{x+2} \dots y_{x+n-1} z_{x+n} \Delta \frac{1}{t_x} \Delta \frac{1}{w_x} \dots \Delta \frac{1}{y_x} \Delta \left(\frac{v_x}{z_x} \right),$$

which must be equivalent to the first member of equation (1). Therefore the same expression, only with u_x instead of v_x , must be equivalent to the first member of the proposed equation, and consequently equal to X ; hence, equating these equals, and integrating, we get

$$u_x = z_x \Sigma y_x \Sigma w_x \dots \Sigma t_x \Sigma \frac{X}{t_{x+1} w_{x+2} \dots y_{x+n-1} z_{x+n}},$$

(each Σ affecting the whole of the expression which follows it) which is a general formula for the integration of any linear equation of differences whose solution can be effected when $X=0$.

In the case of constant coefficients if the auxiliary equation contain equal or imaginary roots, this method is still applicable; it is only necessary to assume for v_x the value belonging to the case of equal or imaginary roots, as will be seen in some of the following instances.

Ex. 1. Suppose the coefficients of the equation to be constant, and $a_1, a_2, \dots a_n$ to be the n roots of its auxiliary equation.

Then $v_x = c_1 a_1^x + c_2 a_2^x + c_3 a_3^x + \dots + c_n a_n^x$,

$$\therefore \Delta \frac{v_x}{a_1^x} = c_2 \left(\frac{a_2}{a_1} \right)^x + c_3 \left(\frac{a_3}{a_1} \right)^x + \dots + c_n \left(\frac{a_n}{a_1} \right)^x,$$

changing, both in this, and in the similar succeeding steps, the arbitrary constants;

$$\Delta \left(\frac{a_1}{a_2} \right)^x \Delta \frac{v_x}{a_1^x} = c_2 \left(\frac{a_3}{a_2} \right)^x + \dots + c_n \left(\frac{a_n}{a_2} \right)^x$$

$$\Delta \left(\frac{a_2}{a_3} \right)^x \Delta \left(\frac{a_1}{a_2} \right)^x \Delta \frac{v_x}{a_1^x} = c_4 \left(\frac{a_4}{a_3} \right)^x + \dots + c_n \left(\frac{a_n}{a_3} \right)^x$$

.....

$$\Delta \left(\frac{a_{n-1}}{a_n} \right)^x \Delta \left(\frac{a_{n-2}}{a_{n-1}} \right)^x \dots \Delta \left(\frac{a_1}{a_2} \right)^x \Delta \frac{v_x}{a_1^x} = 0,$$

in which expression the coefficient of v_{x+n} is evidently

$$\frac{1}{a_n^x} \frac{1}{a_n a_{n-1} \dots a_2 a_1} = \frac{1}{(-1)^n p_n a_n^x},$$

therefore, dividing by this coefficient, and replacing v_x by u_x , we get

$$(-1)^n p_n a_n^x \Delta \left(\frac{a_{n-1}}{a_n} \right)^x \Delta \left(\frac{a_{n-2}}{a_{n-1}} \right)^x \dots \Delta \left(\frac{a_1}{a_2} \right)^x \Delta \frac{u_x}{a_1^x} = X,$$

$$\therefore u_x = \frac{(-1)^n}{p_n} a_1^x \Sigma \left(\frac{a_2}{a_1} \right)^x \dots \Sigma \left(\frac{a_n}{a_{n-1}} \right)^x \Sigma \left(\frac{X}{a_n^x} \right);$$

or, if we choose to introduce the arbitrary constant after each integration,

$$u_x = c_1 a_1^x + c_2 a_2^x + \dots c_n a_n^x, \\ + (-1)^n \frac{1}{p_n} a_1^x \Sigma \left(\frac{a_2}{a_1} \right)^x \Sigma \left(\frac{a_3}{a_2} \right)^x \dots \Sigma \left(\frac{a_n}{a_{n-1}} \right)^x \Sigma \left(\frac{X}{a_n^x} \right).$$

This result may be readily obtained by separation of symbols, and so shewn to agree with that obtained in Art. 80. For the proposed equation is

$$D^n u_x + p_1 D^{n-1} u_x + \dots + p_{n-1} D u_x + p_n u_x = X,$$

$$\text{or } (D - a_n) (D - a_{n-1}) \dots (D - a_1) u_x = X;$$

$$\therefore (D - a_{n-1}) (D - a_{n-2}) \dots (D - a_1) u_x = (D - a_n)^{-1} X = a_n^{-x-1} \Sigma (a_n^{-x} X)$$

$$= a_n^x X_n \text{ suppose, where } X_n = \frac{1}{a_n} \Sigma (a_n^{-x} X);$$

$$\therefore (D - a_{n-2}) (D - a_{n-3}) \dots (D - a_1) u_x = (D - a_{n-1})^{-1} a_n^x X_n$$

$$= a_{n-1}^{-x-1} \Sigma \left(\frac{a_n}{a_{n-1}} \right)^x X_n = a_{n-1}^x X_{n-1}, \text{ where } X_{n-1} = \frac{1}{a_{n-1}} \Sigma \left(\frac{a_n}{a_{n-1}} \right)^x X_n,$$

and so on till we arrive at

$$u_x = (D - a_1)^{-1} a^x X_2 = a_1^{x-1} \Sigma \left(\frac{a_2}{a_1} \right)^x X_2, \text{ when } X_2 = \frac{1}{a_2} \Sigma \left(\frac{a_3}{a_2} \right)^x X_3.$$

Hence restoring the values of $X_2, X_3 \dots X_n$, we get

$$\text{since } a_1 a_2 \dots a_n = (-1)^n p_n,$$

$$u_x = (-1)^n \frac{1}{p_n} a^x \Sigma \left(\frac{a_2}{a_1} \right)^x \Sigma \left(\frac{a_3}{a_2} \right)^x \dots \Sigma \left(\frac{a_n}{a_{n-1}} \right)^x \Sigma \left(\frac{X}{a^x} \right),$$

where each Σ affects the whole of the expression that follows it.

$$\text{Ex. 2.} \quad a^2 u_{x+2} - 2a u_{x+1} + u_x = a^2 X.$$

$$\text{Here } v_x = (c + c^1 x) \frac{1}{a_x}; \quad \therefore \Delta^2 (a^x v_x) = 0,$$

in which expression, the coefficient of v_{x+2} is a^{x+2} ; therefore, dividing by this quantity, and replacing v_x by u_x , we get

$$\frac{1}{a^{x+2}} \Delta^2 (a^x u_x) = X; \quad \therefore u_x = \frac{1}{a^x} \Sigma^2 (X a^{x+2}).$$

$$\text{Ex. 3.} \quad u_{x+2} - 2m u_{x+1} + (m^2 + n^2) u_x = X.$$

$$\text{Here } v_x = c_1 \rho^x \sin x\theta + c_2 \rho^x \cos x\theta,$$

$$\text{where } \rho^2 = m^2 + n^2, \quad \tan \theta = \frac{n}{m};$$

$$\therefore \frac{v_x}{\rho^x \cos x\theta} = c_1 \tan x\theta + c_2;$$

$$\therefore \Delta \left(\frac{v_x}{\rho^x \cos x\theta} \right) = \frac{c_1 \sin \theta}{\cos x\theta \cos (x+1)\theta};$$

$$\therefore \Delta \cos x\theta \cos (x+1)\theta \Delta \left(\frac{v_x}{\rho^x \cos x\theta} \right) = 0,$$

in which expression, the coefficient of v_{x+2} is $\frac{\cos (x+1)\theta}{\rho^{x+2}}$; therefore, dividing by this coefficient, and replacing v_x by u_x , we get

$$\frac{\rho^{x+2}}{\cos (x+1)\theta} \Delta \cos x\theta \cos (x+1)\theta \Delta \left(\frac{u_x}{\rho^x \cos x\theta} \right) = X;$$

$$\therefore u_x = \rho^x \cos x\theta \Sigma \sec x\theta \sec (x+1) \theta \Sigma \left\{ \frac{X \cos (x+1) \theta}{\rho^{x+2}} \right\};$$

to which we may add the terms $c_1 \rho^x \sin x\theta + c_2 \rho^x \cos x\theta$, if we suppose a constant to be added after each integration.

84*. Having given a particular integral of the equation

$$u_{x+2} + A_x u_{x+1} + B_x u_x = 0,$$

to find its complete solution.

Let v_x be a particular integral, then

$$v_{x+2} + A_x v_{x+1} + B_x v_x = 0.$$

Hence, eliminating A_x , we find, putting $\frac{u_x}{v_x} = z_x$,

$$\frac{\Delta z_{x+1}}{\Delta z_x} = \frac{B_x v_x}{v_{x+2}} = w_x, \text{ suppose;}$$

$$\text{or } \Delta \log (\Delta z_x) = \log w_x,$$

$$\therefore \Delta z_x = CPw_{x-1}, \quad (\text{Art. 50})$$

$$\therefore u_x = Cv_x \Sigma (Pw_{x-1}).$$

85. If we know a particular integral of a linear equation of any order that has no term independent of u_x , we may reduce it to another equation of the same kind of the order immediately inferior.

Let $u_x = v_x$ be a particular integral of a linear equation of differences of the n^{th} order reduced to the form

$$\Delta^n u_x + q_1 \Delta^{n-1} u_x + q_2 \Delta^{n-2} u_x + \dots + q_n u_x = f(\Delta) u_x = 0 \dots (1).$$

Assume $u_x = v_x \Sigma w_x$; then

$$\Delta^n u_x = (\Delta + D\Delta')^n v_x \Sigma w_x,$$

where Δ and D affect v_x only, and Δ' affects Σw_x only (Art. 34), and the proposed equation becomes

$$f(\Delta + D\Delta') v_x \Sigma w_x = 0.$$

Now $f(\Delta + D\Delta')$ is a rational integral function of Δ and Δ' , which we wish to arrange according to powers of Δ' , and this may be done at once by Taylor's theorem, which gives

$$f(\Delta) + f_1(\Delta) D\Delta' + \frac{1}{1 \cdot 2} f_2(\Delta) D^2\Delta'^2 + \dots + D^n \Delta'^n.$$

Hence, observing that $f(\Delta) v_x = 0$, since v_x substituted for u_x satisfies the proposed equation, we get the depressed equation

$$w_x f_1(\Delta) v_{x+1} + \frac{1}{1 \cdot 2} \Delta w_x f_2(\Delta) v_{x+2} + \dots + \Delta^{n-1} w_x \cdot v_{x+n} = 0;$$

or, reversing the order of the terms, {since $f_r(\Delta)$ means the same function of Δ , that $\frac{d^r f(x)}{dx^r}$ does of x },

$$\begin{aligned} & v_{x+n} \Delta^{n-1} w_x + (n\Delta v_{x+n-1} + q_1 v_{x+n-1}) \Delta^{n-2} w_x \\ & + \left\{ \frac{n(n-1)}{1 \cdot 2} \Delta^2 v_{x+n-2} + (n-1) q_1 \Delta v_{x+n-2} + q_2 v_{x+n-2} \right\} \Delta^{n-3} w_x + \dots \\ & + \{ n\Delta^{n-1} v_{x+1} + (n-1) q_1 \Delta^{n-2} v_{x+1} + \dots + q_{n-1} v_{x+1} \} w_x = 0 \dots \dots (2), \end{aligned}$$

a linear equation of the $(n-1)^{\text{th}}$ order, of the same form as the original one.

Similarly, if we know another particular value of u_x , z_x , then $\Delta \left(\frac{z_x}{v_x} \right)$ will be a value of w_x in equation (2), which may be depressed to another of the same form of the $(n-2)^{\text{th}}$ order; and if we know r particular solutions of equation (1), we may in this way depress it to an equation of the same form of the $(n-r)^{\text{th}}$ order. As a linear equation of differences of the first order and degree can always be solved, it appears that to obtain the complete integral of a linear equation of the n^{th} order, we must know $n-1$ particular solutions.

Equations of Differences which admit of being reduced to Linear Equations. Simultaneous Equations.

86. Besides linear equations of differences, and such equations as can be reduced to that form, very little is known of equations of differences of the second and higher orders and

degrees. Where such equations can be solved, it is generally by means of particular substitutions adapted to the form of the proposed equation; or by transformations of a more intricate character. The following are instances of equations which admit of being reduced to linear equations with constant coefficients, by particular substitutions.

$$1. \quad u_{x+n} + p_1 v_{x+n} u_{x+n-1} + p_2 v_{x+n} v_{x+n-1} u_{x+n-2} + \dots \\ + p_n v_{x+n} v_{x+n-1} \dots v_{x+1} u_x = 0;$$

where $p_1, p_2, \&c.$ are constants, and v_x is any function of x . Assume $u_x = v_x P v_x$, then the equation becomes divisible by $P v_{x+n}$, and is reduced to the linear equation with constant coefficients,

$$w_{x+n} + p_1 w_{x+n-1} + p_2 w_{x+n-2} + \dots + p_n w_x = 0.$$

$$\text{Ex.} \quad u_{x+2} - 2(x+2)u_{x+1} - 3(x+2)(x+1)u_x = 0.$$

$$u_x = \{a3^x + b(-1)^x\} \times 1.2.3\dots x.$$

$$2. \quad u_{x+n} + p_1 a^x u_{x+n-1} + p_2 a^{2x} u_{x+n-2} + \dots + p_n a^{nx} u_x = 0.$$

Assume $u_x = v_x a^{(1+2+3\dots+x)}$, then any term $p_r a^{rx} u_{x+n-r}$ becomes

$$= p_r v_{x+n-r} a^{\frac{1}{2}x(x+2n+1)} \cdot a^{\frac{1}{2}(n-r)(n-r+1)};$$

therefore the equation becomes divisible by $a^{\frac{1}{2}x(x+2n+1)}$, and is reduced to the linear equation with constant coefficients,

$$a^{\frac{1}{2}(n+1)n} v_{x+n} + p_1 a^{\frac{1}{2}n(n-1)} v_{x+n-1} + p_2 a^{\frac{1}{2}(n-1)(n-2)} v_{x+n-2} + \dots + p_n v_x = 0.$$

$$3. \quad u_{x+2} + \{a + b(-1)^x\} u_{x+1} + cu_x = 0.$$

$$\text{Let } u_x = v_x \sqrt{a + b(-1)^x},$$

$$\text{then } u_{x+1} = v_{x+1} \sqrt{a - b(-1)^x},$$

$$u_{x+2} = v_{x+2} \sqrt{a + b(-1)^x};$$

$$\therefore v_{x+2} + \sqrt{a^2 - b^2} v_{x+1} + cv_x = 0.$$

$$4. \quad u_{x+1} u_x + au_{x+1} + bu_x + c = 0.$$

Assume $u_x + a = \frac{v_{x+1}}{v_x}$, then $u_{x+1} + a = \frac{v_{x+2}}{v_{x+1}}$,

$$\therefore \frac{v_{x+1}}{v_x} \left(\frac{v_{x+2}}{v_{x+1}} - a \right) + b \left(\frac{v_{x+1}}{v_x} - a \right) + c = 0,$$

$$\text{or } v_{x+2} - (a-b)v_{x+1} + (c-ab)v_x = 0.$$

The two arbitrary constants which will appear in the value of v_x must be reduced to a single constant by the condition of the proposed equation being satisfied.

Ex. $u_x u_{x+1} - 2u_x + 1 = 0. \quad u_x = 1 + (x+c)^{-1}.$

5. $u_{x+n} \cdot u_{x+n-1}^p \dots u_x^r = X.$

If we take the logarithm of both sides, and assume $\log u_x = v_x$, we get

$$v_{x+n} + p v_{x+n-1} + q v_{x+n-2} + \dots + r v_x = \log X;$$

from which we can determine v_x , and then we have $u_x = e^{v_x}$.

Thus, if $u_{x+2} u_{x+1}^{-2} \cdot u_x = 1$, then $v_{x+2} - 2v_{x+1} + v_x = 0$;

$$\therefore v_x = c_0 + c_1 x, \text{ and } u_x = C' \cdot C^x.$$

Again if $u_{x+2} \cdot u_{x+1}^{2m} \cdot u_x^{m^2} = a$, then $v_{x+2} + 2m v_{x+1} + m^2 v_x = \log a$;

$$\therefore v_x = \frac{\log a}{(1+m)^2} + (c_0 + c_1 x) (-m)^x,$$

which is the value of $\log u_x$.

6. $u_{x+1}^2 + u_x = 2.$

Assume $u_x = 2 \cos v_x$, then $4 \cos^2 v_{x+1} + 2 \cos v_x = 2$,

$$\therefore 2 \cos^2 v_{x+1} = 1 - \cos v_x = 2 \sin^2 \frac{1}{2} v_x,$$

$$\text{or } \cos v_{x+1} = \cos \frac{1}{2} (\pi - v_x).$$

$$\therefore v_{x+1} + \frac{1}{2} v_x = \frac{1}{2} \pi, \text{ which gives } v_x = \frac{2}{3} \cdot \frac{1}{2} \pi + c \left(-\frac{1}{2}\right)^x,$$

$$\therefore u_x = 2 \cos \left\{ \frac{1}{3} \pi + c \left(-\frac{1}{2}\right)^x \right\}.$$

Similarly, $u_{x+1} - 2u_x^2 + 1 = 0$, by assuming $u_x = \cos v_x$, gives $\cos v_{x+1} = \cos 2v_x$, $\therefore v_{x+1} - 2v_x = 0$, $v_x = C2^x$, and

$$u_x = \cos v_x = \frac{1}{2} (e^{v_x \sqrt{-1}} + e^{-v_x \sqrt{-1}}) = \frac{1}{2} (c^{2^x} + c^{-2^x}) \text{ putting } c = e^{C\sqrt{-1}}.$$

$$7. \quad u_{x+1}u_x - X(u_{x+1} - u_x) + 1 = 0.$$

Let $u_x = \tan v_x$, $\therefore X(\tan v_{x+1} - \tan v_x) = 1 + \tan v_{x+1} \tan v_x$,

$$\text{or } \tan(v_{x+1} - v_x) = \frac{1}{X}, \quad \therefore \Delta v_x = \tan^{-1} \frac{1}{X},$$

$$\text{and } u_x = \tan \left(C + \sum \tan^{-1} \frac{1}{X} \right).$$

$$\text{If } X = 1 + x + x^2, \quad u_x = \frac{x + c}{1 - cx}.$$

Similarly $u_{x+2}u_{x+1}u_x = au_x \pm a(u_{x+1} + u_{x+2})$,

by assuming $u_x = \sqrt{a} \cdot \tan v_x$, becomes

$$\tan v_x (1 - \tan v_{x+1} \tan v_{x+2}) \pm (\tan v_{x+2} + \tan v_{x+1}) = 0,$$

$$\text{or } \tan v_x \pm \tan(v_{x+2} + v_{x+1}) = 0,$$

$\therefore v_{x+2} + v_{x+1} \pm v_x = r\pi$, where r is any integer ;

$$\therefore v_x = \frac{1}{3}r\pi + c \cos \frac{2\pi x}{3} + c' \cdot \sin \frac{2\pi x}{3},$$

$$\text{or } v_x = r\pi + c \left(2 \sin \frac{\pi}{10} \right)^x + c' \left(-2 \sin \frac{3\pi}{10} \right)^x,$$

according as the upper or lower sign is taken ; and then

$$u_x = \sqrt{a} \tan v_x.$$

87. We shall next give some examples of equations which by particular substitutions are reduced to linear equations of the first order.

$$1. \quad (ax + b) \Delta^2 u_x + mx \Delta u_x + mn u_x = 0.$$

Assume $u_x = \Delta^{n-1} v_x$;

$$\therefore (ax + b) \Delta^{n+1} v_x + mx \Delta^n v_x + mn \Delta^{n-1} v_x = 0.$$

But putting $u_x = x - n$ in the formula for transforming $u_{x+n} \Delta^n v_x$ (Art. 34), we have

$$\begin{aligned} x \Delta^n v_x &= \Delta^n \{(x-n) v_x\} - n \Delta^{n-1} v_x, \\ \therefore a \Delta^{n+1} \{(x-n-1) v_x\} - a(n+1) \Delta^n v_x \\ &\quad + b \Delta^{n+1} v_x + m \Delta^n \{(x-n) v_x\} = 0, \\ \text{or } \Delta \{a(x-n-1) + b\} v_x - \{a(n+1) - m(x-n)\} v_x &= \Sigma^n 0, \\ \therefore \{a(x-n) + b\} v_{x+1} + \{(x-n)(m-a) - na - b\} v_x &= \Sigma^n 0, \end{aligned}$$

a linear equation of the first order, which will furnish the value of v_x , and thence the complete integral of the proposed; the supernumerary constants contained in

$$\Sigma^n 0 = c_0 + c_1 x + \dots + c_{n-1} x^{n-1}$$

being determined by substituting the value of u_x in the proposed. For a particular solution, supposing $\Sigma^n 0 = 0$, and putting

$$A_x = \frac{(a-m)(x-n) + na + b}{a(x-n) + b},$$

we get $v_x = C \cdot P A_{x-1}$, and $u_x = C \Delta^{n-1} (P A_{x-1})$.

$$2. \quad (ax^2 + b) \Delta^2 u_x + (r + mx) \Delta u_x + (m-n)(n+1) u_x = 0.$$

Let $u_x = \Delta^{n-1} v_x$, then

$$(ax^2 + b) \Delta^{n+1} v_x + (r + mx) \Delta^n v_x + (m-n)(n+1) \Delta^{n-1} v_x = 0 \dots (1).$$

But if in the formula for transforming $u_{x+n} \Delta^n v_x$ (Art. 34) we make $u_x = x - n$, $(x-n)^2$, successively, we get values of $x \Delta^n v_x$, $x^2 \Delta^{n+1} v_x$; and substituting these values in (1), we find

$$\begin{aligned} &a \Delta^{n+1} \{(x-n-1)^2 v_x\} - a(n+1) \Delta^n \{(2x-2n-1) v_x\} \\ &+ a(n+1) n \Delta^{n-1} v_x + b \Delta^{n+1} v_x + r \Delta^n v_x + m \Delta^n \{(x-n) v_x\} \\ &\quad - mn \Delta^{n-1} v_x + (m-n)(n+1) \Delta^{n-1} v_x = 0, \\ &\text{or } \Delta^{n+1} \{a(x-n-1)^2 + b\} v_x \\ &\quad - \Delta^n \{a(n+1)(2x-2n-1) - m(x-n) - r\} v_x \\ &\quad + \{(a-1)(n+1)n + m\} \Delta^{n-1} v_x = 0, \end{aligned}$$

which is integrable in several cases by the disappearance of one of its terms; thus suppose $a = 1$, $m = 0$, then

$$\Delta \{(x - n - 1)^2 + b\} v_x - \{(n + 1)(2x - 2n - 1) - r\} v_x = \Sigma^n 0,$$

$$\text{or } \{(x - n)^2 + b\} v_{x+1} - \{x^2 - n(n + 1) + b - r\} v_x = \Sigma^n 0,$$

a linear equation of the first order, from which v_x and $u_x = \Delta^{n-1} v_x$ may be determined; and the arbitrary constants contained in $\Sigma^n 0$ must be reduced to the proper number by substituting the value of u_x in the proposed equation. But if we suppose the complementary function $\Sigma^n 0$ to vanish, we may readily obtain two particular values of v_x ; for let

$$\frac{x^2 - n(n + 1) + b - r}{(x - n)^2 + b} = A_x,$$

$$\text{then } v_{x+1} - A_x v_x = 0, \quad v_x = C \cdot PA_{x-1}.$$

Now replace n by $-(n + 1)$ which amounts to assuming $u_x = \Delta^{-n-2} v_x$, then $v_x = C' \cdot PA'_{x-1}$, where

$$\frac{x^2 - n(n + 1) + b - r}{(x + n + 1)^2 + b} = A'_x;$$

$$\text{then } u_x = C \Delta^{n-1} (PA_{x-1}) + C' \Sigma^{n+2} (PA'_{x-1}),$$

the complete integral of the proposed equation.

Several other examples of the like kind may be seen in the Cambridge and Dublin Mathematical Journal.

87*. The solution of $f(D) u_x = b^x$ in the form (Art. 83)

$$u_x = \frac{b^x}{f(b)} + Ca_1^x + \&c.$$

fails, if b be equal to a root of the auxiliary equation $f(a) = 0$. Suppose that $b = a_1$ a root that occurs twice in $f(a) = 0$; then the value of u_x by changing the constants may be put under the form

$$u_x = \frac{1}{f'(b)} (b^x - a_1^x - xha_1^{x-1}) + (c_0 + c_1x) a_1^x + \&c.$$

Now suppose $b = a_1 + h$, where h is very small; then since $f(a_1) = 0$, $f'(a_1) = 0$, we have

$$u_x = \frac{\frac{1}{2}x(x-1)h^2a_1^{x-2} + \&c.}{\frac{1}{2}h^2f''(a_1) + \&c.} + (c_0 + c_1x)a_1^x + \&c.;$$

therefore, making $h = 0$ or $b = a_1$, we get

$$u_x = \frac{x(x-1)}{f''(a_1)}a_1^{x-2} + (c_0 + c_1x)a_1^x + \&c.$$

And if the root to which b becomes equal occur r times, the first term in the value of u_x will be

$$x(x-1)\dots(x-r+1)a_1^{x-r} \div f^{(r)}(a_1).$$

$$\text{Hence in } u_{x+2} - 2a_1u_{x+1} + a_1^2u_x = a_1^x,$$

$$\text{since } f(a) = (a - a_1)^2 \quad f''(a_1) = 2,$$

$$u_x = \frac{1}{2}x(x-1)a_1^{x-2} + (c_0 + c_1x)a_1^x;$$

and in examples 3 and 4 of Art. 83 if $\theta = 0$, the first term in the value of u_x becomes

$$\frac{1}{2}x(x-1)a^{x-2}, \text{ and } \frac{1}{24}x(x-1)(x-2)(x-3)a^{x-4},$$

respectively.

88. In simultaneous equations, instead of one equation between the independent variable x , and the successive values or differences of the dependent function u_x , we have given two or more equations between x and two or more of its functions u_x , v_x , &c. and their successive values or differences; the object, as before, being to determine each of the unknown functions in terms of x . When the proposed equations are linear with constant coefficients, we may separate the symbols of operation from those of quantity; and then obtain by the ordinary processes of elimination an equation containing only one of the unknown functions; for as the symbols of operation here employed combine according to the same laws as ordinary algebraical quantities, they may be treated precisely as if they were symbols of quantity.

$$\begin{aligned} 1. \quad & u_{x+2} + cv_{x+1} - c^3u_x = a^x, \\ & v_{x+2} - u_{x+1} - cv_x = a^{-x}. \end{aligned}$$

These may be written

$$\begin{aligned} (D^2 - c^3) u_x + cDv_x &= a^x, \\ (D^2 - c) v_x - Du_x &= a^{-x}, \end{aligned}$$

which give by the elimination of v_x

$$(D^4 - c^3D^2 + c^4) u_x = (D^2 - c) a^x - cDa^{-x};$$

$$\therefore u_x = \frac{(a^2 - c) a^x}{a^4 - c^3a^2 + c^4} - \frac{ca^{-x+3}}{1 - c^3a^2 + c^4a^4} + \text{complementary function.}$$

Now if c be greater than 2,

$$D^4 - c^3D^2 + c^4 = (D^2 - \alpha^2)(D^2 - \beta^2),$$

and consequently the complementary function

$$= c_1 a^x + c_2 (-\alpha)^x + c_3 \beta^x + c_4 (-\beta)^x.$$

If $c=2$, the values of α and β become equal to one another; or imaginary if $c < 2$; and the corresponding values of the complementary function may be determined in the usual manner.

The value of u_x being known, we have

$$cv_x = \frac{a^x - (D^2 - c^3) u_x}{D} = a^{x-1} - u_{x+1} + c^3 u_{x-1}.$$

$$\begin{aligned} 2. \quad & u_{x+1} = 2z_x + 12u_x, \quad y_{x+1} = 14z_x, \\ & z_{x+1} = y_x + 6z_x + 7u_x. \end{aligned}$$

Upon introducing the symbol D , and eliminating u_x and y_x , we find $(D-6)(D+2)(D-14)z_x = 0$, which gives

$$z_x = a \cdot 6^x + b \cdot (-2)^x + c(14)^x;$$

$$\text{then } y_x = 14z_{x-1} = \frac{7a}{3} 6^x - 7b(-2)^x + c(14)^x,$$

$$u_x = -\frac{a}{3} 6^x - \frac{b}{7} (-2)^x + c(14)^x.$$

89. It may be observed that the knowledge of the generating function of the second member of a linear equation will sometimes lead to the solution of the equation. For since (Art. 28)

$$G\{\psi(D)u_x\} = \psi\left(\frac{1}{t}\right)Gu_x; \quad \therefore Gu_x = \frac{G\{\psi(D)u_x\}}{\psi\left(\frac{1}{t}\right)}.$$

Now suppose $\psi(D)u_x = X$ to be an equation of Differences, and let $\phi(t)$ be the generating function of X , and therefore of its equal $\psi(D)u_x$, then

$$Gu_x = \frac{\phi(t)}{\psi\left(\frac{1}{t}\right)};$$

and if this value of Gu_x can be developed in a series of powers of t , and have w_x for the coefficient of t^x , then the solution of $\psi(D)u_x = X$ is $u_x = w_x + \text{complementary function}$.

Thus, for $u_{x+2} - 2 \cos \theta u_{x+1} + u_x = \sin x\theta$,

$$\phi(t) = \frac{t \sin \theta}{1 - 2t \cos \theta + t^2} = \dots + t^x \sin x\theta + \dots \quad (\text{Trig. Art. 155})$$

$$\begin{aligned} \phi(t) \div \psi\left(\frac{1}{t}\right) &= \frac{t^3 \sin \theta}{(1 - 2t \cos \theta + t^2)^2} = \dots \\ &+ \frac{t^x}{2 \sin^2 \theta} \{\sin x\theta - x \cos(x-1)\theta \sin \theta\} + \dots \end{aligned}$$

the latter expansion being derived from the former by differentiating relative to θ , and reducing;

$\therefore u_x = \frac{1}{2 \sin^2 \theta} \{\sin x\theta - x \cos(x-1)\theta \sin \theta\} + \text{complementary function}$

$$= -\frac{x}{2 \sin \theta} \cos(x-1)\theta + c \cos x\theta + c' \sin x\theta.$$

Again, let the proposed equation be $(D-1)^{n-2}u_x = x$, then (Art. 26) $\phi(t) = t(1-t)^{-2}$,

$$\begin{aligned} \therefore \frac{\phi(t)}{\psi\left(\frac{1}{t}\right)} &= \frac{t(1-t)^{-2}}{\left(\frac{1}{t}-1\right)^{n-2}} = t^{n-1}(1-t)^{-n} \\ &= t^{n-1} \left\{ 1 + nt + \frac{n(n+1)}{1 \cdot 2} t^2 + \dots + \frac{n(n+1) \dots x}{1 \cdot 2 \cdot 3 \dots (x-n+1)} t^{x-n+1} + \dots \right\}; \end{aligned}$$

$$\therefore w_x = \frac{x(x-1)\dots n}{1.2.3\dots(x-n+1)} = \frac{x(x-1)\dots(x-n+2)}{1.2.3\dots(n-1)};$$

$$\therefore u_x = \frac{x(x-1)\dots(x-n+2)}{1.2.3\dots(n-1)} + \alpha + \beta x + \dots + \lambda x^{n-3},$$

which agrees with the result obtained by direct integration of $\Delta^{n-2}u_x = x$.

Equations of Partial Differences and Mixed Differences.

90. If in $u_{x,y}$ a function of two independent variables x and y , x be changed into $x+h$, and y into $y+k$, the resulting value $u_{x+h,y+k}$ is called the first total successive value of $u_{x,y}$, and is denoted by $Du_{x,y}$, so that $Du_{x,y} = u_{x+h,y+k}$; consequently

$$D^2u_{x,y} = u_{x+2h,y+2k}, \text{ \&c. } D^n u_{x,y} = u_{x+nh,y+nk};$$

and as $u_{x+h,y+k} = e^{\frac{d}{dx} + \frac{d}{dy}} u_{x,y}$, D is here equivalent to $e^{\frac{d}{dx} + \frac{d}{dy}}$. Also if from the altered value of the function we subtract the original value, the result is called the first total difference of $u_{x,y}$, and is denoted by $\Delta u_{x,y}$, so that

$$\Delta u_{x,y} = u_{x+h,y+k} - u_{x,y} = Du_{x,y} - u_{x,y} = (D-1)u_{x,y}.$$

Consequently we have, as in the case of a function of one variable, Δ equivalent to $D-1$; and $1+\Delta$ equivalent to D ; hence we must have

$$\Delta^2 u_{x,y} = (D-1)^2 u_{x,y}, \quad D^2 u_{x,y} = (1+\Delta)^2 u_{x,y}, \text{ \&c. ;}$$

and generally

$$\Delta^n u_{x,y} = (D-1)^n u_{x,y}, \quad D^n u_{x,y} = (1+\Delta)^n u_{x,y};$$

which become by development

$$\Delta^n u_{x,y} = D^n u_{x,y} - nD^{n-1}u_{x,y} + \frac{n(n-1)}{1.2} D^{n-2}u_{x,y} - \text{\&c.}$$

$$= u_{x+nh,y+nk} - nu_{x+(n-1)h,y+(n-1)k} + \text{\&c.},$$

$$D^n u_{x,y} = u_{x,y} + n\Delta u_{x,y} + \frac{n(n-1)}{1.2} \Delta^2 u_{x,y} + \text{\&c.};$$

where Δ , D express total operations that refer to both of the independent variables.

If there be given $\Delta u_{x,y} = f(x, y)$, a rational and integral function of x and y , we may find the value of $u_{x,y}$ by the method of Art. 44; thus if $\Delta u_{x,y} = x + y + 1$, then

$$u_{x,y} = \frac{1}{2}(x^2 + y^2) + C_{x-y},$$

where C_{x-y} denotes any function of $x - y$, the increments of both variables being taken equal to unity.

91. Again, if in $u_{x,y}$ we change x into $x + h$ without altering y , the result $u_{x+h,y}$ is called the first partial successive value relative to x , and is denoted by $D_x u_{x,y}$, so that $D_x u_{x,y} = u_{x+h,y}$; consequently

$$D_x^2 u_{x,y} = u_{x+2h,y}, \text{ \&c., } D_x^n u_{x,y} = u_{x+nh,y};$$

and as $u_{x+h,y} = e^{h \frac{d}{dx}} u_{x,y}$, D_x is equivalent to $e^{h \frac{d}{dx}}$. Exactly in the same way we have, relative to the other variable y ,

$$D_y u_{x,y} = u_{x,y+k}, \quad D_y^n u_{x,y} = u_{x,y+nk},$$

D_y being equivalent to $e^{k \frac{d}{dy}}$; hence $D = D_x \cdot D_y$. Also the differences $u_{x+h,y} - u_{x,y}$, $u_{x,y+k} - u_{x,y}$, are called the partial differences of $u_{x,y}$ relative to x and y respectively, and are denoted by Δ_x , Δ_y ; so that

$$\Delta_x u_{x,y} = u_{x+h,y} - u_{x,y} = (D_x - 1) u_{x,y};$$

$$\Delta_y u_{x,y} = u_{x,y+k} - u_{x,y} = (D_y - 1) u_{x,y};$$

and generally

$$\Delta_x^n u_{x,y} = (D_x - 1)^n u_{x,y}, \quad \Delta_y^n = (D_y - 1)^n u_{x,y};$$

and the symbols Σ_x , Σ_y , which are equivalent to Δ_x^{-1} , Δ_y^{-1} , are used to denote the operation of integrating $u_{x,y}$ with respect to the variable x only, and the variable y only, respectively.

In the investigations which follow, the values of h and k will be taken equal to unity, unless the contrary be stated.

92. By Equations of Partial Differences are meant relations between a function of two independent variables $u_{x,y}$, some of its partial successive values or differences, and the variables x and y . As the symbols of operation here employed combine in subjection to the same fundamental laws as algebraical quantities, we may, by separating the symbols, resolve these equations when linear or otherwise, by the same processes as ordinary equations of differences of the like class: but we shall first employ an elementary method.

Ex. 1. $u_{x+n,y} + p_1 u_{x+n-1,y+1} + \dots + p_n u_{x,y+n} = 0(1)$, the coefficients being constant, and the sum of the subscribed indices the same in every term. If the second member be q instead of zero, it is easily seen that the value of $u_{x,y}$ determined from (1) would have to be increased by $\frac{q}{f(1)}$.

Assume $u_{x,y} = a^x \cdot C_{x+y}$, where C_{x+y} is any function of the binomial $x + y$, then the first member of (1) becomes

$$a^x (a^n + p_1 a^{n-1} + \dots + p_n) C_{x+y+n}$$

which is reduced to zero if a be taken equal to any one of the roots real or imaginary, $a_1, a_2, \dots a_n$ of the auxiliary equation

$$f(a) = a^n + p_1 a^{n-1} + \dots + p_n = 0.$$

Hence, if $a_1, a_2, \&c.$, be all unequal, the complete value of $u_{x,y}$ involving n arbitrary functions is

$$u_{x,y} = a_1^x C'_{x+y} + a_2^x C''_{x+y} + \dots + a_n^x C^{(n)}_{x+y},$$

which shews that arbitrary functions of the binomial $x + y$ here stand in the places of the arbitrary constants that enter into the solutions of ordinary equations of Differences. But if $a_2 = a_1$, first suppose $a_2 = a_1 + h$ where h is very small, then

$$\begin{aligned} a_1^x C'_{x+y} + a_2^x C''_{x+y} &= a_1^x (C'_{x+y} + C''_{x+y}) \\ &+ \frac{h}{a_1} C''_{x+y} \cdot x a_1^x \{1 + \frac{1}{2}(x-1) \frac{h}{a_1} + \&c.\}, \end{aligned}$$

$$= a_1^x \cdot c'_{x+y} + c''_{x+y} \cdot x a_1^x \{1 + \frac{1}{2}(x-1) \frac{h}{a_1} + \&c.\},$$

replacing the arbitrary functions

$$C'_{x+y} + C''_{x+y} \quad \text{and} \quad \frac{h}{a_1} C''_{x+y}$$

by other arbitrary functions c'_{x+y} and c''_{x+y} ; now make $h = 0$, then the equal roots a_1, a_2 produce in the value of $u_{x,y}$ the terms $a_1^x (c'_{x+y} + x c''_{x+y})$. If a_1, a_2 be a pair of imaginary roots and

$$= \rho (\cos \theta \pm \sqrt{-1} \sin \theta),$$

then it is easily seen that $a_1^x C'_{x+y} + a_2^x C''_{x+y}$ takes the form

$$\rho^x (c_{x+y} \cos x\theta + c'_{x+y} \sin x\theta).$$

It may be observed that the assumption $u_{x,y} = a^x \cdot C_{x+y}$ is symmetrical with respect to x and y , for it may be replaced by

$$u_{x,y} = \left(\frac{1}{a}\right)^y \cdot a^{x+y} \cdot C_{x+y} = b^y \cdot C'_{x+y}.$$

Ex. 1. $u_{x+1,y} - a u_{x,y+1} = b$. Here $u_{x,y} = a^x \cdot C_{x+y} + \frac{b}{1-a}$.

Ex. 2. $u_{x+2,y} - 2a u_{x+1,y+1} + a^2 u_{x,y+2} = b$;

$$u_{x,y} = a^x (c_{x+y} + x \cdot c'_{x+y}) + \frac{b}{(1-a)^2}.$$

Ex. 3. $u_{x+2,y} - 2a \cos \theta u_{x+1,y+1} + a^2 u_{x,y+2} = b$,

$$u_{x,y} = a^x (c_{x+y} \cos x\theta + c'_{x+y} \sin x\theta) + \frac{b}{1-2a \cos \theta + a^2}.$$

93. The preceding results can of course be readily obtained by separating the symbols; but that method may be applied with still greater advantage to the case where the second member is V a function of x and y . For the proposed equation may be written

$$\{(D_x D_y^{-1})^n + p_1 (D_x D_y^{-1})^{n-1} + \dots + p_n\} D_y^n u_{x,y} = V,$$

$$\text{instead of } u_{x+n,y} + p_1 u_{x+n-1,y+1} + \dots + p_n u_{x,y+n} = V.$$

Let a be a root that occurs singly, and b a root that occurs m times, in the auxiliary equation $z^n + p_1 z^{n-1} + \dots + p_n = 0$: then by resolution into partial fractions we shall have

$$\frac{1}{z^n + p_1 z^{n-1} + \dots + p_n} = \frac{A}{z-a} + \frac{B_m}{(z-b)^m} + \frac{B_{m-1}}{(z-b)^{m-1}} + \dots + \frac{B_1}{z-b} + \&c.,$$

where $A, B_m, \&c.$ are known in terms of $p_1, p_2, \&c.$; hence substituting $D_x D_y^{-1}$ for z , we get

$$u_{x,y} = \left\{ \frac{A D_y}{D_x - a D_y} + \frac{B_m D_y^m}{(D_x - b D_y)^m} + \frac{B_{m-1} D_y^{m-1}}{(D_x - b D_y)^{m-1}} + \dots + \frac{B_1 D_y}{D_x - b D_y} \right\} D_y^{-n} V;$$

$$\text{but } A (D_x - a D_y)^{-1} D_y^{-n+1} V = A (a D_y)^{x-1} \Sigma_x (a D_y)^{-x} D_y^{-n+1} V; \&c.;$$

since the operations denoted by D_x, D_y , are independent of one another:

$$\therefore u_{x,y} = A (a D_y)^{x-1} \Sigma_x (a D_y)^{-x} D_y^{-n+1} V + B_m (b D_y)^{x-m} \Sigma_x^m (b D_y)^{-x} D_y^{-n+m} V + \&c.$$

Instead of reserving the complementary functions under the sign of integration, we may obtain them explicitly by supposing $V=0$; then since

$$\Sigma_x 0 = c_y, \quad \Sigma_x^m 0 = c_y^0 + x c_y' + x^2 c_y'' + \dots + x^{m-1} c_y^{(m-1)},$$

$$\text{and } D_y^{x-1} c_y = c_{x+y-1}, \quad D_y^{x-m} c_y = c_{x+y-m},$$

the part of the complementary function due to the root a and the m roots b will be, changing the arbitrary functions,

$$A a^x c_{x+y} + B_m b^x \{c_{x+y}^0 + x c_{x+y}' + \dots + x^{m-1} c_{x+y}^{(m-1)}\},$$

and similar terms will be introduced by the other roots of the auxiliary equation.

If any of the roots $a, b, \&c.$ be imaginary, these results may be transformed into real expressions, as shewn above.

Ex. 1. $u_{x+1,y} - u_{x,y+1} = xy,$

or $(D_x - D_y) u_{x,y} = xy, \therefore u_{x,y} = (D_x - D_y)^{-1} xy = (D_y)^{x-1} \Sigma_x D_y^{-x} xy;$

but $D_y^{-x} xy = x(y-x) = x(y+1) - x(x+1);$

$$\begin{aligned} \therefore u_{x,y} &= D_y^{x-1} \left\{ \frac{(x-1)x(y+1)}{2} - \frac{(x-1)x(x+1)}{3} + C_y \right\} \\ &= \frac{1}{6} (x-1)x(x+3y-2) + C_{x+y}. \end{aligned}$$

Ex. 2. $u_{x+2,y} - 2u_{x+1,y+1} + u_{x,y+2} = x+y,$

or $(D_x - D_y)^2 u_{x,y} = x+y;$

$$\begin{aligned} \therefore u_{x,y} &= (D_x - D_y)^{-2} (x+y) = D_y^{x-2} \Sigma_x^2 D_y^{-x} (x+y) \\ &= D_y^{x-2} \Sigma_x^2 y = D_y^{x-2} \left\{ \frac{1}{2} (x-1) xy + x C_y + C'_y \right\} \\ &= \frac{1}{2} (x-1)x(y+x-2) + x c_{x+y} + c'_{x+y}. \end{aligned}$$

Ex. 3. $u_{x+3,y} - 12u_{x+1,y+2} + 16u_{x,y+3} = c^{x+y},$

or $(D_x - 2D_y)^2 (D_x + 4D_y) u_{x,y} = c^{x+y};$

and since D_x, D_y , applied to c^{x+y} are each equivalent to multiplying it by c ,

$$\therefore u_{x,y} = \frac{c^{x+y}}{(-c)^2 \cdot 5c} + \&c. = \frac{1}{5} c^{x+y-3} + 2^x (c_{x+y} + x c'_{x+y}) + (-4)^x c''_{x+y}.$$

Ex. 4. $u_{x+2,y} - 2a \cos \theta u_{x+1,y+1} + a^2 u_{x,y+2} = V.$

$$\begin{aligned} u_{x,y} &= a^{x-2} \frac{\sin(x-1)\theta}{\sin \theta} D_y^{x-1} \Sigma_x (a^{-x} \cos x\theta D_y^{-x-1} V) \\ &\quad - a^{x-2} \frac{\cos(x-1)\theta}{\sin \theta} D_y^{x-1} \Sigma_x (a^{-x} \sin x\theta D_y^{-x-1} V) \\ &\quad + a^x (\cos x\theta \cdot c_{x+y} + \sin x\theta \cdot c'_{x+y}). \end{aligned}$$

Ex. 5. $(D_x - a)(D_y - b) u_{x,y} = V,$

or $u_{x+1,y+1} - a u_{x,y+1} - b u_{x+1,y} + a b u_{x,y} = V;$

$$\therefore (D_y - b) u_{x,y} = (D_x - a)^{-1} V = a^{x-1} \Sigma_x a^{-x} V + a^{x-1} C_y,$$

$$\begin{aligned} u_{x,y} &= a^{x-1} b^{y-1} \Sigma_y (b^{-y} \Sigma_x a^{-x} V) + a^{x-1} b^{y-1} C'_y + a^{x-1} C_y \\ &= a^{x-1} b^{y-1} \Sigma_y (b^{-y} \Sigma_x a^{-x} V) + b^y c'_x + a^x c''_y. \end{aligned}$$

If $V = c^{x-y}$, since D_x, D_y are in that case equivalent respectively to c and c^{-1} ;

$$u_{x,y} = \frac{c^{x-y}}{(c-a)(c^{-1}-b)} + b^y c'_x + a^x c''_y.$$

Equations of Mixed Differences.

94. Equations of Mixed Differences are those in which the differential coefficients of the dependent variable u_x , as well as its differences or successive values, are involved along with the principal variable x . When the equation is linear with respect to the successive values and their differential coefficients, and the coefficients of the several terms are constant, solutions may be obtained by proceeding in the same way as for linear equations of differences of the like kind. It is plain that any differential equation

$$f(x, y, \frac{dy}{dx}, \frac{d^2y}{dx^2}, \&c.) = 0,$$

which admits of a solution $y = X$, will lead by substituting

$$y = u_{x+n} + p_1 u_{x+n-1} + \dots + p_n u_x,$$

where $p_1, p_2, \&c.$ are constants, to a linear equation of differences of the kind that has been solved.

$$\begin{aligned} 1. \quad \frac{d}{dx}(u_{x+n} + p_1 u_{x+n-1} + \dots + p_n u_x) \\ + m(u_{x+n} + p_1 u_{x+n-1} + \dots + p_n u_x) = X. \end{aligned}$$

Multiplying both sides by e^{mx} and integrating, we get

$$u_{x+n} + p_1 u_{x+n-1} + \dots + p_n u_x = e^{-mx} \int dx e^{mx} X,$$

an ordinary linear equation of differences. Thus suppose the proposed equation to be

$$\frac{du_{x+1}}{dx} + a \frac{du_x}{dx} + bu_{x+1} + au_x = 0,$$

then $u_x = Cm^x + C'(-a)^x$, where $m = e^{-b}$.

$$2. \quad \frac{d}{dx} (u_{x+1} - 2u_x) + 3u_{x+1} - 6u_x = e^{mx},$$

$$\text{then } u_{x+1} - 2u_x = \frac{e^{mx}}{m+3} + C_0 e^{-3x};$$

$$\therefore u_x = \frac{e^{mx}}{(m+3)(e^m - 2)} + C \cdot e^{-3x} + C' \cdot 2^x.$$

$$3. \quad (1 + ae^{x+1}) u_{x+1} - b(1 + ae^x) \frac{d}{dx} u_x + bu_x = c.$$

$$\text{Assume } u_x(1 + ae^x) = v_x;$$

$$\therefore ae^x u_x + (1 + ae^x) \frac{d}{dx} u_x - \frac{dv_x}{dx} = 0,$$

and multiplying this by b and adding it to the proposed, we get

$$v_{x+1} - b \frac{dv_x}{dx} + bv_x = c.$$

$$\text{Now, let } v_x = e^{mx} + k;$$

$$\therefore e^{m(x+1)} + k - mb \cdot e^{mx} + be^{mx} + bk = c:$$

let $k + bk = c$, then $e^m - (m-1)b = 0$; and if μ be a value of m which satisfies this equation, then a particular value of u_x is

$$\frac{1}{1 + ae^x} \left(Ce^{\mu x} + \frac{c}{1+b} \right).$$

95. The following are examples of Equations of Mixed Partial Differences.

$$1. \quad u_{x+1,y} - \frac{d}{dy} u_{x,y} = c,$$

$$\text{or } D_x u_{x,y} - \frac{d}{dy} u_{x,y} = c,$$

$$\therefore u_{x,y} = \left(D_x - \frac{d}{dy} \right)^{-1} c = \left(\frac{d}{dy} \right)^{x-1} \left\{ \Sigma_x \left(\frac{d}{dy} \right)^{-x} c + \frac{d}{dy} C_y \right\},$$

putting $\frac{d}{dy} C_y$ for an arbitrary function of y ;

$$\therefore u_{x,y} = c + \left(\frac{d}{dy}\right)^x C_y;$$

and if $c = 0$, we have for the equation

$$u_{x+1,y} - \frac{d}{dy} u_{x,y} = 0, \quad u_{x,y} = \left(\frac{d}{dy}\right)^x C_y.$$

$$2. \quad u_{x+1,y} - a \frac{d}{dy} u_{x,y} = V;$$

$$\therefore u_{x,y} = \left(D_x - a \frac{d}{dy}\right)^{-1} V = \left(a \frac{d}{dy}\right)^{x-1} \Sigma_x \left(a \frac{d}{dy}\right)^{-x} V.$$

Suppose $V = xe^y$, then the effect of any power, positive or negative, of $\frac{d}{dy}$ upon V , is to multiply it by 1;

$$\therefore \Sigma_x \left(a \frac{d}{dy}\right)^{-x} V = e^y \Sigma_x a^{-x} x = \frac{e^y}{a^{x-1}} \frac{x(1-a) - 1}{(1-a)^2} + a \frac{d}{dy} C_y,$$

$$\therefore u_{x,y} = e^y \frac{x(1-a) - 1}{(1-a)^2} + a^x \left(\frac{d}{dy}\right)^x C_y.$$

$$3. \quad u_{x+2,y} - 2a \frac{d}{dy} u_{x+1,y} + a^2 \left(\frac{d}{dy}\right)^2 u_{x,y} = V;$$

$$\therefore u_{x,y} = \left(D_x - a \frac{d}{dy}\right)^{-2} V = \left(a \frac{d}{dy}\right)^{x-2} \Sigma_x^2 \left(a \frac{d}{dy}\right)^{-x} V.$$

Let $V = xe^y$, then proceeding as in the last example,

$$u_{x,y} = a^{x-2} e^y \Sigma^2 (a^{-x} x) = e^y \frac{x(1-a) - 2}{(1-a)^3} + a^x \left(\frac{d}{dy}\right)^x C_y + xa^x \left(\frac{d}{dy}\right)^x C_y^1.$$

$$4. \quad \Delta_x u_{x,y} - a \frac{d}{dy} u_{x,y} = c.$$

As this may be written

$$\Delta_x (u_{x,y} - cx) - a \frac{d}{dy} (u_{x,y} - cx) = 0,$$

we may integrate it supposing $c=0$, and then add cx to the result. Assume $u_{x,y} = a^x e^{-\frac{y}{a}} v_{x,y}$;

$$\therefore a^x e^{-\frac{y}{a}} (a v_{x+1,y} - v_{x,y}) - a^x e^{-\frac{y}{a}} \left(-v_{x,y} + a \frac{d}{dy} v_{x,y} \right) = 0;$$

$$\text{or } v_{x+1,y} - \frac{d}{dy} v_{x,y} = 0; \quad \therefore v_{x,y} = \left(\frac{d}{dy} \right)^x C_y. \quad (\text{Ex. 1});$$

$$\therefore u_{x,y} = cx + a^x e^{-\frac{y}{a}} \left(\frac{d}{dy} \right)^x C_y.$$

Continued Fractions.—Functional Equations.

96. The determination of the values of continued fractions in certain cases gives rise to equations of differences which can be integrated as linear equations.

1. To determine the value of the continued fraction,

$$u_x = \frac{c}{a + \frac{c}{a + \dots \frac{c}{a}}} \text{ to } x \text{ fractional terms};$$

$$\therefore u_{x+1} = \frac{c}{a + u_x}, \text{ or } u_{x+1} (u_x + a) = c;$$

Assume $u_x + a = \frac{v_{x+1}}{v_x}$, then $v_{x+2} - a v_{x+1} - c v_x = 0$;

$$\therefore v_x = c_1 \alpha^x + c_2 \beta^x, \alpha \text{ and } \beta \text{ being roots of } k^2 - ak - c = 0;$$

$$\therefore u_x = \frac{c_1 \alpha^{x+1} + c_2 \beta^{x+1}}{c_1 \alpha^x + c_2 \beta^x} - a = \frac{c_1 \alpha^x (\alpha - a) + c_2 \beta^x (\beta - a)}{c_1 \alpha^x + c_2 \beta^x}$$

$$= -\alpha \beta \frac{c_1 \alpha^{x-1} + c_2 \beta^{x-1}}{c_1 \alpha^x + c_2 \beta^x}, \text{ since } \alpha + \beta = a;$$

$$\text{or } u_x = c \cdot \frac{\alpha^{x-1} + c_3 \beta^{x-1}}{\alpha^x + c_3 \beta^x}, \text{ putting } c_3 = \frac{c_2}{c_1};$$

$$\therefore u_1 = \frac{c}{a} = c \frac{1 + c_3}{\alpha + c_3 \beta}, \quad \therefore c_3 = -\frac{\beta}{\alpha},$$

$$\therefore u_x = c \frac{\alpha^x - \beta^x}{\alpha^{x+1} - \beta^{x+1}}.$$

If α and β be imaginary, so that

$$k^2 - ak - c = k^2 - 2k\rho \cos \theta + \rho^2,$$

$$\text{then } u_x = -\rho \frac{\sin x\theta}{\sin (x+1)\theta}.$$

2. To find the value of the continued fraction

$$u_x = \frac{c}{a + \frac{c}{a + \dots \frac{c}{a + \frac{c}{b}}}},$$

to x fractional terms, the last term being $\frac{c}{b}$.

Proceeding as in the last example, we find

$$u_x = c \cdot \frac{\alpha^{x-1} + c_3 \beta^{x-1}}{\alpha^x + c_3 \beta^x},$$

$$\text{and } u_1 = \frac{c}{b} = c \cdot \frac{1 + c_3}{\alpha + c_3 \beta}, \quad \therefore c_3 = -\frac{b - \alpha}{b - \beta},$$

$$\therefore u_x = c \cdot \frac{b(\alpha^{x-1} - \beta^{x-1}) + c(\alpha^{x-2} - \beta^{x-2})}{b(\alpha^x - \beta^x) + c(\alpha^{x-1} - \beta^{x-1})}.$$

If $c = -1$, $\alpha = 2 \cos \theta$, $b = \cos \theta$, then it will be found that

$$u_x = -\frac{\cos (x-1)\theta}{\cos x\theta}.$$

3. To find the value of the continued fraction to x fractional terms,

$$u_x = \frac{1}{a + \frac{1}{b + \frac{1}{a + \frac{1}{b + \dots}}}}.$$

$$\text{Here } u_{x+2} = \frac{1}{a + \frac{1}{b + u_x}}, \quad \text{or } \frac{u_{x+2}}{1 - au_{x+2}} = b + u_x.$$

$$\text{Assume } 1 - au_x = \frac{v_{x-1}}{v_{x+1}}, \quad \therefore 1 - au_{x+2} = \frac{v_{x+1}}{v_{x+3}},$$

$$\therefore \frac{1}{a} \left(1 - \frac{v_{x+1}}{v_{x+3}} \right) \frac{v_{x+3}}{v_{x+1}} = b + \frac{1}{a} \left(1 - \frac{v_{x-1}}{v_{x+1}} \right),$$

$$\therefore v_{x+3} - v_{x+1} = abv_{x+1} + v_{x+1} - v_{x-1},$$

$$\text{or } v_{x+3} - (2 + ab) v_{x+1} + v_{x-1} = 0,$$

of which the auxiliary equation is

$$k^4 - (2 + ab) k^2 + 1 = (k^2 - \sqrt{ab} k - 1) (k^2 + \sqrt{ab} k - 1) = 0.$$

Let $\alpha, -\frac{1}{\alpha}, -\alpha, \frac{1}{\alpha}$, be the roots of this equation, then

$$v_x = \alpha^x \{c_1 + c_2 (-1)^x\} + \frac{1}{\alpha^x} \{c_3 + c_4 (-1)^x\},$$

$$\therefore au_x = 1 - \frac{\alpha^{x-1} + p_x \alpha^{-(x-1)}}{\alpha^{x+1} + p_x \alpha^{-(x+1)}},$$

$$\text{where } p_x = \frac{c_3 + c_4 (-1)^{x-1}}{c_1 + c_2 (-1)^{x-1}} = C + C' (-1)^{x-1};$$

$$\therefore au_1 = 1 = 1 - \frac{1 + p_1}{\alpha^2 + p_1 \alpha^2}, \quad \therefore p_1 = -1 = C + C',$$

$$au_2 = \frac{ab}{1 + ab} = 1 - \frac{\alpha + p_2 \alpha^{-1}}{\alpha^3 + p_2 \alpha^{-3}},$$

$$\therefore ab + 1 = \alpha^2 - 1 + \alpha^{-2} = \frac{\alpha^3 + p_2 \alpha^{-3}}{\alpha + p_2 \alpha^{-1}}, \quad \therefore \alpha - \frac{1}{\alpha} = \sqrt{ab};$$

$$\therefore p_2 = 1 = C - C', \quad \therefore C = 0, \quad C' = -1, \text{ and } p_x = (-1)^x,$$

$$\therefore au_x = 1 - \frac{\alpha^{x-1} + (-1)^x \alpha^{-x+1}}{\alpha^{x+1} + (-1)^x \alpha^{-x-1}}.$$

4. Suppose b to be essentially negative and $= -c$, so that the continued fraction is

$$\frac{1}{a +} \frac{1}{-c +} \frac{1}{a +} \frac{1}{-c +} \dots,$$

and let $\sqrt{ab} = \sqrt{-1} \sqrt{ac} = 2 \sqrt{-1} \sin \theta$, then

$$k^2 - 2 \sqrt{-1} \sin \theta k - 1 = 0, \text{ and } \alpha = \cos \theta + \sqrt{-1} \sin \theta;$$

therefore, according as x is odd, or even,

$$au_x = 1 - \frac{\sin(x-1)\theta}{\sin(x+1)\theta}, \text{ or } 1 - \frac{\cos(x-1)\theta}{\cos(x+1)\theta}.$$

Functional Equations.

97. If a function $\phi(x)$ is of such a nature that when it is twice performed on a quantity, the result is the quantity itself, that is $\phi\{\phi(x)\}$ or $\phi^2(x) = x$, then it is called a periodic function of the second order; and if $\phi(x)$ be such that $\phi^n(x) = x$, then $\phi(x)$ is termed a periodic function of the n^{th} order. Any symmetrical function of x and y , put equal to zero, will by resolution give for the value of y a function of x , which is a periodic function of the second order. Consequently,

$$\frac{a}{x}, \quad a-x, \quad \frac{x}{x-1}, \quad \frac{1-x}{1+x}, \quad \sqrt{1-x^2},$$

are all periodic functions of the second order. Examples of periodic functions of the third order are $\frac{1}{1-x}$, $\sqrt{1-x^2}$; and of the fourth order $\frac{1}{2} \frac{1}{1-x}$, $\frac{2}{2-x}$. Functional equations involving these expressions may be frequently solved by a simple elimination; as will be seen in the following instances.

Ex. 1. $\{f(x)\}^2 \cdot f\left(\frac{1-x}{1+x}\right) = c^2x;$

here, since $\frac{1-x}{1+x}$ is a periodic function of the second order,

replacing x by $\frac{1-x}{1+x}$, we get

$$\left\{f\left(\frac{1-x}{1+x}\right)\right\}^2 f(x) = c^2 \cdot \frac{1-x}{1+x};$$

hence dividing the square of the proposed equation by this, we have

$$\{f(x)\}^3 = \frac{1+x}{1-x} c^2 x^2, \text{ or } f(x) = \left(\frac{1+x}{1-x} c^2 x^2\right)^{\frac{1}{3}}.$$

Ex. 2. $f(x) + a \cdot f\left(\frac{1}{1-x}\right) = \frac{1}{x};$

here since $\frac{1}{1-x}$ is a periodic function of the third order, replacing x twice in succession by $\frac{1}{1-x}$, we get

$$f\left(\frac{1}{1-x}\right) + af\left(1 - \frac{1}{x}\right) = 1 - x,$$

$$f\left(1 - \frac{1}{x}\right) + af(x) = \frac{x}{x-1}.$$

Now multiply the second of these equations by $-a$, and the third by a^2 , and we find by adding them together,

$$(1 + a^3)f(x) = \frac{1}{x} + a(x-1) + \frac{a^2x}{x-1}.$$

98. Other functional equations may be sometimes solved by the aid of the Calculus of Finite Differences, or by particular artifices.

Ex. 1. $f(x) + f(a-x) = c^2.$

Assume $x = u_x, \quad a - x = u_{x+1};$

$$\therefore u_{x+1} + u_x = a, \quad \text{or } u_x = C(-1)^x + \frac{1}{2}a.$$

But $f(u_{x+1}) + f(u_x) = c^2$, which may be written

$$v_{x+1} + v_x = c^2;$$

$$\therefore v_x = C'(-1)^x + \frac{1}{2}c^2 = \frac{C'}{C}(u_x - \frac{1}{2}a) + \frac{1}{2}c^2 = f(u_x);$$

$$\therefore f(x) = m(x - \frac{1}{2}a) + \frac{1}{2}c^2,$$

m being an arbitrary constant.

In the same way we may find a particular solution of the equation

$$f(a+x) \cdot f(a-x) = a^2 - x^2;$$

or we may use for that purpose the following artifice,

$$\frac{f(a+x)}{a+x} \cdot \frac{f(a-x)}{a-x} = 1, \quad \text{or } \phi(x) \cdot \phi(-x) = 1,$$

which is satisfied by $\phi(x) = C^x$;

$$\therefore f(a+x) = (a+x) C^x = (a+x) C^{(a+x)-a};$$

$$\therefore f(x) = x C^{x-a}.$$

Ex. 2. $f(bx) = n f(x).$

Let $bx = u_{x+1}, \quad x = u_x;$

$$\therefore u_{x+1} - bu_x = 0, \quad \text{or} \quad u_x = \frac{1}{b} b^x = x;$$

$$\text{but } f(u_{x+1}) - n f(u_x) = 0, \quad \text{or } v_{x+1} - n v_x = 0;$$

$$\therefore v_x = n^x. \phi(\cos 2\pi x) = n^{\frac{\log cx}{\log b}} \times \phi \left\{ \cos \left(\frac{2\pi}{\log b} \log cx \right) \right\} \quad (\text{Art. 41}),$$

$$\text{but } v_x = f(u_x) = f(x),$$

$$\therefore f(x) = (cx)^{\frac{\log n}{\log b}} \times \phi \left\{ \cos \left(\frac{2\pi \log cx}{\log b} \right) \right\}.$$

Ex. 3. $f(a+bx) = n f(x).$

Assume $a+bx = u_{x+1}, \quad x = u_x$; then proceeding as in the last Example we have

$$f(x) = \left(cx - \frac{ca}{1-b} \right)^{\frac{\log n}{\log b}} \times \phi \left\{ \cos \frac{2\pi}{\log b} \log \left(cx - \frac{ca}{1-b} \right) \right\};$$

and for a particular value, when $c=1$, and $\phi(\cos 2\pi x) = 1$,

$$f(x) = \left(x - \frac{a}{1-b} \right)^{\frac{\log n}{\log b}}.$$

Ex. 4. $f(x) \cdot f(y) = f(x+y) + f(x-y).$

Expanding the second member by Taylor's theorem and dividing both sides by $f(x)$, we get

$$f(y) = 2 \left\{ 1 + \frac{1}{2} \frac{y^2}{f(x)} \cdot \frac{d^2 f(x)}{dx^2} + \frac{1}{1 \cdot 2 \cdot 3 \cdot 4} \cdot \frac{y^4}{f(x)} \cdot \frac{d^4 f(x)}{dx^4} + \&c. \right\}.$$

But as $f(y)$ does not involve x , the coefficients of y^2, y^4 , &c. in the second member must be constants; let b be the value of the coefficient of y^2 , then

$$\frac{d^2 f(x)}{dx^2} = b \cdot f(x),$$

$$\therefore \frac{d^4 f(x, y)}{dx^4} = b \cdot \frac{d^2 f(x)}{dx^2} = b^2 \cdot f(x), \text{ \&c.};$$

$$\therefore f(y) = 2 \left(1 + \frac{1}{2} b y^2 + \frac{1}{1 \cdot 2 \cdot 3 \cdot 4} b^2 y^4 + \text{\&c.} \right),$$

or replacing b by another constant $-c^2$,

$$f(y) = 2 \left(1 - \frac{1}{2} c^2 y^2 + \frac{1}{1 \cdot 2 \cdot 3 \cdot 4} c^4 y^4 - \text{\&c.} \right) = 2 \cos cy.$$

99. We shall conclude this Section by two or three Problems, the solution of which is effected by equations of Differences.

1. To find the value of $u_x = \sqrt{2 - \sqrt{2 - \sqrt{2} \dots}}$ to x terms.

Since $u_{x+1} = \sqrt{2 - u_x}$, $u_{x+1}^2 + u_x = 2$;

of which the integral is $u_x = 2 \cos \left\{ c \left(-\frac{1}{2} \right)^x + \frac{1}{3} \pi \right\}$; (Art. 86)

$$\therefore u_1 = 2 \cos \left(-\frac{c}{2} + \frac{\pi}{3} \right) = \sqrt{2} = 2 \cos \frac{\pi}{4}, \quad \therefore c = \frac{\pi}{6};$$

$$\therefore u_x = 2 \cos \left\{ \frac{\pi}{6} \left(-\frac{1}{2} \right)^x + \frac{\pi}{3} \right\} = 2 \sin \frac{\pi}{6} \{ 1 - \left(-\frac{1}{2} \right)^x \}.$$

2. A swan breeds three cygnets in its second year, and four every succeeding year; and the young ones all breed according to the same law; required the number at the end of the x^{th} year.

Let u_x = number at end of x^{th} year;
then in $(x+1)^{\text{th}}$ year, there will be bred three each by those in their second year, and four each by all the others;

$$\therefore u_{x+1} = u_x + 3(u_{x-1} - u_{x-2}) + 4u_{x-2} = u_x + 3u_{x-1} + u_{x-2};$$

$$\therefore u_{x+3} - u_{x+2} - 3u_{x+1} - u_x = 0;$$

$$\therefore m^3 - m^2 - 3m - 1 = m(m^2 - 1) - (m + 1)^2 = 0;$$

$$\therefore m = -1, \text{ and } 1 \pm \sqrt{2};$$

$$\therefore u_x = a(-1)^x + b(1 + \sqrt{2})^x + c(1 - \sqrt{2})^x,$$

$$u_0 = 0 = a + b + c,$$

$$u_1 = 1 = -a + b(1 + \sqrt{2}) + c(1 - \sqrt{2}),$$

$$u_2 = 4 = a + b(3 + 2\sqrt{2}) + c(3 - 2\sqrt{2});$$

$$\therefore a = 1, \quad b = \frac{1}{2\sqrt{2}}(3 - \sqrt{2}), \quad c = -\frac{1}{2\sqrt{2}}(3 + \sqrt{2});$$

$$\therefore u_x = (-1)^x + \frac{1}{2\sqrt{2}}(3 - \sqrt{2})(1 + \sqrt{2})^x - \frac{1}{2\sqrt{2}}(3 + \sqrt{2})(1 - \sqrt{2})^x.$$

If the law of increase be one in the second and every succeeding year,

$$\sqrt{5}u_x = \frac{1}{2^x}(1 + \sqrt{5})^x - \frac{1}{2^x}(1 - \sqrt{5})^x.$$

3. If a person possessed of a given capital of £ n , spends in the first year the whole of the interest receivable at the end of that year; and in the second year twice the interest receivable at the end of the second year; and in the third year thrice the interest receivable at the end of that year, and so on; to find how long it will be before his capital is exhausted.

Let u_x = reduced capital in pounds at the end of the x^{th} year, r = interest of £1 for a year; then ru_x = interest receivable at end of $(x + 1)^{\text{th}}$ year, and $(x + 1) ru_x$ = expenditure in $(x + 1)^{\text{th}}$ year, therefore reduced capital at end of $(x + 1)^{\text{th}}$ year, is

$$u_x + ru_x - (x + 1) ru_x = (1 - rx) u_x;$$

$$\therefore u_{x+1} - (1 - rx) u_x = 0;$$

$$\therefore u_x = n(1 - r)(1 - 2r) \dots \{1 - (x - 1)r\},$$

since when $x = 1$, $u_x = n$. Now this vanishes when $x = 1 + \frac{1}{r}$, which expresses the number of years the capital will last.

100. The following are instances of Geometrical Problems, which admit of being solved by Finite Differences.

1. To find a curve in which the chord QP joining the extremities of any two ordinates PN , QM at a given distance from one another, when produced shall meet the axis of the abscissæ at a constant distance from the foot of either ordinate.

Let $MN = h$, $TN = \frac{h}{n-1}$, $PN = y_x$, $QM = y_{x+h}$, (fig. 6)

$$\text{then } (y_{x+h} - y_x) \div h = y_x \div \frac{h}{n-1};$$

$$\therefore y_{x+h} - ny_x = 0; \quad \therefore y_x = Cn^{\frac{x}{h}};$$

or $y = Cn^{\frac{x}{h}}$ is the equation to the curve.

2. To find a curve such that the portion of a straight line drawn through a fixed point in its plane and terminated by the curve, shall be of a constant length. Let $\rho = f(\theta)$ be its equation, then we must have for all values of θ ,

$$f(\theta) + f(\theta + \pi) = c.$$

As π is here the increment of θ , let $\theta = \pi z$, and let $f(\pi z)$ be denoted by u_z , then $u_{z+1} + u_z = c$;

$$\therefore u_z = \frac{1}{2}c + C_z(-1)^z,$$

where C_z denotes any function of z that does not alter in value by the change of z into $z+1$. Now $(-1)^z = \cos \pi z$, and $F(\cos 2\pi z)$ fulfils the condition to which C_z is subject;

$$\therefore \rho = f(\theta) = \frac{1}{2}c + F(\cos 2\theta) \cdot \cos \theta,$$

the equation to the required curve. If $F(\cos 2\theta) = b$, this becomes $\rho = \frac{1}{2}c + b \cos \theta$; which represents a circle if $b = 0$.

3. To find a curve such that any n radii drawn through a fixed point in its plane so as to make equal angles with one

another, and terminating in the curve, shall have their sum invariable.

If $\rho = f(\theta)$ be the equation to the curve, and h denote the angle between two consecutive radii, we have

$$f(\theta) + f(\theta + h) + \dots + f\{\theta + (n-1)h\} = c \dots (1),$$

or making $\theta = hz$, and $f(hz) = u_z$,

$$u_{z+n-1} + u_{z+n-2} + \dots + u_z = c \dots \dots \dots (1),$$

the solution of which is

$$u_z = \frac{1}{n} c + C'_z \cos \frac{2\pi z}{n} + C''_z \cos \frac{4\pi z}{n} \\ + C'''_z \cos \frac{6\pi z}{n} + \&c. + C^{(n-1)}_z \cos \frac{(2n-2)\pi z}{n},$$

for it is seen by summing the resulting series that any one of the values $\cos \frac{2\pi z}{n}$, $\cos \frac{4\pi z}{n}$, &c. for u_z , satisfies (1), when $c = 0$.

Now suppose $h = \frac{2\pi}{m}$, then $\frac{2\pi z}{n} = \frac{mhz}{n} = \frac{m\theta}{n}$, and C'_z may be replaced by $f_1(\cos 2\pi z) = f_1(\cos m\theta)$, &c.; consequently the required equation to the curve is

$$\rho = \frac{1}{n} c + f_1(\cos m\theta) \cos \frac{m\theta}{n} + f_2(\cos m\theta) \cos \frac{2m\theta}{n} + \dots \\ + f_{n-1}(\cos m\theta) \cos \frac{(n-1)m\theta}{n}.$$

If $m = n$, $f_1(\cos m\theta) = b$, and if the rest of the arbitrary functions vanish, we get

$$\rho = \frac{1}{n} c + b \cos \theta.$$

4. To find a curve such that the product of the two segments of any straight line drawn through a fixed point in its plane to meet the curve, shall be constant. If $\rho = f(\theta)$ be the equation to the curve, then

$$f(\theta) \cdot f(\theta + \pi) = c^2; \text{ let } \theta = \pi z, \text{ and } f(\pi z) = u_z;$$

$$\therefore u_{z+1} \cdot u_z = c^2, \text{ which gives (Art. 86)}$$

$$u_z = c C_z^{(-1)^z} = c \cdot \{F(\cos 2\pi z)\}^{\cos \pi z},$$

since C_z is any function of z that does not alter in value when z is changed into $z + 1$, and $(-1)^z = \cos \pi z$;

$$\therefore \rho = c \{F(\cos 2\theta)\}^{\cos \theta}.$$

Now $F(\cos 2\theta)$ denotes any function of $\cos 2\theta$ whose value does not alter by the change of θ into $\pi + \theta$; and may therefore be replaced by

$$\left\{ \frac{F(\cos 2\theta) + \cos \theta}{F(\cos 2\theta) - \cos \theta} \right\}^{\frac{n}{\cos \theta}},$$

which evidently has the property of not being altered by the substitution of $\pi + \theta$ for θ ;

$$\therefore \frac{\rho}{c} = \left\{ \frac{F(\cos 2\theta) + \cos \theta}{F(\cos 2\theta) - \cos \theta} \right\}^n,$$

which gives algebraic curves by assigning an algebraic form to the arbitrary function F . Thus if $n = \frac{1}{2}$, and

$$F(\cos 2\theta) = \frac{\sqrt{a^2 - b^2 \sin^2 \theta}}{b}, \text{ then } \frac{\rho}{c} = \frac{\sqrt{a^2 - b^2 \sin^2 \theta} + b \cos \theta}{\sqrt{a^2 - b^2}},$$

the equation to a circle. Several other interesting Problems of this description may be found in Herschel's Examples.

The following example supplies an omission at the end of Art. 84*. It is evident that $v_x = a^x$ satisfies the equation

$$u_{x+2} - a(1 + B_x)u_{x+1} + a^2 B_x u_x = 0,$$

$$\therefore w_x = CB_x \text{ and } u_x = a^x \{C' + C \sum (PB_{x-1})\}.$$

$$\text{Suppose } B_x = \frac{x+1}{x+3}, \text{ then } PB_{x-1} = \frac{2 \cdot 3 \dots x}{4 \cdot 5 \dots (x+2)} = \frac{1}{(x+1)(x+2)};$$

$$\therefore \sum (PB_{x-1}) = \frac{C}{x+1}; \text{ and } u_x = a^x \left(C' + \frac{C}{x+1} \right) \text{ is the solution of}$$

$$(x+3)u_{x+2} - 2a(x+2)u_{x+1} + a^2(x+1)u_x = 0.$$

SECTION IV.

SUMMATION OF SERIES.

Integration of the General Term.

101. ONE of the most direct and important applications of the Calculus of Finite Differences, is the general method which it furnishes of assigning the sum of any number of Terms of a Series, of the general term of which we are able to take the integral. There will be two cases to consider, according as the general term is given explicitly in terms of the index, or is only given by means of an equation of differences. We shall begin with the former case, by shewing that the sum of any number of terms ending with the general term u_x , is equal to the integral of the following term, together with a constant.

Let S_x denote the sum of the first x terms of a series whose general term is u_x ,

$$\text{then } S_x = u_1 + u_2 + \dots + u_x,$$

$$S_{x+1} = u_1 + u_2 + \dots + u_x + u_{x+1};$$

$$\therefore S_{x+1} - S_x = \Delta S_x = u_{x+1};$$

$$\therefore S_x = \Sigma u_{x+1} + C.$$

$$\text{Making } x=0, \quad S_0 = 0 = \Sigma u_{x=1} + C;$$

or, as it is usually written, $0 = \Sigma u_1 + C$,

$$\therefore S_x = \Sigma u_{x+1} - \Sigma u_1.$$

In general the arbitrary constant will be determined by the term with which we make the series commence. Thus, if we

make it begin with u_r instead of u_1 , this amounts to supposing the sum of the terms preceding u_r , that is, Σu_r to be zero; and therefore to determine the constant, we have

$$S_{r-1} = \Sigma u_{x=r} + C = 0.$$

102. As the summation of series, where the general term is given explicitly as a function of the index, thus resolves itself into the cases of integration treated of in Section II., it will only be necessary to give a few numerical examples; every expression integrated in that Section gives the sum of a series of which it is the general term.

1. To find the sum of the first x terms of any progression of figurate numbers. In the r^{th} order the general term is

$$u_x = \frac{x(x+1)(x+2) \dots (x+r-2)}{1 \cdot 2 \cdot 3 \dots (r-1)};$$

$$\therefore \Sigma u_{x+1} = \frac{x(x+1) \dots (x+r-1)}{1 \cdot 2 \cdot 3 \dots r} + C, \quad (\text{Art. 45})$$

$$\Sigma u_1 = 0 + C = 0;$$

$$\therefore S_x = \frac{x(x+1) \dots (x+r-1)}{1 \cdot 2 \cdot 3 \dots r}.$$

Similarly, for the series of inverse figurate numbers, except when $r=2$, that is, for the series $1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots + \frac{1}{x}$.

$$\text{For we have } u_x = \frac{1 \cdot 2 \cdot 3 \dots (r-1)}{x(x+1) \dots (x+r-2)};$$

$$\therefore \Sigma u_{x+1} = C - \frac{1 \cdot 2 \cdot 3 \dots (r-1)}{(r-2)(x+1) \dots (x+r-2)}, \quad (\text{Art. 47})$$

$$\Sigma u_1 = C - \frac{r-1}{r-2} = 0,$$

$$S_x = (r-1) \left\{ \frac{1}{r-2} - \frac{1 \cdot 2 \cdot 3 \dots (r-3)}{(x+1)(x+2) \dots (x+r-2)} \right\},$$

$$\text{and } S_\infty = \frac{r-1}{r-2}.$$

2. To find the sums of the squares and cubes of the natural numbers. Here $u_x = x^2 = (x-1)x + x$;

$$\therefore \Sigma u_{x+1} = \frac{(x-1)x(x+1)}{3} + \frac{x(x+1)}{2} + C;$$

$$\text{and } \Sigma u_1 = 0 + C = 0; \therefore S_x = \frac{x(x+1)(2x+1)}{6}.$$

Again, $u_x = x^3 = x(x^2-1) + x$;

$$\therefore \Sigma u_{x+1} = \frac{(x-1)x(x+1)(x+2)}{4} + \frac{x(x+1)}{2} + C,$$

$$\Sigma u_1 = 0 + C = 0;$$

$$\therefore S_x = \frac{x(x+1)}{2} \left\{ \frac{(x-1)(x+2)}{2} + 1 \right\} = \left\{ \frac{x(x+1)}{2} \right\}^2.$$

Similarly, $1^2 + 3^2 + 5^2 + \dots + (2x-1)^2 = \frac{x(4x^2-1)}{3}.$

$$1^3 + 3^3 + 5^3 + \dots + (2x-1)^3 = 2x^4 - x^2.$$

3. To sum the series $1^2 - 2^2 + 3^2 - 4^2 + \dots \pm x^2$.

Here $u_x = (-1)^{x-1} x^2$;

$$\Sigma u_x = \frac{(-1)^{x-1} x^2}{-2} - \frac{(2x+1)(-1)^x}{4} + \frac{2(-1)^{x+1}}{-8} + C; \text{ (Art. 59)}$$

$$\therefore \Sigma u_{x+1} = (-1)^{x+1} \left\{ \frac{(x+1)^2}{2} - \frac{2x+3}{4} + \frac{1}{4} \right\} + C;$$

$$\therefore S_x = (-1)^{x+1} \frac{x(x+1)}{2}. \quad \text{Similarly,}$$

$$2 \cdot 1^2 + 2^2 \cdot 2^2 + 2^3 \cdot 3^2 + \dots + 2^x x^2 = 2^{x+1} (x+1) (x-3) + (2^{x+1}-1) 6.$$

4. To find the sum of x terms of the series

$$\frac{1}{2^2-3^2} + \frac{1}{4^2-3^2} + \frac{1}{6^2-3^2} + \&c.$$

$$u_x = \frac{1}{(2x)^2-3^2} = \frac{1}{(2x-3)(2x+3)}, \text{ which falls under Art. 48,}$$

$$S_x = \frac{1}{18} - \frac{x+1}{(2x+1)(2x+3)} - \frac{1}{6} \cdot \frac{6x+5}{(2x-1)(2x+1)(2x+3)};$$

$$\text{and } S_\infty = \frac{1}{18}. \quad \text{Similarly,}$$

$$\frac{5}{1 \cdot 2 \cdot 3} + \frac{8}{2 \cdot 3 \cdot 4} + \frac{11}{3 \cdot 4 \cdot 5} + \&c. \text{ to } x \text{ terms} = 2 - \frac{3x+4}{(x+1)(x+2)}.$$

$$5. \quad 2 \cdot 4 \cdot 7 + 4 \cdot 7 \cdot 13 + 8 \cdot 13 \cdot 25 + 16 \cdot 25 \cdot 49 + \&c. \text{ to } x \text{ terms.}$$

$$\text{Here } u_x = 2^x (3 \cdot 2^{x-1} + 1) (3 \cdot 2^x + 1).$$

$$\text{Assume } \Sigma u_{x+1} = A (3 \cdot 2^{x-1} + 1) (3 \cdot 2^x + 1) (3 \cdot 2^{x+1} + 1); \quad (\text{Art. 55})$$

$$\begin{aligned} \therefore u_{x+1} &= A (3 \cdot 2^x + 1) (3 \cdot 2^{x+1} + 1) \{3 \cdot 2^{x+2} + 1 - (3 \cdot 2^{x+1} + 1)\} \\ &= \frac{21}{4} A \cdot 2^{x+1} (3 \cdot 2^x + 1) (3 \cdot 2^{x+1} + 1); \text{ which gives } A = \frac{4}{21}, \end{aligned}$$

$$\therefore \Sigma u_{x+1} = \frac{4}{21} (3 \cdot 2^{x-1} + 1) (3 \cdot 2^x + 1) (3 \cdot 2^{x+1} + 1) + C;$$

$$\Sigma u_1 = \frac{4}{21} \cdot \frac{5}{2} \cdot 4 \cdot 7 + C = 0;$$

$$\therefore S_x = \frac{4}{21} (3 \cdot 2^{x-1} + 1) (3 \cdot 2^x + 1) (3 \cdot 2^{x+1} + 1) - \frac{40}{3}.$$

$$6. \quad \frac{2}{3 \cdot 3} - \frac{4}{3 \cdot 9} + \frac{8}{9 \cdot 15} - \frac{16}{15 \cdot 33} + \&c.$$

$$\text{Here } u_x = \frac{(-2)^x}{\{(-2)^x - 1\} \{(-2)^{x+1} - 1\}};$$

$$\therefore S_x = \frac{1}{9} + \frac{1}{3} \cdot \frac{1}{(-2)^{x+1} - 1}, \quad (\text{Art. 55}), \text{ and } S_\infty = \frac{1}{9}.$$

$$7. \quad \frac{16}{2 \cdot 3 \cdot 4} - \frac{21}{3 \cdot 4 \cdot 5} \cdot \frac{2}{3} + \frac{26}{4 \cdot 5 \cdot 6} \cdot \left(\frac{2}{3}\right)^2 - \&c. \text{ to } x \text{ terms,}$$

$$\text{Here } u_x = \frac{16 + 5(x-1)}{(x+1)(x+2)(x+3)} \left(-\frac{2}{3}\right)^{x-1}.$$

$$\text{Assume } \Sigma u_{x+1} = \frac{A}{(x+2)(x+3)} \left(-\frac{2}{3}\right)^x, \quad (\text{Art. 56})$$

$$\text{then } A = -3, \text{ and } \therefore S_x = \frac{1}{2} - \frac{3}{(x+2)(x+1)} \left(-\frac{2}{3}\right)^x.$$

Similarly for the sum of x terms of the series

$$\frac{19}{1 \cdot 2 \cdot 3} \cdot \frac{1}{4} + \frac{28}{2 \cdot 3 \cdot 4} \cdot \frac{1}{8} + \frac{39}{3 \cdot 4 \cdot 5} \cdot \frac{1}{16} + \frac{52}{4 \cdot 5 \cdot 6} \cdot \frac{1}{32} + \dots$$

$$u_x = \frac{x^2 + 6x + 12}{x(x+1)(x+2)} \left(\frac{1}{2}\right)^{x+1}, \quad S_x = 1 - \frac{x+4}{(x+1)(x+2)} \left(\frac{1}{2}\right)^{x+1}.$$

$$8. \quad \frac{1}{\cos \theta \cos 2\theta} + \frac{1}{\cos 2\theta \cos 3\theta} + \frac{1}{\cos 3\theta \cos 4\theta} + \&c.$$

to x terms.

$$\text{Here } u_x = \frac{1}{\cos x\theta \cos (x+1)\theta}; \text{ therefore } (\text{Art. 53})$$

$$\Sigma u_{x+1} = \frac{\tan (x+1)\theta}{\sin \theta} + C, \quad \Sigma u_1 = \frac{\tan \theta}{\sin \theta} + C = 0,$$

$$\therefore S_x = \frac{\tan(x+1)\theta - \tan\theta}{\sin\theta}.$$

9. $1 \cos \theta + 2 \cos 2\theta + 3 \cos 3\theta + \&c.$, to x terms.

Here $u_{x+1} = (x+1) \cos(x+1)\theta$,

$$\Sigma u_{x+1} = (x+1) \frac{\sin(x+\frac{1}{2})\theta}{2 \sin \frac{1}{2}\theta} - \frac{1}{2 \sin \frac{1}{2}\theta} \Sigma \sin(x+\frac{3}{2})\theta \quad (\text{Art. 59})$$

$$= \frac{(x+1) \sin(x+\frac{1}{2})\theta}{2 \sin \frac{1}{2}\theta} + \frac{\cos(x+1)\theta}{(2 \sin \frac{1}{2}\theta)^2} + C,$$

$$\Sigma u_1 = \frac{1}{2} + \frac{\cos\theta}{(2 \sin \frac{1}{2}\theta)^2} + C = 0, \quad \therefore C = -\frac{1}{(2 \sin \frac{1}{2}\theta)^2},$$

$$\begin{aligned} \therefore S_x &= \frac{x \sin(x+\frac{1}{2})\theta}{2 \sin \frac{1}{2}\theta} + \frac{1}{2 \sin^2 \frac{1}{2}\theta} \{\sin(x+\frac{1}{2})\theta \sin \frac{1}{2}\theta - \sin^2 \frac{1}{2}(x+1)\theta\} \\ &= \frac{x \sin(x+\frac{1}{2})\theta}{2 \sin \frac{1}{2}\theta} - \frac{1}{2} \left(\frac{\sin \frac{1}{2}x\theta}{\sin \frac{1}{2}\theta} \right)^2. \end{aligned}$$

Hence also, if $S_x = 1 \cos^2 \theta + 2 \cos^2 2\theta + 3 \cos^2 3\theta + \&c.$

then $2S_x = 1(1 + \cos 2\theta) + 2(1 + \cos 4\theta) + \&c.$

$$\therefore S_x = \frac{x(x+1)}{4} + \frac{x \sin(2x+1)\theta}{4 \sin \theta} - \frac{1}{4} \left(\frac{\sin x\theta}{\sin \theta} \right)^2.$$

10. $a \cos \theta + 2a^2 \cos 2\theta + 3a^3 \cos 3\theta + \&c.$, to x terms.

Here $u_x = xa^x \cos x\theta = xv_x$, suppose;

$$\therefore \Sigma u_x = x \Sigma v_x - \Sigma^2 v_{x+1}$$

$$= \frac{x}{c} (a^2 D^{-1} - 1) v_x - \frac{1}{c^2} (a^2 D^{-1} - 1)^2 v_{x+1}; \quad (\text{Art. 52})$$

$$\therefore \Sigma u_{x+1} = \frac{x+1}{c} (a^2 v_x - v_{x+1}) - \frac{1}{c^2} (a^4 v_x - 2a^2 v_{x+1} + v_{x+2}) + C,$$

$$0 = \frac{1}{c} (a^2 - a \cos \theta) - \frac{1}{c^2} (a^4 - 2a^3 \cos \theta + a^2 \cos 2\theta) + C;$$

$$\therefore C = \frac{a}{c} \cos \theta - 2 \left(\frac{a \sin \theta}{c} \right)^2;$$

$$\therefore S_x = \frac{x+1}{c} (a^3 v_x - v_{x+1}) - \frac{1}{c^2} (a^4 v_x - 2a^2 v_{x+1} + v_{x+2}) + \frac{a}{c} \cos \theta - 2 \frac{a^2}{c^2} \sin^2 \theta;$$

where $v_x = a^x \cos x\theta$. Similarly, if the general term be $x^2 a^x \cos x\theta$.

$$11. \quad \tan^{-1} \frac{1}{1+1+1^2} + \tan^{-1} \frac{1}{1+2+2^2} + \tan^{-1} \frac{1}{1+3+3^2} + \&c. \\ \text{to } x \text{ terms;}$$

$$u_x = \tan^{-1} \frac{1}{1+x+x^2} = \tan^{-1} \frac{\Delta x}{1+x(x+1)},$$

$$\therefore \Sigma u_x = \tan^{-1} x + C, \quad (\text{Art. 54})$$

$$\therefore S_x = \Sigma u_{x+1} - \Sigma u_1 = \tan^{-1} (x+1) - \frac{1}{4} \pi.$$

$$12. \quad \tan^{-1} \left(\frac{\theta^3}{4+3\theta^2} \right) + 2 \tan^{-1} \left(\frac{\theta^3}{32+6\theta^2} \right) + 4 \tan^{-1} \left(\frac{\theta^3}{256+12\theta^2} \right) + \&c.$$

$$\text{Here } u_{x+1} = 2^x \tan^{-1} \frac{\theta^3}{4 \cdot 2^{3x} + 3 \cdot 2^x \theta^2};$$

$$\therefore \Sigma u_{x+1} = 2^x \tan^{-1} \frac{\theta}{2^x} + C, \quad (\text{Art. 54})$$

$$\therefore S_x = \Sigma u_{x+1} - \Sigma u_1 = 2^x \tan^{-1} \frac{\theta}{2^x} - \tan^{-1} \theta, \text{ and } S_\infty = \theta - \tan^{-1} \theta.$$

And we may similarly shew for the series whose $(x+1)^{\text{th}}$ term is

$$3^x \tan^{-1} \left(\frac{8\theta^3 3^x}{27 \cdot 3^{4x} + 18\theta^2 \cdot 3^{2x} - \theta^4} \right),$$

$$\text{that } S_x = 3^x \tan^{-1} \frac{\theta}{3^x} - \tan^{-1} \theta, \text{ and } S_\infty = \theta - \tan^{-1} \theta.$$

$$13. \quad \log \cot \theta + \frac{1}{2} \log \cot 2\theta + \frac{1}{4} \log \cot 4\theta + \&c. \text{ to } x \text{ terms.}$$

$$\text{Here } u_{x+1} = \frac{1}{2^x} \log \cot 2^x \theta = 2\Delta \frac{1}{2^x} \log (2 \sin 2^x \theta);$$

$$\therefore S_x = \frac{1}{2^{x-1}} \log (2 \sin 2^x \theta) - 2 \log (2 \sin \theta).$$

Recurring Series.

103. We next come to the case where the general term is not given explicitly in terms of its index, but only certain relations between the consecutive terms, or these and their indices are expressed.

The equation of differences which determines the form of the general term, cannot always be solved; when however the equation is linear with constant coefficients, its solution, as we know, can be effected; and this happens for Recurring Series—the most remarkable class of this sort of series.

This equation when integrated will involve the same number of arbitrary constants as is expressed by its order: implying that that number of consecutive terms may be regarded as indeterminate, and the remaining terms formed according to the law which the equation to the series expresses.

A recurring series is a series in which an equation of the first degree with constant coefficients, holds good between a certain definite number of consecutive terms, in whatever part of the series they be taken. For example, in the series

$$3 + 5 + 9 + 17 + 33 + \&c.$$

we have $9 = 3 \cdot 5 - 2 \cdot 3$, $17 = 3 \cdot 9 - 2 \cdot 5$, &c.; and in general

$$u_{x+2} = 3u_{x+1} - 2u_x. \quad (1)$$

The integral of this is $u_x = c \cdot 2^x + c'$; and assuming a and b for the first two terms of the series of which u_x is the general term,

$$a = 2c + c', \quad b = 4c + c'; \quad \therefore c = \frac{1}{2}(b - a), \quad c' = 2a - b,$$

$$\text{so that } u_x = (b - a) 2^{x-1} + 2a - b;$$

and the series that has (1) for its equation is, in its most general form, $a + b + (3b - 2a) + (7b - 6a) + \&c.$

But for the series $\frac{a}{n+1} + \frac{a^2}{n+2} + \frac{a^3}{n+3} + \&c.$ we have

$$(x + n + 2) u_{x+2} - 2a(x + n + 1) u_{x+1} + a^2(x + n) u_x = 0,$$

an equation with variable coefficients, giving $u_x = \frac{a^x(c + c'x)}{x + n}.$

104. The general equation of every recurring series is

$$u_{x+n} + p_1 u_{x+n-1} + p_2 u_{x+n-2} + \dots + p_{n-1} u_{x+1} + p_n u_x = 0.$$

The series of coefficients which connects any term with the preceding ones is called the Scale of Relation. Thus

$$1 + p_1 + p_2 + \dots + p_n = 0$$

is the scale of relation of the recurring series whose equation is

$$u_{x+n} + p_1 u_{x+n-1} + \dots + p_n u_x = 0 \dots\dots\dots (1).$$

105. A recurring series may generally be resolved into two or more geometric progressions.

For if $a_1, a_2, a_3, \dots a_n$ be the roots of the equation

$$a^n + p_1 a^{n-1} + p_2 a^{n-2} + \dots + p_n = 0 \dots\dots\dots (2),$$

the complete integral of the equation of the series is

$$u_x = c_1 a_1^x + c_2 a_2^x + \dots + c_n a_n^x, \text{ (Art. 77)}$$

$$\therefore u_1 = c_1 a_1 + c_2 a_2 + c_3 a_3 + \dots + c_n a_n,$$

$$u_2 = c_1 a_1^2 + c_2 a_2^2 + c_3 a_3^2 + \dots + c_n a_n^2,$$

$$\dots\dots\dots$$

$$u_n = c_1 a_1^n + c_2 a_2^n + c_3 a_3^n + \dots c_n a_n^n,$$

and the proposed series consequently is transformed into

$$c_1 (a_1 + a_1^2 + \dots + a_1^n) + c_2 (a_2 + a_2^2 + \dots + a_2^n) + \&c. \\ + c_n (a_n + a_n^2 + \dots a_n^n).$$

106. In the particular cases in which equation (2) has equal or impossible roots, the recurring series can no longer be resolved into geometric progressions; for the complete integral of the equation of the series becomes in those two cases, respectively,

$$u_x = (c_0 + c_1 x + c_2 x^2 + \dots + c_{r-1} x^{r-1}) a_1^x + c_{r+1} a_{r+1}^x + \dots + c_n a_n^x,$$

$$u_x = (c_0 + c_1 x + \dots + c_{r-1} x^{r-1}) \rho^x \cos x\theta$$

$$+ (c'_0 + c'_1 x + \dots + c'_{r-1} x^{r-1}) \rho^x \sin x\theta + c_{2r+1} a_{2r+1}^x + \dots + c_n a_n^x.$$

107. Hence, to find the general term of a recurring series, we must integrate the equation expressing the relation between its successive terms, and determine the arbitrary constants by making the general term u_x coincide with a sufficient number of given terms. When a series is known to be recurring, its equation may be determined by assuming it to be of the form (1); and then forming a sufficient number of equations for finding the coefficients p_1, p_2 , &c. p_n , by substituting the given terms in order.

108. To find the sum of x terms of a recurring series.

First, let its general term be of the form

$$u_x = c_1 a_1^x + c_2 a_2^x + \dots + c_n a_n^x; \text{ then } S_x = \sum u_{x+1} - \sum u_1 \\ = c_1 \cdot \frac{a_1^{x+1} - a_1}{a_1 - 1} + c_2 \frac{a_2^{x+1} - a_2}{a_2 - 1} + \dots + c_n \cdot \frac{a_n^{x+1} - a_n}{a_n - 1}.$$

Secondly, let the general term be

$$u_x = (c_0 + c_1 x + c_2 x^2 + \dots + c_{r-1} x^{r-1}) a_1^x + c_{r+1} a^{x+r+1} + \&c.;$$

then since

$$\sum x^n a^x = \frac{a^x x^n}{a-1} - \frac{a^{x+1} \Delta x^n}{(a-1)^2} + \frac{a^{x+2} \Delta^2 x^n}{(a-1)^3} - \&c. \text{ (Art. 59),}$$

$$\sum u_{x+1} = c_0 \frac{a_1^{x+1}}{a_1 - 1} + c_1 \left\{ \frac{a_1^{x+1} (x+1)}{a_1 - 1} - \frac{a_1^{x+2}}{(a_1 - 1)^2} \right\} + \&c.;$$

$$\therefore S_x = \sum u_{x+1} - \sum u_1, \text{ is determined.}$$

Thirdly, let

$$u_x = \rho^x (c_0 \cos x\theta + c_0' \sin x\theta) + \rho^x x (c_1 \cos x\theta + c_1' \sin x\theta) \\ + \rho^x x^2 (c_2 \cos x\theta + c_2' \sin x\theta) + \&c.,$$

then each term to be integrated will be of the form

$$x^n \rho^x \cos (x\theta + \alpha),$$

the integral of which may be found by Art. 58, because

$$\sum^n \rho^2 \cos (x\theta + \alpha)$$

is always assignable.

Ex. 1. To find the sum of x terms of the series

$$1 + 5 + 17 + 53 + \&c.$$

Let the equation be $u_{x+2} + pu_{x+1} + qu_x = 0$;

then $17 + 5p + q = 0$, $53 + 17p + 5q = 0$, which give $p = -4$,

$$q = 3, \text{ so that } u_{x+2} - 4u_{x+1} + 3u_x = 0.$$

Let $u_x = a^x$; then $a^2 - 4a + 3 = 0$; $a = 3$ or 1 ;

$$\therefore u_x = c_1 3^x + c_2, \quad 1 = 3c_1 + c_2, \quad 5 = 9c_1 + c_2,$$

which give $c_1 = \frac{2}{3}$, $c_2 = -1$, so that $u_x = 2 \cdot 3^{x-1} - 1$;

$$\therefore S_x = 2 \cdot \frac{3^x}{3-1} - x + C, \quad 0 = 1 + C;$$

$$\therefore S_x = 3^x - x - 1.$$

Ex. 2. $2 - a - a^2 + 2a^3 - a^4 - a^5 + 2a^6 - a^7 - \&c.$

$$u_x = 2a^{x-1} \cos \frac{(2x-2)\pi}{3};$$

$$S_x = \frac{2a^{x+1} \cos \frac{1}{3}(2x-2)\pi - 2a^2 \cos \frac{2}{3}\pi x + a + 2}{a^2 + a + 1}.$$

Ex. 3. $1 + 2 + 3 + 8 + 13 + 30 + 55 + 116 + \&c.$

$$u_{x+3} - 3u_{x+1} - 2u_x = 0, \quad u_x = \frac{4}{9} 2^x + \frac{1}{9} (3x-4)(-1)^x.$$

$$S_x = \frac{8}{9} 2^x - \frac{3}{4} + (-1)^x \cdot \frac{6x-5}{36}.$$

Ex. 4. $1 + 4 + 18 + 80 + 356 + \&c.$, to x terms.

Here $u_{x+2} - 4u_{x+1} - 2u_x = 0$;

$\therefore 2\sqrt{6}u_x = (2 + \sqrt{6})^x - (2 - \sqrt{6})^x$; and

$$2\sqrt{6}S_x = \frac{(2 + \sqrt{6})^{x+1} - (2 + \sqrt{6})}{1 + \sqrt{6}} - \frac{(2 - \sqrt{6})^{x+1} - (2 - \sqrt{6})}{1 - \sqrt{6}}.$$

Application of the Integral Calculus to the Summation of Series.

109. As integrals are often expressed by Series, so, conversely, the latter may be represented by integrals; and it is often desirable to find the integral of which a proposed series is one of the developments, in order to subject it to the methods which we possess for calculating, at least approximately, the value of any integral taken between assigned limits. We proceed therefore to notice one or two processes given by Euler for effecting this; they consist chiefly in performing certain operations on the series, by which it is transformed into another series which we are able to sum, or which is similar to the proposed one.

110. Series which proceed according to the powers of some quantity t , affected with coefficients consisting of factors in arithmetic progression either in the numerator or denominator, may be summed by the aid of the Integral Calculus, the denominators being taken away by differentiation and the numerators by integration.

Ex. 1. Let $s = \frac{t}{2} + \frac{2t^2}{3} + \frac{3t^3}{4} + \&c.$ (to ∞),

$$\therefore \frac{d}{dt}(st) = t + 2t^2 + 3t^3 + \&c.$$

$$\therefore \int dt \cdot \frac{1}{t} \frac{d}{dt}(st) = t + t^2 + t^3 + \&c. = \frac{t}{1-t};$$

$$\therefore \frac{1}{t} \frac{d}{dt}(st) = \frac{1}{(1-t)^2}, \quad st = \int dt \frac{t}{(1-t)^2} = \log(1-t) + \frac{1}{1-t} + C,$$

$$\therefore s = \frac{\log(1-t)}{t} + \frac{1}{1-t}, \text{ since } C = -1.$$

Ex. 2. $s = \frac{a+b}{\alpha+\beta} t + \frac{a+2b}{\alpha+2\beta} t^2 + \dots \frac{a+nb}{\alpha+n\beta} t^n + \dots$ to m terms;

$$\therefore \beta \frac{d}{dt} (st^{\frac{\alpha}{\beta}}) = \dots + (a+nb) t^{n+\frac{\alpha}{\beta}-1} + \dots$$

$$\frac{\beta}{b} \int dt \left\{ t^{\frac{\alpha}{\beta}-\frac{\alpha}{\beta}} \frac{d}{dt} (st^{\frac{\alpha}{\beta}}) \right\} = \dots + t^{n+\frac{\alpha}{b}} + \dots = \frac{t^m-1}{t-1} t^{\frac{\alpha}{b}+1};$$

$$\therefore s = \frac{b}{\beta} t^{-\frac{\alpha}{\beta}} \int dt t^{\frac{\alpha}{\beta}-\frac{\alpha}{b}} \cdot \frac{d}{dt} \left\{ \frac{t^m-1}{t-1} \cdot t^{\frac{\alpha}{b}+1} \right\}.$$

111. Series like the above in which the coefficient of every term is the same function of the index of that term, may be readily summed by separation of symbols. Thus if we put $t=e^\theta$, and denote $\frac{d}{d\theta}$ by d , so that $f(d) e^{m\theta} = f(m) e^{m\theta}$, the series in Ex. 2, taken to infinity, may be written

$$\begin{aligned} s &= \frac{a+b\bar{d}}{\alpha+\beta\bar{d}} (e^\theta + e^{2\theta} + e^{3\theta} + \dots) = \frac{a+b\bar{d}}{\alpha+\beta\bar{d}} \frac{e^\theta}{1-e^\theta} \\ &= \left\{ \frac{a\beta-b\alpha}{\beta} (\alpha+\beta d)^{-1} + \frac{b}{\beta} \right\} \frac{e^\theta}{1-e^\theta} \\ &= \frac{a\beta-b\alpha}{\beta^2} e^{-\frac{a\theta}{\beta}} \int d\theta \frac{e^{\left(\frac{\alpha}{\beta}+1\right)\theta}}{1-e^\theta} + \frac{b}{\beta} \cdot \frac{e^\theta}{1-e^\theta} \\ &= \frac{a\beta-b\alpha}{\beta^2} t^{-\frac{\alpha}{\beta}} \int dt \frac{t^{\frac{\alpha}{\beta}}}{1-t} + \frac{b}{\beta} \cdot \frac{t}{1-t}, \end{aligned}$$

which may be easily shewn to coincide with the preceding result when m is infinite. And in general the series

$$f(n) t^n + f(n+r) t^{n+r} + f(n+2r) t^{n+2r} + \&c. \text{ to } m \text{ terms,}$$

becomes by putting $t=e^\theta$, $\frac{d}{d\theta} = d$,

$$f(d) (e^{n\theta} + e^{(n+r)\theta} + e^{(n+2r)\theta} + \&c.) = f(d) e^{n\theta} \cdot \frac{1-e^{mr\theta}}{1-e^{r\theta}}.$$

Thus the infinite series

$$\begin{aligned}
 & \frac{5t^3}{1 \cdot 2 \cdot 3} + \frac{6t^4}{2 \cdot 3 \cdot 4} + \frac{7t^5}{3 \cdot 4 \cdot 5} + \&c. \text{ becomes} \\
 & \frac{d+2}{d(d-1)(d-2)} (e^{3\theta} + e^{4\theta} + e^{5\theta} + \&c.) = \frac{d+2}{d(d-1)(d-2)} \frac{e^{3\theta}}{1-e^\theta} \\
 & = \left(\frac{1}{d} - \frac{3}{d-1} + \frac{2}{d-2} \right) \frac{e^{3\theta}}{1-e^\theta} \\
 & = \int d\theta \frac{e^{3\theta}}{1-e^\theta} - 3e^\theta \int d\theta \frac{e^{2\theta}}{1-e^\theta} + 2e^{2\theta} \int d\theta \frac{e^\theta}{1-e^\theta} \\
 & = \int dt \frac{t^2}{1-t} - 3t \int dt \frac{t}{1-t} + 2t^2 \int dt \frac{1}{1-t} \\
 & = \frac{5t^2}{2} - t - (1-3t+2t^2) \log(1-t).
 \end{aligned}$$

112. In the following instances the coefficient of any term of the series is not a function of the index of an invariable form.

$$\begin{aligned}
 \text{Ex. 1. } s &= 1 + \frac{a}{\alpha+\beta} t + \frac{a(a+b)}{(\alpha+\beta)(\alpha+2\beta)} t^2 \\
 &+ \frac{a(a+b)(a+2b)}{(\alpha+\beta)(\alpha+2\beta)(\alpha+3\beta)} t^3 + \&c. \text{ (to } \infty),
 \end{aligned}$$

$$\begin{aligned}
 \frac{1}{b} \int dt s t^{\frac{a}{b}-2} &= \frac{t^{\frac{a}{b}-1}}{a-b} + \frac{t^{\frac{a}{b}}}{\alpha+\beta} + \frac{a}{(\alpha+\beta)(\alpha+2\beta)} t^{\frac{a}{b}+1} \\
 &+ \frac{a(a+b)}{(\alpha+\beta)(\alpha+2\beta)(\alpha+3\beta)} t^{\frac{a}{b}+2} + \&c.;
 \end{aligned}$$

$$\begin{aligned}
 \therefore \frac{\beta}{b} \frac{d}{dt} \left(t^{\frac{a}{\beta}-\frac{a}{b}+1} \cdot \int dt s t^{\frac{a}{b}-2} \right) &= \frac{a}{a-b} t^{\frac{a}{\beta}-1} + t^{\frac{a}{\beta}} + \frac{a}{\alpha+\beta} t^{\frac{a}{\beta}+1} \\
 &+ \frac{a(a+b)}{(\alpha+\beta)(\alpha+2\beta)} t^{\frac{a}{\beta}+2} + \&c. = \frac{a}{a-b} t^{\frac{a}{\beta}-1} + t^{\frac{a}{\beta}} s; \dots (1)
 \end{aligned}$$

hence, performing the differentiation, we find

$$(\beta t - b t^2) \frac{ds}{dt} + (\alpha - at) s = \alpha,$$

a linear equation of the first order for finding s .

For the sum of m terms, we shall evidently have from equation (1)

$$\frac{\beta}{b} \frac{d}{dt} \left(t^{\frac{\alpha}{\beta} - \frac{\alpha}{b} + 1} \int dt s t^{\frac{\alpha}{b} - 2} \right) = \frac{\alpha}{a-b} t^{\frac{\alpha}{\beta} - 1} \\ + t^{\frac{\alpha}{\beta}} \left\{ s - \frac{a(a+b) \dots \{a + (m-2)b\}}{(\alpha + \beta)(\alpha + 2\beta) \dots \{\alpha + (m-1)\beta\}} t^{m-1} \right\}.$$

$$\text{Ex. 2. } \frac{t^a}{a} + \frac{t^{a+b}}{a(a+b)} + \frac{t^{a+2b}}{a(a+b)(a+2b)} + \&c. \text{ (to } \infty \text{)}.$$

$$s = e^{\frac{1}{b}t} \left(c + \int dt e^{-\frac{1}{b}t} \cdot t^{a-1} \right).$$

If we suppose $a = b = 3$, we get $s = e^{\frac{1}{3}t^2} - 1$.

$$\text{Hence } \frac{1}{3} + \frac{1}{3 \cdot 6} + \frac{1}{3 \cdot 6 \cdot 9} + \&c. = \sqrt[3]{e} - 1.$$

$$\text{Ex. 3. } s = 1 + \frac{\alpha}{1} \cdot \frac{\beta}{\gamma} t + \frac{\alpha(\alpha+1)}{1 \cdot 2} \frac{\beta(\beta+1)}{\gamma(\gamma+1)} t^2 \\ + \frac{\alpha(\alpha+1)(\alpha+2)}{1 \cdot 2 \cdot 3} \frac{\beta(\beta+1)(\beta+2)}{\gamma(\gamma+1)(\gamma+2)} t^3 + \&c. \text{ (to } \infty \text{)}.$$

$$(t^2 - t) \frac{d^2 s}{dt^2} + \{(\alpha + \beta + 1)t - \gamma\} \frac{ds}{dt} + \alpha \beta s = 0,$$

which is satisfied by $s = (1 - t)^{-\alpha}$, when $\beta = \gamma$.

113. When consecutive denominators have only a single factor in common, or none at all, the following is a convenient mode of summing the series.

$$\text{Ex. } \frac{1}{a(a+b)(a+2b)} + \frac{1}{(a+2b)(a+3b)(a+4b)} \\ + \&c. \text{ (to } \infty \text{)}.$$

$$\text{Since } \frac{2b^2}{(a+xb)\{a+(x+1)b\}\{a+(x+2)b\}} = \frac{1}{a+xb} \\ - \frac{2}{a+(x+1)b} + \frac{1}{a+(x+2)b},$$

resolving all the terms by means of this formula, by putting $x=0, 2, 4, \&c.$, we find

$$2b^2S = \frac{1}{a} - \frac{2}{a+b} + \frac{2}{a+2b} - \frac{2}{a+3b} + \frac{2}{a+4b} - \&c. \\ \therefore 2b^2S = 2 \int_0^1 dx \frac{x^{a-1}}{1+x^b} - \frac{1}{a}.$$

$$\text{Similarly, } \frac{1}{a(a+b)(a+2b)(a+3b)} \\ + \frac{1}{(a+3b)(a+4b)(a+5b)(a+6b)} + \&c. \text{ (to } \infty) \\ = \frac{1}{6b^3} \left(\frac{1}{a} - 3 \int_0^1 dx \frac{x^{a+b-1}}{1+x^b+x^{2b}} \right).$$

114. We may obtain an expression for the sum of the series

$$1 + \frac{x}{1} \cdot \frac{d\phi(z)}{dz} + \frac{x^2}{1 \cdot 2} \frac{d^2\{\phi(z)\}^2}{dz^2} + \frac{x^3}{1 \cdot 2 \cdot 3} \frac{d^3\{\phi(z)\}^3}{dz^3} + \&c.,$$

where $\phi(z)$ is any function of z , by Lagrange's Theorem.

For if $y = z + x\phi(y)$, then

$$y = z + \frac{x}{1} \phi(z) + \frac{x^2}{1 \cdot 2} \frac{d\{\phi(z)\}^2}{dz} + \&c.$$

$$\therefore 1 + \frac{x}{1} \frac{d\phi(z)}{dz} + \frac{x^2}{1 \cdot 2} \frac{d^2\{\phi(z)\}^2}{dz^2} + \&c. = \frac{dy}{dz} = \frac{1}{1 - x \frac{d}{dy} \phi(y)},$$

where $\frac{d}{dy} \phi(y)$ must be obtained in terms of x and z , from the equation $y = z + x\phi(y)$. Suppose, for example,

$$\phi(y) = \frac{1}{2}(y^2 - 1), \text{ then } \frac{d\phi(y)}{dy} = y,$$

$$\text{and } y = z + \frac{x}{2}(y^2 - 1), \quad \therefore 1 - xy = \sqrt{1 - 2zx + x^2};$$

$$\therefore 1 + \frac{x}{1} \cdot \frac{1}{2} \frac{d(z^2 - 1)}{dz} + \frac{x^2}{1 \cdot 2} \cdot \frac{1}{2^2} \cdot \frac{d^2(z^2 - 1)^2}{dz^2} + \dots = \frac{1}{\sqrt{1 - 2zx + x^2}}.$$

Hence if

$$(1 - 2zx + x^2)^{-\frac{1}{2}} = 1 + Z_1 x + Z_2 x^2 + \dots + Z_n x^n + \dots,$$

$$\text{then } Z_n = \frac{1}{2^n} \frac{d^n(z^2 - 1)^n}{dz^n}.$$

115. The function Z_n has some remarkable properties, of which the following may be noticed.

From the formula

$$\int dx \frac{1}{\sqrt{(a + bx)(c + ex)}} = \frac{2}{\sqrt{eb}} \log \left\{ \sqrt{a + bx} + \sqrt{\frac{b}{e}(c + ex)} \right\}$$

it is easily shewn that

$$\int_{-1}^{+1} dz (1 - 2arz + a^2 r^2)^{-\frac{1}{2}} \cdot \left(1 - \frac{2a}{r} z + \frac{a^2}{r^2}\right)^{-\frac{1}{2}} = \frac{1}{a} \log \left(\frac{1+a}{1-a}\right),$$

$$\begin{aligned} \therefore \int_{-1}^{+1} dz (1 + Z_1 ar + Z_2 a^2 r^2 + \dots) \cdot \left(1 + Z_1 \frac{a}{r} + Z_2 \frac{a^2}{r^2} + \dots\right) \\ = \frac{2}{a} \left(a + \frac{a^3}{3} + \frac{a^5}{5} + \&c.\right). \end{aligned}$$

Hence the definite integral of every term of the product forming the first member that is not independent of r , must equal zero;

$$\therefore \int_{-1}^{+1} dz (Z_n Z_m) = 0, \text{ and } \int_{-1}^{+1} dz (Z_n)^2 = \frac{2}{2n+1}.$$

116. To find an expression for the sum of the reciprocals of the n^{th} powers of the values of y , in the equation

$$z - y + \phi(y) = 0;$$

where $\phi(y)$ denotes any function of y , and z a quantity independent of y .

Let $z - y + \phi(y) = C(a_1 - y)(a_2 - y) \dots (a_m - y)$,

$$\therefore \frac{1 - \frac{d}{dy} \phi(y)}{z - y + \phi(y)} = \frac{1}{a_1 - y} + \frac{1}{a_2 - y} + \dots + \frac{1}{a_m - y} \dots \dots (1)$$

Now developing the two members of this equation in powers of y , the general term of the second member is

$$\left(\frac{1}{a_1^{n+1}} + \frac{1}{a_2^{n+1}} + \dots + \frac{1}{a_m^{n+1}} \right) y^n = s_{n-1} \cdot y^n,$$

and by Taylor's theorem the general term of the first member, putting $1 - \frac{d}{dy} \phi(y) = f(y)$, and regarding $f(y) \div (z - y)$ as a function of z and $\phi(y)$ as the increment of z , is

$$\frac{f(y) \phi^r(y)}{[r]} \frac{d^r}{dz^r} \left(\frac{1}{z - y} \right).$$

But if $f(y) = A_0 + A_1 y + \dots + A_n y^n + \dots$,

$$\text{since } \frac{1}{z - y} = \frac{1}{z} + \frac{y}{z^2} + \dots + \frac{y^n}{z^{n+1}} + \dots,$$

$$\therefore \frac{f(y)}{z - y} = \dots + \frac{A_0 + A_1 z + \dots + A_n z^n}{z^{n+1}} y^n + \dots,$$

therefore the general term of $\frac{f(y)}{z - y}$ is $\frac{f(z)}{z^{n+1}} y^n$, whatever be the form of $f(z)$, if $\frac{f(z)}{z^{n+1}}$ be restricted to such terms only as involve negative powers of z ; consequently the general term of $\frac{f(y) \phi^r(y)}{z - y}$ is $\frac{f(z) \phi^r(z)}{z^{n+1}} \cdot y^n$, and therefore of

$$\frac{1}{[r]} \frac{d^r}{dz^r} \left\{ \frac{f(y) \phi^r(y)}{z - y} \right\} \text{ it is } \frac{1}{[r]} \frac{d^r}{dz^r} \left\{ \frac{f(z) \phi^r(z)}{z^{n+1}} \right\} y^n;$$

hence equating coefficients of y^n on both sides of equation (1),

$$s_{n-1} = \frac{f(z)}{z^{n+1}} + \frac{d}{dz} \left\{ \frac{f(z) \phi(z)}{z^{n+1}} \right\} + \frac{1}{1 \cdot 2} \frac{d^2}{dz^2} \left\{ \frac{f(z) \phi^2(z)}{z^{n+1}} \right\} + \&c.$$

Now putting for $f(z)$ its value, the general term may be resolved into

$$\frac{1}{\lfloor r} \frac{d^r}{dz^r} \left\{ \frac{\phi^r(z)}{z^{n+1}} \right\} - \frac{1}{\lfloor r} \frac{d^r}{dz^r} \left\{ \frac{\phi'(z) \cdot \phi^r(z)}{z^{n+1}} \right\},$$

or

$$\frac{1}{\lfloor r-1} \frac{d^{r-1}}{dz^{r-1}} \left\{ \frac{\phi^{r-1}(z) \phi'(z)}{z^{n+1}} \right\} - \frac{n+1}{\lfloor r} \frac{d^{r-1}}{dz^{r-1}} \left\{ \frac{\phi^r(z)}{z^{n+1}} \right\}$$

$$- \frac{1}{\lfloor r} \frac{d^r}{dz^r} \left\{ \frac{\phi^r(z) \phi'(z)}{z^{n+1}} \right\},$$

of which the first and last will be destroyed by the corresponding parts in the preceding and succeeding terms; therefore, changing n into $n-1$,

$$s_{-n} = \frac{1}{z^n} - \frac{n}{1} \frac{\phi(z)}{z^{n+1}} - \frac{n}{1.2} \frac{d}{dz} \left\{ \frac{\phi^2(z)}{z^{n+1}} \right\} - \dots - \frac{n}{\lfloor r} \frac{d^{r-1}}{dz^{r-1}} \left\{ \frac{\phi^r(z)}{z^{n+1}} \right\} - \&c.$$

But the second member of this equation is what Lagrange's Theorem gives for the development of y^{-n} , in the equation $z - y + \phi(y) = 0$; therefore s_{-n} is equal to the sum of those terms of the development of y^{-n} which involve negative powers of z .

117. By the help of the preceding Proposition it may be shewn that Lagrange's Theorem, applied to the solution of the equation

$$z - y + \phi(y) = 0,$$

gives an approximation to the least root. By what precedes, we have

$$\frac{s_{-n}}{s_{-n-r}} = \frac{\frac{1}{z^n} - \frac{n}{1} \frac{\phi(z)}{z^{n+1}} - \frac{n}{1.2} \frac{d}{dz} \left\{ \frac{\phi^2(z)}{z^{n+1}} \right\} - \&c.}{\frac{1}{z^{n+r}} - \frac{n+r}{1} \frac{\phi(z)}{z^{n+r+1}} - \frac{n+r}{1.2} \frac{d}{dz} \left\{ \frac{\phi^2(z)}{z^{n+r+1}} \right\} - \&c.},$$

where each series is restricted to those terms which involve negative powers of z ; and the number of those terms increases with n , and if n be supposed very great, then each series may

be taken ad infinitum. But whatever be the value of n , if both series go on ad infinitum, the value of the second member is

$$\frac{y^{-n}}{y^{-n-r}} = y^r = z^r + \frac{r}{1} z^{r-1} \phi(z) + \frac{r}{1 \cdot 2} \frac{d}{dz} \{z^{r-1} \phi^2(z)\} + \&c.,$$

as given by Lagrange's Theorem, and which may be verified by actual multiplication; therefore when n is infinite,

$$\text{limit of } \frac{s_{-n}}{s_{-n-r}} = y^r,$$

(y^r denoting that function of z which is given for the development of y^r by Lagrange's Theorem).

But if a_1 be the least of the roots $a_1, a_2, a_3, \&c.$, we have also (Theory of Algebraical Equations, Art. 156)

$$\text{limit of } \frac{s_{-n}}{s_{-n-r}} = a_1^r, \text{ when } n \text{ is infinite;}$$

$$\therefore y^r = a_1^r, \text{ and } y = a_1 \text{ the least root.}$$

118. To find the sums of the series

$$\frac{\sin \alpha}{1^2 + k^2} + \frac{2 \sin 2\alpha}{2^2 + k^2} + \frac{3 \sin 3\alpha}{3^2 + k^2} + \&c. \text{ (to } \infty \text{),}$$

$$\frac{\cos \alpha}{1^2 + k^2} + \frac{\cos 2\alpha}{2^2 + k^2} + \frac{\cos 3\alpha}{3^2 + k^2} + \&c. \text{ (to } \infty \text{),}$$

$$\text{or the values of } {}^1S^\infty \frac{n \sin n\alpha}{n^2 + k^2}, \quad {}^1S^\infty \frac{\cos n\alpha}{n^2 + k^2}.$$

First, we have (*Trigon.* Art. 155)

$$\frac{(1-p^2) e^{-kx}}{1-2p \cos x + p^2} = e^{-kx} + 2(p \cos x + p^2 \cos 2x + p^3 \cos 3x + \&c.) e^{-kx},$$

and integrating both sides from $x=0$ to $x=\alpha$, and putting $p=1$,

$$\text{we get} \quad \pi = \frac{1}{k} (1 - e^{-k\alpha}) +$$

$$2e^{-k\alpha} {}^1S^\infty \frac{n \sin n\alpha}{k^2 + n^2} - 2ke^{-k\alpha} {}^1S^\infty \frac{\cos n\alpha}{k^2 + n^2} + 2 {}^1S^\infty \frac{k}{k^2 + n^2}.$$

Again, integrating from $x = 0$ to $x = 2\pi$, and putting $p = 1$, we find

$$\pi (e^{-2k\pi} + 1) = \frac{1 - e^{-2k\pi}}{k} + 2 {}^1S^\infty \frac{k}{k^2 + n^2} - 2e^{-2k\pi} {}^1S^\infty \frac{k}{k^2 + n^2},$$

$$\text{which gives } 2 {}^1S^\infty \frac{k}{k^2 + n^2} = \pi \cdot \frac{1 + e^{-2k\pi}}{1 - e^{-2k\pi}} - \frac{1}{k},$$

and substituting this value in the foregoing result, and calling the two sums we are in search of, S_1 and S_2 , we have

$$\frac{2\pi e^{-2k\pi + k\alpha}}{1 - e^{-2k\pi}} = \frac{1}{k} - 2S_1 + 2kS_2;$$

and changing the sign of k ,

$$\frac{-2\pi e^{-k\alpha}}{1 - e^{-2k\pi}} = -\frac{1}{k} - 2S_1 - 2kS_2;$$

therefore, adding and subtracting,

$$S_1 = {}^1S^\infty \frac{n \sin n\alpha}{k^2 + n^2} = \frac{\pi}{2} \frac{e^{k(\pi-\alpha)} - e^{-k(\pi-\alpha)}}{e^{k\pi} - e^{-k\pi}},$$

$$S_2 = {}^1S^\infty \frac{\cos n\alpha}{k^2 + n^2} = \frac{\pi}{2k} \frac{e^{k(\pi-\alpha)} + e^{-k(\pi-\alpha)}}{e^{k\pi} - e^{-k\pi}} - \frac{1}{2k^2}.$$

These formulæ were first given by Poisson, and may be considered as embracing the chief results which have hitherto been obtained relating to the summation of series of the sines and cosines of multiple angles.

119. The definite integrals required in the preceding investigation may be found as follows. Suppose

$$y = \frac{(1 - p^2)e^{-kx}}{1 - 2p \cos x + p^2},$$

to be the equation to a curve; and let it be proposed to find the limiting value of its area, from $x = 0$ to $x = \alpha$, on the supposition that p approaches continually to unity. First, we observe that

all the ordinates will be ultimately evanescent except those corresponding to $x = 0, 2\pi, 4\pi, \&c.$, which will be very large, because for those values $\frac{1+p}{1-p}$ becomes a factor of the expression for y . We will begin by supposing that $\alpha < 2\pi$; and therefore the only ordinates that we are concerned with are those immediately succeeding that through the origin; so that, making $p = 1 - \rho$, and then supposing x very small, we get successively

$$y = \frac{\rho(2-\rho)e^{-kx}}{\rho^2 + 4(1-\rho)\sin^2 \frac{1}{2}x} = \frac{2\rho}{\rho^2 + x^2}, \text{ ultimately;}$$

$$\therefore \int_0^{\alpha} dxy = 2 \tan^{-1} \frac{\alpha}{\rho}, \text{ and } \int_0^{\alpha} dxy_{p=1} = \pi, \text{ making } \rho = 0.$$

Next, suppose that the area is to be found from $x=0$ to $x=2\pi$; then, besides the area just found, there will be another portion immediately preceding the point for which $x=2\pi$; to find this latter portion put $x = 2\pi - x'$, and call the ordinate y' ;

$$\text{then } y' = \frac{(1-p^2)e^{-2k\pi+kx'}}{1-2p\cos x' + p^2} = e^{-2k\pi} y,$$

therefore the portion of the area immediately preceding the second limit $= \pi e^{-2k\pi}$;

$$\therefore \int_0^{2\pi} dxy_{p=1} = \pi + \pi e^{-2k\pi}.$$

120. As the following important theorems in Definite Integrals are based upon the preceding investigations, we shall here give the proofs of them. As in Art. 118 we have

$$y = \frac{(1-p^2)f(x)}{1-2p\cos(c-x)\frac{\pi}{\alpha} + p^2} = f(x) + 2^1 S^{\infty} p^n \cos n(c-x) \frac{\pi}{\alpha} \cdot f(x);$$

$$\therefore \int_{-\alpha}^{+\alpha} dxy_{p=1} = \int_{-\alpha}^{+\alpha} dx f(x) + 2^1 S^{\infty} \int_{-\alpha}^{+\alpha} dx \cos n(c-x) \frac{\pi}{\alpha} \cdot f(x).$$

Now, suppose c to be less than α , then for all values of x between $x = -\alpha$ and $x = +\alpha$, y is evanescent, except when $x = c$; if therefore we write $x = c + z$, supposing z exceedingly small,

and then integrate with respect to z , from $z=0$ to $z=\gamma$ any small finite value, and double the result, and make $p=1$, we shall obtain the value of $\int_{-a}^{+a} dx y_{p=1}$. But making $p=1-\rho$, we have

$$y = \frac{\rho(2-\rho)f(c+z)}{\rho^2 + 4(1-\rho)\sin^2 \frac{z\pi}{2a}} = \frac{2\rho f(c)}{\rho^2 + \frac{\pi^2 z^2}{a^2}} \text{ ultimately ;}$$

$$\therefore \int_0^\gamma dz y = \frac{2\alpha f(c)}{\pi} \tan^{-1} \frac{\pi\gamma}{\rho\alpha} ; \quad \int_0^\gamma dz y_{p=1} = \alpha f(c) ;$$

$$\therefore 2\alpha f(c) = \int_{-a}^{+a} dx f(x) + 2^1 S^\infty \int_{-a}^{+a} dx \cos n(c-x) \frac{\pi}{a} \cdot f(x). \quad (1)$$

121. Hence

$$\begin{aligned} f(c) = & \frac{1}{2a} \int_{-a}^{+a} dx f(x) + \frac{1}{a} \int_{-a}^{+a} dx \left\{ \cos(c-x) \frac{\pi}{a} \cdot f(x) \right. \\ & \left. + \cos(c-x) \frac{2\pi}{a} \cdot f(x) + \cos(c-x) \frac{3\pi}{a} \cdot f(x) + \dots \right\} ; \end{aligned}$$

therefore, making α infinite,

$$f(c) = \frac{1}{\pi} \text{ limit of } \frac{\pi}{a} \int_{-a}^{+a} dx \left\{ \cos(c-x) \frac{\pi}{a} + \cos(c-x) \frac{2\pi}{a} + \&c. \right\} f(x),$$

$$\text{or, } f(c) = \frac{1}{\pi} \int_0^\infty dz \left\{ \int_{-a}^{+a} dx \cos(c-x) z \cdot f(x) \right\} \text{ (Integ. Cal. Art. 116),}$$

a theorem given by Fourier, and included, as we see, in the theorem (1), which was first given by Poisson.

122. When in physical questions a definite integral arises whose value cannot be exhibited in finite terms, or in a form convenient for numerical calculation, the method of Quadratures is used as a substitute for Integration. This method consists in taking a series of values of the function to be integrated, multiplying these by the differences of the corresponding values of the independent variable, and adding together all the results. The sum of such results approximates to the value of the de-

finite integral, as the intervals of the independent variable are diminished. The method is equivalent to adopting, for the area of a curve which a definite integral gives, the sum of the areas of a system of inscribed or circumscribed polygons.

Let the equal intervals into which the independent variable x is divided be taken equal to unity, and let $f(x)$ be the function of x to be integrated between assigned limits a and b . Take a value of it $f(a + n - \frac{1}{2})$ at the middle of the n^{th} of the intervals into which $b - a$ is divided. If the intervals be very small, $f(x)$ may be considered constant from

$$x = a + n - 1 \text{ to } x = a + n,$$

or $\Delta^2 f(x)$ may be neglected; and the value of the definite integral is $\Sigma f(a + n - \frac{1}{2})$, where n is to receive in succession the values

$$1, 2, 3, \dots b - a.$$

For greater accuracy, suppose $\Delta^2 f(x)$ to be constant from

$$x = a \text{ to } x = b,$$

or $\Delta^3 f(x)$ to be neglected. Then

$$f(a + n - \frac{1}{2} + z) = f(a + n - \frac{1}{2}) + z\Delta f(a + n - \frac{1}{2}) + \frac{z^2}{2}\Delta^2 f(x),$$

and the integral for the interval in question is

$$\begin{aligned} & \int dz f(a + n - \frac{1}{2} + z), \text{ from } z = -\frac{1}{2} \text{ to } z = \frac{1}{2}, \\ & = f(a + n - \frac{1}{2}) + \frac{1}{6} \left(\frac{1}{8} + \frac{1}{8} \right) \Delta^2 f(x) = f(a + n - \frac{1}{2}) + \frac{1}{24} \Delta^2 f(x). \end{aligned}$$

The integral, consequently, from a to b , is

$$\Sigma \{ f(a + n - \frac{1}{2}) + \frac{1}{24} \Delta^2 f(x) \};$$

n receiving the same successive values as in the former case. In calculating the value of a definite integral by this method, the intervals are to be taken smaller as the variation of the function is more rapid.

Convergency and Divergency of Series.

123. A series $u_1 + u_2 + u_3 + \dots + u_n + \dots$ (to ∞) is called convergent, if the sum s_n of any number n of its terms approaches continually to a finite quantity s as its limit, when n is indefinitely increased; and divergent in the contrary case.

When a series is convergent, the sum of any number of consecutive terms after the n^{th} continually tends to zero, as n increases. For

$$u_{n+1} + u_{n+2} + \dots + u_{n+m} = s_{n+m} - s_n;$$

therefore, as n increases, the value of the first member continually approaches to $s - s$, or zero, as its limit. This being true when $m = 1$, we see also that u_{n+1} , or the general term u_n continually tends to zero as n increases; that is, each term is greater than the following one; but this, although a necessary condition, is not sufficient to insure the convergency of a series. Thus in the series

$$\frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n};$$

$u_n = \frac{1}{n}$ continually approaches to zero as n increases; but

$$s_{n+m} - s_n = \frac{1}{n+1} + \frac{1}{n+2} + \dots + \frac{1}{n+m}$$

is evidently greater than $\frac{m}{n+m} > \frac{1}{2}$, if $m = n$; and therefore the sum of n consecutive terms after the n^{th} , does not diminish indefinitely as n increases; consequently the series in question is divergent.

124. In the geometrical progression

$$a + ax + ax^2 + \dots + ax^{n-1} + \dots \quad (1),$$

$$s_n = a \cdot \frac{1 - x^n}{1 - x}, \quad s_{n+m} = a \cdot \frac{1 - x^{n+m}}{1 - x};$$

$$\therefore s_{n+m} - s_n = ax^n \cdot \frac{1 - x^m}{1 - x}.$$

Hence if $x < 1$, when n is infinite $x^n = 0$,

$$\therefore s = \frac{a}{1-x}; \text{ also } u_n = 0, s_{n+m} - s_n = 0;$$

both which results shew that the series is convergent.

But if $x > 1$, then $u_n = ax^{n-1}$ increases indefinitely with n , which alone shews that the series is divergent.

Hence the geometrical progression (1) is convergent or divergent, according as x is less or greater than unity; and it may be used as the test of the convergency or divergency of other series. For if a proposed series can be shewn to have no term greater than the corresponding term of (1) when $x < 1$, then that series is convergent; or if a proposed series can be shewn to have no term less than the corresponding term of (1) when $x > 1$, then that series is divergent.

Thus in the series for e the base of the natural logarithms

$$1 + \frac{1}{1} + \frac{1}{1 \cdot 2} + \frac{1}{1 \cdot 2 \cdot 3} + \dots + \frac{1}{\underline{n-1}} + \dots$$

the terms which follow the n^{th} , viz.

$$\frac{1}{\underline{n}} + \frac{1}{\underline{n+1}} + \frac{1}{\underline{n+2}} + \&c.$$

are evidently less than the corresponding terms of the geometrical progression

$$\frac{1}{\underline{n}} + \frac{1}{\underline{n}} \cdot \frac{1}{n} + \frac{1}{\underline{n}} \cdot \frac{1}{n^2} + \&c.$$

the sum of which is

$$\frac{1}{\underline{n}} \cdot \frac{1}{1 - \frac{1}{n}} = \frac{1}{\underline{n-1}} \cdot \frac{1}{n-1},$$

and continually tends to zero as n increases. Therefore the proposed series is convergent; and the error in taking the aggregate of its first n terms for its sum, is less than the quotient of the n^{th} term divided by $n-1$.

125. From the measure of convergency or divergency which a geometrical progression furnishes, we shall now proceed to deduce one or two other tests as given by Cauchy (*Cours d'Analyse Algébrique*).

The series $u_1 + u_2 + \dots + u_n + u_{n+1} \dots$ is convergent if the limit of the ratio of u_{n+1} to u_n when n is infinite, be less than unity; and divergent in the contrary case. Suppose that as n increases indefinitely, the ratio $\frac{u_{n+1}}{u_n}$ continually tends to become equal to a finite quantity k as its limit; and let i denote a number less than the difference between k and 1, so that the quantities $k-i$ and $k+i$ are both less than 1 when k is less than 1, and both greater than 1 in the contrary case. Then by giving n a sufficiently large value the ratio $\frac{u_{n+1}}{u_n}$ may be made to lie between $k-i$ and $k+i$ for all superior values of n ; and in the series

$$\begin{array}{ccccccc} u_n + u_n (k-i) + u_n (k-i)^2 + \dots & & & & & & \\ u_n + u_{n+1} & + & u_{n+2} & + & \dots & & \\ u_n + u_n (k+i) + u_n (k+i)^2 + \dots & & & & & & \end{array}$$

every term of the second will be intermediate in value to the corresponding terms of the first and third. But the first and third series evidently decrease indefinitely if u_n does so, as n is increased without limit, provided k be less than 1; therefore the second series which is intermediate in value to these is convergent under the same circumstances; and therefore the proposed series is convergent provided the limit of the ratio of u_{n+1} to u_n when n is infinite, be less than 1. Also the three series above written will all increase indefinitely if k be greater than 1, and the proposed series is in that case divergent. The test of convergency or divergency is here presented in the form most convenient for application; but it may be changed into the following: if $k=1$, the test gives no result in either form.

126. The series $u_1 + u_2 + \dots + u_n + \dots$ is convergent, or will become so, if the superior limit of $(u_n)^{\frac{1}{n}}$ be less than 1, when n is infinite; and divergent in the contrary case.

Let k denote the superior limit of $(u_n)^{\frac{1}{n}}$ when n is infinite; and first suppose $k < 1$; also, let a be any magnitude between k and 1, so that $k < a < 1$; then when n is increased indefinitely, $(u_n)^{\frac{1}{n}}$ cannot approach indefinitely near to k without finally becoming constantly less than a . Therefore it will be possible to take for n so large a value, that for that and all superior values, we may constantly have

$$(u_n)^{\frac{1}{n}} < a, \quad u_n < a^n.$$

Consequently, the proposed series will finish by always having its terms less than the corresponding terms of the geometrical progression

$$a + a^2 + \dots + a^n + a^{n+1} + \dots;$$

and as this series is convergent, a being < 1 , it follows that the proposed series will *à fortiori* end by being convergent.

Secondly, suppose $k > 1$; and take, as before, a between 1 and k , so that $k > a > 1$. Then, when n is indefinitely increased, $(u_n)^{\frac{1}{n}}$ cannot approach indefinitely near to k without finally becoming constantly $> a$; we shall therefore be able to satisfy the condition $(u_n)^{\frac{1}{n}} > a$, or $u_n > a^n$, by taking n sufficiently large; and consequently we shall always find in the series

$$u_1 + u_2 + \dots + u_n + u_{n+1} + \dots,$$

an indefinite number of terms greater than the corresponding terms of the geometrical progression

$$a + a^2 + \dots + a^n + a^{n+1} + \dots,$$

which is divergent, a being > 1 ; and therefore the proposed series will end by being divergent.

127. It may be shewn that if as n increases indefinitely u_n remains positive, and the ratio $\frac{u_{n+1}}{u_n}$ continually tends to become equal to a finite quantity k as its limit, the expression $(u_n)^{\frac{1}{n}}$

continually approaches to the same limit. For as in Art. 125, by giving to m a value large enough, we may make each of the ratios $\frac{u_{m+1}}{u_m}, \frac{u_{m+2}}{u_{m+1}}, \dots, \frac{u_{m+n}}{u_{m+n-1}}$ differ from k by a quantity as small as we please; and consequently the geometric mean between these ratios $\left(\frac{u_{m+n}}{u_m}\right)^{\frac{1}{n}}$ may, by sufficiently increasing n , be made to differ from k by a quantity i as small as we please. We shall therefore have, taking n sufficiently large,

$$(u_{m+n})^{\frac{1}{n}} = \{u_{n(1+\frac{m}{n})}\}^{\frac{1}{n}} = (k \pm i) (u_m)^{\frac{1}{n}}.$$

Now suppose n infinite, so that i becomes equal to zero, and we get

$$\text{limit of } (u_n)^{\frac{1}{n}}_{n=\infty} = k = \text{limit of } \left(\frac{u_{n+1}}{u_n}\right)_{n=\infty}.$$

128. Suppose that the series $u_1 + u_2 + u_3 + \dots$ consists of both positive and negative terms; then if $v_1, v_2, v_3, \&c.$ be the numerical values of these terms, so that $u_1 = \pm v_1, u_2 = \pm v_2, \&c.$, it is evident that the sum of the proposed series can never surpass that of the series $v_1 + v_2 + v_3 + \dots$; if therefore the latter series be convergent, that is, if $\left(\frac{v_{n+1}}{v_n}\right)_{n=\infty} < 1$, the proposed series will be convergent; or if the latter series finish by having terms greater than any assignable magnitude, that is, if $\left(\frac{v_{n+1}}{v_n}\right)_{n=\infty} > 1$, the same thing will happen to the proposed series, which will consequently be divergent. Hence the above test is applicable to series consisting of both positive and negative terms, provided we use the numerical values of the terms without regard to signs; and it fails, as in the preceding case, when $k = 1$.

$$129. \text{ Let } a_0 + a_1x + a_2x^2 + \dots + a_nx^n + \dots \quad (1)$$

be a series arranged according to positive and ascending powers of the variable x ; the coefficients being positive or negative; then, by what has been proved, this series will be convergent or divergent, according as kx ,

where $k = (a_n)^{\frac{1}{n}} = \frac{a_{n+1}}{a_n}$ when n is infinite,

is numerically less or greater than 1. Hence for all values of x between $-\frac{1}{k}$ and $+\frac{1}{k}$ the series will be convergent, and for all values of x beyond those limits it will be divergent.

$$\text{Ex. 1. } \frac{a}{1}x + \frac{a(a-1)}{1 \cdot 2}x^2 + \frac{a(a-1)(a-2)}{1 \cdot 2 \cdot 3}x^3 + \&c.$$

$$\text{Here } \frac{a_{n+1}}{a_n} = \frac{a-n}{n+1} = -1, \text{ when } n \text{ is infinite;}$$

therefore the series is convergent or divergent, according as x lies between $+1$ and -1 , or without those limits.

$$\text{Ex. 2. } \frac{a}{1} + \frac{a^2}{2} + \frac{a^3}{3} + \&c.,$$

$$u_n = \frac{a^n}{n}, \quad \therefore \frac{u_{n+1}}{u_n} = \frac{na}{n+1} = a \text{ when } n \text{ is infinite;}$$

therefore the series is convergent or divergent, according as $a < \text{or} > 1$.

$$\text{Ex. 3. } \frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n},$$

$$\frac{u_{n+1}}{u_n} = \frac{1}{n+1} = 0, \text{ when } n \text{ is infinite,}$$

therefore the series converges. Similarly, it will appear that the series for $\sin x$ and $\cos x$ are always convergent; and that for $\tan^{-1} x$ convergent for all values of x situated between $+1$ and -1 .

130. The series $u_1 + u_2 + \dots + u_n + \dots$ is convergent, or will become so, if the inferior limit of $\frac{\log u_n}{\log \frac{1}{n}}$ be > 1 , when n

is infinite; and divergent in the contrary case.

Let k denote the inferior limit of $\frac{\log u_n}{\log \frac{1}{n}}$ when n is infinite,

and first suppose $k > 1$; also let a be any number between k and 1, so that $k > a > 1$. Then when n is indefinitely increased,

$\frac{\log u_n}{\log \frac{1}{n}}$ or its equal $\frac{\log \frac{1}{u_n}}{\log n}$ cannot approach indefinitely near to

k without finally becoming constantly greater than a . Therefore it will be possible to take for n so large a value, that for that and all superior values we may constantly have

$$\log \frac{1}{u_n} > a \log n > \log n^a, \quad \text{or } \frac{1}{u_n} > n^a, \quad \text{or } u_n < \frac{1}{n^a}.$$

Consequently the proposed series will finish by always having its terms less than the corresponding terms of the converging series (Ex. Art. 131)

$$\frac{1}{1^a} + \frac{1}{2^a} + \frac{1}{3^a} + \dots + \frac{1}{n^a} + \frac{1}{(n+1)^a} + \&c.$$

and therefore will itself end by being, *à fortiori*, convergent.

Similarly, if $k < 1$, it may be shewn that the proposed series will finish by being divergent.

131. The series $u_0 + u_1 + u_2 + u_3 + u_4 + u_5 + \&c. \dots\dots (1)$,
each term of which is less than the immediately preceding term,

and the series $u_0 + 2u_1 + 4u_2 + 8u_3 + 16u_4 + \&c. \dots\dots (2)$,
are convergent or divergent at the same time.

Suppose series (1) to be convergent and that its sum = s ,

$$\text{then } u_0 = u_0, \quad 2u_1 = 2u_1, \quad 4u_2 < 2u_2 + 2u_2,$$

$$8u_3 < 2u_4 + 2u_5 + 2u_6 + 2u_7, \quad \&c.;$$

$$\therefore u_0 + 2u_1 + 4u_2 + 8u_3 + 16u_4 + \&c.$$

$$< u_0 + 2(u_1 + u_2 + u_3 + \&c.) < 2s - u_0,$$

consequently series (2) is convergent. Next, suppose series (1) divergent, then

$$u_0 = u_0, \quad 2u_1 > u_1 + u_2, \quad 4u_3 > u_3 + u_4 + u_5 + u_6, \quad \&c.;$$

$$\therefore u_0 + 2u_1 + 4u_3 + 8u_7 + \&c. > u_0 + u_1 + u_2 + u_3 + \&c.,$$

and is therefore divergent.

Ex. Let series (1) be $\frac{1}{1^m} + \frac{1}{2^m} + \frac{1}{3^m} + \frac{1}{4^m} + \&c. \dots (3)$,

$$\text{then series (2) is } \frac{1}{1^m} + \frac{1}{2^{m-1}} + \frac{1}{4^{m-1}} + \frac{1}{8^{m-1}} + \&c.,$$

a geometric progression convergent when $m > 1$, and divergent in the contrary case; consequently series (3) will be convergent if $m > 1$, and divergent if $m =$ or < 1 .

132. The alternating series $u_1 - u_2 + u_3 - u_4 + \&c.$, is convergent, if the numerical value of the terms decreases without limit.

For, by writing it in the forms

$$u_1 - (u_2 - u_3) - (u_4 - u_5) - \&c.,$$

$$u_1 - u_2 + (u_3 - u_4) + \&c.;$$

we see that it is $> u_1 - u_2$ and $< u_1$, and therefore is convergent.

Thus the sum of the series $\frac{1}{1} - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \&c.$,

lies between 1 and $\frac{1}{2}$; also the series $\frac{1}{1^m} - \frac{1}{2^m} + \frac{1}{3^m} - \frac{1}{4^m} + \&c.$,

is convergent for all positive values of m .

133. It is also shewn in the Work from which these tests of convergency are taken, that two converging series all whose terms are positive, will by their addition or multiplication produce new converging series, whose sums result from the addition or multiplication of the sums of the former. If the proposed

series be imaginary, having its general term of the form $v_n + w_n \sqrt{-1}$, it may be resolved into the two series

$$v_1 + v_2 + \dots + v_n + \dots, \quad \sqrt{-1} (w_1 + w_2 + \dots + w_n + \dots) \dots (1),$$

to which the foregoing tests may be applied separately. If we assume

$$v_n + w_n \sqrt{-1} = \rho_n (\cos \theta_n + \sqrt{-1} \sin \theta_n),$$

then $v_n^2 + w_n^2 = \rho_n^2$, so that v_n and w_n are both less than ρ_n ; if therefore the series of moduli $\rho_1 + \rho_2 + \dots + \rho_n + \dots$ be convergent, then each of the series (1) will be convergent.

Interpolation of Series.

134. When a series of values of a quantity has been obtained either by observation or calculation, it is of great importance to be able to insert other values between them, such as would have resulted from a similar observation or calculation, without the labour of performing these. This is the object of Interpolation; and in this the Calculus of Finite Differences finds one of its chief uses. More strictly, Interpolation of Series is the inserting among the terms of a given series, new terms subject to the same law as the first. In doing this, the terms of the series are considered as particular values of the function which expresses its general term, corresponding to a given regular succession of indices; and it is the business of Interpolation to discover that general term; or at least to assign such a function of the index as shall represent the given series of values, and, approximately, all intermediate values. The problem thus requiring us to assign the analytical expression of a function from a limited number of its numerical values, is plainly indeterminate; it is the same as to form the equation to a curve which shall pass through a limited number of points, whose abscissæ represent the values of the independent variable, and the ordinates those of the function, without giving the species of the curve; which, as is evident, may be done in an infinite variety of ways. But if the given terms are numerous and near to each other, the expression

for the general term, within the limits of the given quantities, may be found to a great degree of accuracy.

135. There are two principal cases to be considered; first, when the given values of $f(x)$, namely,

$$f(x_1), f(x_1 + h), f(x_1 + 2h), \dots f\{x_1 + (n-1)h\},$$

$$\text{or } u_1, u_2, u_3, \dots u_n,$$

as we shall write them, correspond to values of the independent variable, $x_1, x_1 + h, x_1 + 2h$, &c., $x_1 + (n-1)h$, in arithmetical progression. And secondly, when the given values $f(x_1), f(x_2), \dots f(x_n)$, or $u_1, u_2, \dots u_n$, correspond to values $x_1, x_2, \dots x_n$, of the independent variable, not obeying any assigned law.

136. Having given u_1, u_2, u_3 , &c. u_n , n values of a function $f(x)$, corresponding to the n values of the independent variable $x_1, x_1 + h, \dots x_1 + (n-1)h$, to find an expression for any intermediate value $f(x_1 + k)$.

$$\text{Since } f(x + nh) = f(x) + n\Delta f(x) + \frac{n(n-1)}{1 \cdot 2} \Delta^2 f(x) + \&c.$$

(Art. 23), changing x into x_1 , and then replacing nh by k , and n by $\frac{k}{h}$ we get

$$f(x_1 + k) = u_1 + \frac{k}{h} \Delta u_1 + \frac{k(k-h)}{1 \cdot 2h^2} \Delta^2 u_1 + \&c. \\ + \frac{k(k-h) \dots \{k - (n-2)h\}}{[n-1] \cdot h^{n-1}} \Delta^{n-1} u_1;$$

in which, since

$$\left. \begin{aligned} \Delta u_1 &= u_2 - u_1, \\ \Delta^2 u_1 &= u_3 - 2u_2 + u_1, \\ &\dots\dots\dots \\ \Delta^{n-1} u_1 &= u_n - (n-1)u_{n-1} + \frac{(n-1)(n-2)}{1 \cdot 2} u_{n-2} - \&c. \dots \pm u_1, \end{aligned} \right\} (2)$$

if we make $k=0, h, 2h$, &c., $(n-1)h$, the second member assumes the n values $u_1, u_2, \dots u_n$; and not only this, but if we assume for k any value whatever between 0 and $(n-1)h$, we

shall obtain the value of the function corresponding to that value of the independent variable.

137. In applying the above formula, the simplest mode is not to calculate Δu_1 , $\Delta^2 u_1$, &c., by equations (2), but by continued subtraction of the given terms; that is, we must write down the series of given values $u_1, u_2, \dots u_n$, and subtract each from the succeeding one; next subtract each of these differences from the succeeding difference; then perform the same operation upon the new differences; and so on, till the process terminates; the first terms of these series of differences are the values of Δu_1 , $\Delta^2 u_1$, ... $\Delta^{n-1} u_1$. Unless the terms of the given series by continued subtraction lead to a constant difference, the expression for $f(x_1 + k)$ will have as many terms as the given series of values has.

Ex. 1. Having given the values of $\sin 30^\circ$, $\sin 31^\circ$, $\sin 32^\circ$, $\sin 33^\circ$, to find $\sin (30^\circ + k')$, k being between 0 and $180'$. Here

| | | | |
|------------------|----------|------------|------------|
| $u_1 = .5$ | Δ | | |
| $u_2 = .5150381$ | 150381 | Δ^2 | |
| $u_3 = .5299193$ | 148812 | -1569 | Δ^3 |
| $u_4 = .5446390$ | 147197 | -1615 | -46; |

$$\therefore \Delta u_1 = 0.0150381, \Delta^2 u_1 = -0.0001569, \Delta^3 u_1 = -0.0000046,$$

$$\begin{aligned} \therefore \sin (30^\circ + k') &= .5 + \frac{k}{60} \Delta u_1 + \frac{k(k-60)}{2 \cdot 60^2} \Delta^2 u_1 \\ &\quad + \frac{k(k-60)(k-120)}{2 \cdot 3 \cdot 60^3} \Delta^3 u_1. \end{aligned}$$

If $k = 20$, it will be found that $\sin 30^\circ 20' = .5050299$, which is too large only by a unit in the seventh place of decimals.

Ex. 2. Having given

$$\log 3.14 = 0.496929, \quad \log 3.15 = 0.498310,$$

$$\log 3.16 = 0.499687, \quad \log 3.17 = 0.501059;$$

$$\text{shew that } \log 3.14159 = 0.497149.$$

Ex. 3. Having given u_1, u_2, u_3, u_4 , four right ascensions (declinations, longitudes, &c.) of the Moon at intervals of 12 hours, to find its value t hours after the time corresponding to the second value. Here $h = 12$, $k = 12 + t$, and if the required right ascension $= u_2 + \delta$, then

$$u_2 + \delta = u_1 + \left(1 + \frac{t}{12}\right) \Delta u_1 + \frac{(t+12)t}{2 \cdot 12^2} \Delta^2 u_1 + \frac{(t+12)t(t-12)}{2 \cdot 3 \cdot 12^3} \Delta^3 u_1,$$

$$\text{or } \delta = \frac{t}{12} \left(\Delta^1 + \frac{1}{2} \Delta^2 - \frac{1}{6} \Delta^3 \right) + \frac{1}{2} \left(\frac{t}{12} \right)^2 \Delta^2 + \frac{1}{6} \left(\frac{t}{12} \right)^3 \Delta^3,$$

when developed in powers of $\frac{t}{12}$, Δ^1 being written for Δu_1 , Δ^2 for $\Delta^2 u_1$, &c.

138. Between every two consecutive terms of a given series, to interpolate any number of equidistant terms.

Let u_1, u_2, u_3 , &c., u_n be the given series, and let $m-1$ be the number of equidistant terms to be inserted between every two consecutive terms; then the new series will be

$$u_1, \quad u_{\frac{m+1}{m}}, \quad u_{\frac{m+2}{m}}, \quad \dots \quad u_{\frac{2m-1}{m}}, \quad u_2, \quad u_{\frac{2m+1}{m}}, \quad \&c.$$

If therefore v_{r+1} denote the $r+1^{\text{th}}$ term of this series, we have

$$v_{r+1} = u_{\frac{m+r}{m}} = f\left(1 + \frac{r}{m}\right),$$

$$\text{or } v_{r+1} = u_1 + \frac{r}{m} \Delta u_1 + \frac{r(r-m)}{1 \cdot 2m^2} \Delta^2 u_1 + \&c.$$

Hence, taking r from 1 to $m-1$, we get the terms inserted between u_1 and u_2 ; next taking r from $m+1$ to $2m-1$, we get the terms between u_2 and u_3 ; and so on. The differences Δu_1 , $\Delta^2 u_1$, &c., are to be computed by continued subtraction as in Art. (137); and the series for v_{r+1} will have as many terms as the proposed series has, unless those terms by continued subtraction lead to a constant difference.

Ex. To insert three equidistant terms between every two

consecutive ones of the series 1, 7, 15, 28, 49, &c. Here $m = 4$, and

$$\begin{array}{rcl} \Delta, & 6, & 8, \quad 13, \quad 21, \\ \Delta^2 & & 2, \quad 5, \quad 8, \\ \Delta^3 & & 3, \quad 3, \end{array}$$

$$\therefore v_{r+1} = 1 + \frac{6r}{4} + \frac{r(r-4)}{16} + \frac{r(r-4)(r-8)}{128} = \frac{128 + 192r - 4r^2 + r^3}{128};$$

and the series is 1, $\frac{317}{128}$, $\frac{504}{128}$, $\frac{695}{128}$, 7, $\frac{1113}{128}$, &c.

139. The formula of Art. 136 may be presented under a different form by changing k into $x - x_1$, which gives

$$\begin{aligned} f(x) = u_1 + \frac{x - x_1}{h} \Delta u_1 + \frac{(x - x_1)(x - x_1 - h)}{1 \cdot 2 \cdot h^2} \Delta^2 u_1 + \&c. \\ + \frac{(x - x_1)(x - x_1 - h) \dots \{x - x_1 - (n-2)h\}}{\lfloor n-1 \rfloor \cdot h^{n-1}} \Delta^{n-1} u_1; \end{aligned}$$

where $f(x)$ is a function of x which, as x assumes the n values x_1 , $x_1 + h$, &c., $x_1 + (n-1)h$, successively assumes the corresponding values u_1 , u_2, \dots, u_n ; and for any other value of x within, or not far beyond, the limits x_1 and $x_1 + (n-1)h$, it gives the value of the corresponding interpolated term. If we put $h = 1$, the formula is adapted to the case where the increment of the principal variable is unity.

140. In any series of consecutive equidistant values of a function, where one is deficient to insert that one.

Let u_1 , u_2 , u_3 , &c., u_n be the values of the function corresponding to the values x_1 , $x_1 + h, \dots, x_1 + (n-1)h$ for x . Then assuming that $\Delta^{n-1} u_1 = 0$, or that the $(n-2)^{\text{th}}$ differences are constant, which will almost always be the case in tabulated results, we have

$$\Delta^{n-1} u_1 = u_n - (n-1) u_{n-1} + \frac{(n-1)(n-2)}{1 \cdot 2} u_{n-2} - \&c. \pm u_1 = 0,$$

an equation of the first degree from which any one of the values as u_r may be found, if the rest be known. Having thus completed the system of values, we may interpolate any intermediate term $f(x_1 + k)$ by the method of Art. 136. If two values out of n are deficient, then we must suppose

$$\Delta^{n-2}u_1 = 0, \quad \Delta^{n-2}u_2 = 0, \quad \text{or}$$

$$u_{n-1} - (n-2)u_{n-2} + \frac{(n-2)(n-3)}{1 \cdot 2}u_{n-3} - \&c. \pm u_1 = 0,$$

$$u_n - (n-2)u_{n-1} + \frac{(n-2)(n-3)}{1 \cdot 2}u_{n-2} - \&c. \pm u_2 = 0;$$

which equations will suffice to determine any two of the values in terms of the rest. In the same way any number of deficient terms may be inserted.

Ex. 1. Given the cube roots of 121, 122, 124, 125, to find that of 123.

$$u_1 = 4.946088, \quad u_2 = 4.959675, \quad u_4 = 4.986631. \quad u_5 = 5,$$

$$\Delta^4 u_1 = u_5 - 4u_4 + 6u_3 - 4u_2 + u_1 = 0;$$

$$\therefore u_3 = \frac{1}{6} \{4(u_2 + u_4) - u_1 - u_5\} = 4.973190.$$

Ex. 2. Given u_1, u_2, u_5, u_6 , to find u_3, u_4 ,

$$\Delta^4 u_1 = u_6 - 4u_4 + 6u_3 - 4u_2 + u_1 = 0.$$

$$\Delta^4 u_2 = u_6 - 4u_5 + 6u_4 - 4u_3 + u_2 = 0;$$

$$\therefore u_3 = \frac{1}{10} (-3u_1 + 10u_2 + 5u_5 - 2u_6),$$

$$u_4 = \frac{1}{10} (-2u_1 + 5u_2 + 10u_5 - 3u_6).$$

Ex. 3. Given the logarithms of 121, 122, 125, 126 equal respectively to 2.0827854, 2.0863598, 2.0969100, 2.1003705; shew that the logarithms of 123, 124, are equal respectively to 2.0899051, 2.0934217.

141. We next come to the case where the given values

$$f(x_1), f(x_2), \&c., f(x_n), \quad \text{or } u_1, u_2, \dots, u_n,$$

correspond to values x_1, x_2, \dots, x_n , not obeying any assigned law; and it is required to determine a rational integral function of $n-1$ dimensions, $f(x)$, which shall assume the n given values u_1, u_2, \dots, u_n , when for x the values x_1, x_2, x_3 , &c., x_n are successively substituted.

Since $f(x)$ is of $(n-1)$ dimensions, we may assume

$$\frac{f(x)}{(x-x_1)(x-x_2)\dots(x-x_n)} = \frac{C_1}{x-x_1} + \frac{C_2}{x-x_2} + \dots + \frac{C_n}{x-x_n};$$

$$\therefore f(x) = C_1(x-x_2)(x-x_3)\dots(x-x_n)$$

$$+ C_2(x-x_1)(x-x_3)\dots(x-x_n) + \&c. + C_n(x-x_1)(x-x_2)\dots(x-x_{n-1}).$$

Now make $x = x_1, x_2$, &c., x_n , successively; and observing that the corresponding values of the first member are u_1, u_2, \dots, u_n , we get

$$u_1 = C_1(x_1-x_2)(x_1-x_3)\dots(x_1-x_n),$$

$$u_2 = C_2(x_2-x_1)(x_2-x_3)\dots(x_2-x_n)$$

$$\dots\dots\dots$$

$$u_n = C_n(x_n-x_1)(x_n-x_2)\dots(x_n-x_{n-1});$$

$$\begin{aligned} \therefore f(x) &= u_1 \frac{(x-x_2)(x-x_3)\dots(x-x_n)}{(x_1-x_2)(x_1-x_3)\dots(x_1-x_n)} \\ &+ u_2 \frac{(x-x_1)(x-x_3)\dots(x-x_n)}{(x_2-x_1)(x_2-x_3)\dots(x_2-x_n)} + \&c. \\ &+ u_n \frac{(x-x_1)(x-x_2)\dots(x-x_{n-1})}{(x_n-x_1)(x_n-x_2)\dots(x_n-x_{n-1})}, \end{aligned}$$

which is Lagrange's Theorem for interpolation.

Ex. To find a function of x which, when $x = 1, 3, 6, 12$, shall assume the values $1, 7, 10, -8$.

$$\begin{aligned} f(x) &= -\frac{(x-3)(x-6)(x-12)}{2 \cdot 5 \cdot 11} + 7 \cdot \frac{(x-1)(x-6)(x-12)}{2 \cdot 3 \cdot 9} \\ &- \frac{(x-1)(x-3)(x-12)}{3 \cdot 3} - 4 \cdot \frac{(x-1)(x-3)(x-6)}{11 \cdot 9 \cdot 3}. \end{aligned}$$

142. To determine the maximum or minimum value of a function, from three of its values near its maximum or minimum, and the three corresponding values of the independent variable.

If u_1, u_2, u_3 be the given values of u , and x_1, x_2, x_3 those of x , we have

$$u = u_1 \frac{(x - x_2)(x - x_3)}{(x_1 - x_2)(x_1 - x_3)} + u_2 \frac{(x - x_1)(x - x_3)}{(x_2 - x_1)(x_2 - x_3)} + u_3 \frac{(x - x_1)(x - x_2)}{(x_3 - x_1)(x_3 - x_2)}.$$

Hence, putting $\frac{du}{dx} = 0$, we find

$$u_1 (x_3 - x_2) (2x - x_2 - x_3) + u_2 (x_3 - x_1) (2x - x_1 - x_3) \\ + u_3 (x_1 - x_2) (2x - x_1 - x_2) = 0;$$

$$\therefore x = \frac{u_1 (x_2^2 - x_3^2) - u_2 (x_1^2 - x_3^2) + u_3 (x_1^2 - x_2^2)}{2u_1 (x_2 - x_3) - 2u_2 (x_1 - x_3) + 2u_3 (x_1 - x_2)},$$

the value of x at which u is a maximum or minimum. This formula is useful in various Astronomical problems, as for instance, to determine the meridian altitude of a heavenly body, when an observation exactly on the meridian cannot be obtained.

MATHEMATICAL PUBLICATIONS

OF THE

REV. J. HYMERS, D.D.

LATE FELLOW AND TUTOR OF ST JOHN'S COLLEGE, CAMBRIDGE.

THE ELEMENTS OF THE THEORY OF ASTRONOMY.

Second Edition, revised and enlarged.

8vo. 14s.

- A TREATISE ON CONIC SECTIONS, AND THE APPLICATION OF ALGEBRA TO GEOMETRY. Third Edition, revised and enlarged.

8vo. 9s.

- A TREATISE ON ANALYTICAL GEOMETRY OF THREE DIMENSIONS, containing the Theory of Curve Surfaces and of Curves of Double Curvature. Third Edition, revised and enlarged.

8vo. 10s. 6d.

- A TREATISE ON THE THEORY OF ALGEBRAICAL EQUATIONS. Third Edition, revised and enlarged.

8vo. 10s. 6d.

- A TREATISE ON THE INTEGRAL CALCULUS, containing the integration of explicit functions of one variable, together with the Theory of Definite Integrals, and of Elliptic Functions. Third Edition, revised and enlarged.

8vo. 10s. 6d.

- A TREATISE ON DIFFERENTIAL EQUATIONS, AND ON THE CALCULUS OF FINITE DIFFERENCES. Second Edition, enlarged.

8vo.

- A TREATISE ON PLANE AND SPHERICAL TRIGONOMETRY, and on Trigonometrical Tables and Logarithms, together with a selection of Problems and the solutions of the same. Fourth Edition, revised.

8vo. 8s. 6d.

LIST OF WORKS in GENERAL LITERATURE

PUBLISHED BY

Messrs. LONGMAN, BROWN, GREEN, LONGMANS, and ROBERTS,

39, PATERNOSTER ROW, LONDON.

CLASSIFIED INDEX.

Agriculture and Rural Affairs.

| | |
|-------------------------------|----|
| Baydon on Valuing Rents, &c. | 3 |
| Cecil's Stud Farm | 6 |
| Hoskyns's Talpa | 10 |
| Loudon's Agriculture | 12 |
| Low's Elements of Agriculture | 13 |

Arts, Manufactures, and Architecture.

| | |
|-------------------------------------|----|
| Bonnie on the Screw Propeller | 4 |
| Brande's Dictionary of Science, &c. | 4 |
| " Organic Chemistry | 6 |
| Chevel on Colour | 6 |
| Cressy's Civil Engineering | 6 |
| Fairbairn's Informa. for Engineers | 7 |
| Gwilt's Encyclo. of Architecture | 8 |
| Harford's Plates from N. Angelo | 8 |
| Humphreys's Pencil Illustrations | 11 |
| Jameson's Sacred & Legendary Art | 11 |
| " Commoplace-Book | 11 |
| König's Pictorial Life of Luther | 8 |
| Loudon's Rural Architecture | 13 |
| MacDougall's Theory of War | 13 |
| Maian's Aphorisms on Drawing | 14 |
| Moseley's Engineering | 16 |
| Piesse's Art of Perfumery | 17 |
| Richardson's Art of Horsemanship | 18 |
| Scrivener on the Iron Trade | 19 |
| Stark's Printing | 22 |
| Steam-Engine, by the Artisan Club | 4 |
| Ure's Dictionary of Arts, &c. | 23 |

Biography.

| | |
|------------------------------------|----|
| Arago's Autobiography | 22 |
| " Lives of Scientific Men | 22 |
| Bodenstedt and Wagner's Schamyl | 22 |
| Buckingham's (J. S.) Memoirs | 5 |
| Bunsen's Hippolytus | 6 |
| Cockayne's Marshal Torrence | 6 |
| Crosse's (Andrew) Memorials | 22 |
| Foster's De Foe and Churchill | 22 |
| Green's Princesses of England | 8 |
| Harford's Life of Michael Angelo | 8 |
| Hayward's Chesterfield and Selwyn | 22 |
| Holcroft's Memoirs | 22 |
| Lardner's Cabinet Cyclopædia | 12 |
| Maunder's Biographical Treasury | 12 |
| Memoir of the Duke of Wellington | 22 |
| Memoirs of James Montgomery | 15 |
| Merrill's Memoirs of Cicero | 15 |
| Mountain's (Col.) Memoirs | 16 |
| Perry's (Admiral) Memoirs | 17 |
| Rogers's Life and Genius of Fuller | 22 |
| Russell's Memoirs of Moore | 16 |
| Sonthey's Life of Wesley | 20 |
| " Life and Correspondence | 20 |
| " Select Correspondence | 20 |
| Stephen's Ecclesiastical Biography | 20 |
| Strickland's Queens of England | 21 |
| Sydney Smith's Memoirs | 20 |
| Symonds's (Admiral) Memoirs | 21 |
| Taylor's Loyola | 21 |
| " Wesley | 21 |
| Waterton's Autobiography & Essays | 24 |

Books of General Utility.

| | |
|-----------------------------------|----|
| Acton's Bread-Book | 3 |
| " Cookery | 3 |
| Black's Treatise on Brewing | 4 |
| Cabinet Gazetteer | 6 |
| " Lawyer | 6 |
| Curt's Invalid's Own Book | 6 |
| Gilbart's Logic for the Million | 6 |
| Hints on Etiquette | 6 |
| How to Nurse Sick Children | 10 |
| Hudson's Executor's Guide | 10 |
| " on Making Wills | 10 |
| Kesteven's Domestic Medicine | 12 |
| Lardner's Cabinet Cyclopædia | 12 |
| Loudon's Lady's Country Companion | 13 |
| Maunder's Treasury of Knowledge | 14 |
| " Biographical Treasury | 14 |
| " Geographical Treasury | 14 |

| | |
|----------------------------------|----|
| Maunder's Scientific Treasury | 14 |
| " Treasury of History | 14 |
| " Natural History | 14 |
| Piesse's Art of Perfumery | 17 |
| Pocket and the Stind | 8 |
| Pycroft's English Reading | 17 |
| Roe's Medical Guide | 18 |
| Rich's Comp. to Latin Dictionary | 18 |
| Richardson's Art of Horsemanship | 18 |
| Riddle's Latin Dictionaries | 18 |
| Rogers's English Thesaurus | 18 |
| Rowton's Debater | 18 |
| Short Whist | 21 |
| Thomson's Interest Tables | 21 |
| Webster's Domestic Economy | 24 |
| West on Children's Diseases | 24 |
| Willrich's Popular Tables | 24 |
| Wilmot's Blackstone | 24 |

Botany and Gardening.

| | |
|---------------------------------------|----|
| Hassall's British Freshwater Algae | 9 |
| Hooker's British Flora | 9 |
| " Guide to Kew Gardens | 9 |
| Lindley's Introduction to Botany | 9 |
| " Theory of Horticulture | 11 |
| Loudon's Hortus Britannicus | 13 |
| " Amateur Gardener | 13 |
| " Trees and Shrubs | 12 |
| " Gardening | 12 |
| " Plants | 13 |
| " Self-Instruction for Gardeners, &c. | 13 |
| Pereira's Materia Medica | 17 |
| River's Rose-Amateur's Guide | 17 |
| Wilson's British Mosses | 24 |

Chronology.

| | |
|-----------------------------------|----|
| Blair's Chronological Tables | 4 |
| Brewer's Historical Atlas | 4 |
| Bunsen's Ancient Egypt | 6 |
| Calendars of English State Papers | 5 |
| Haydn's Britain's India | 9 |
| Jacquinet's Chronology | 11 |
| Nicolas's Chronology of History | 12 |

Commerce and Mercantile Affairs.

| | |
|-----------------------------------|----|
| Gilbart's Treatise on Banking | 8 |
| Lorimer's Young Master Mariner | 12 |
| Maclean's Banking | 13 |
| McCulloch's Commerce & Navigation | 14 |
| Scrivener on Iron Trade | 19 |
| Thomson's Interest Tables | 21 |
| Tooke's History of Prices | 21 |

Criticism, History, and Memoirs.

| | |
|-------------------------------------|----|
| Blair's Chron. and Hist. Tables | 4 |
| Brewer's Historical Atlas | 4 |
| Bunsen's Ancient Egypt | 6 |
| " Hippolytus | 6 |
| Burton's History of Scotland | 6 |
| Calendars of English State Papers | 5 |
| Chapman's Gustavus Adolphus | 6 |
| Conybeare and Howson's St. Paul | 6 |
| Connolly's Sappers and Miners | 6 |
| Gleig's Leipzig Campaign | 22 |
| Gurney's Historical Sketches | 8 |
| Herschel's Essays and Addresses | 9 |
| Jeffrey's (Lord) Contributions | 11 |
| Kemble's Anglo-Saxons | 11 |
| Lardner's Cabinet Cyclopædia | 12 |
| Macaulay's Crit. and Hist. Essays | 12 |
| " History of England | 12 |
| " Speeches | 13 |
| Mackintosh's Miscellaneous Works | 13 |
| " History of England | 13 |
| McCulloch's Geographical Dictionary | 14 |
| Maunder's Treasury of History | 14 |
| Memoir of the Duke of Wellington | 22 |
| Merrill's History of Rome | 16 |
| " Roman Republic | 16 |
| Milner's Church History | 16 |
| Moore's (Thomas) Memoirs, &c. | 16 |
| Mure's Greek Literature | 18 |
| Perry's Franks | 17 |

| | |
|------------------------------------|----|
| Raikes's Journal | 17 |
| Ranke's Ferdinand & Maximilian | 17 |
| Riddle's Latin Dictionaries | 18 |
| Rogers's Essays from Edinb. Review | 18 |
| Rogers's English Thesaurus | 18 |
| Schmitt's History of Greece | 18 |
| Southey's Doctor | 20 |
| Stephen's Ecclesiastical Biography | 20 |
| " Lectures on French History | 20 |
| Sydney Smith's Works | 20 |
| " Select Works | 22 |
| " Lectures | 22 |
| " Memoirs | 20 |
| Taylor's Loyola | 21 |
| " Wesley | 21 |
| Thirlwall's History of Greece | 21 |
| Thomson's Historical Notes | 21 |
| Thornbury's Shakespeare's England | 21 |
| Townsend's State Trials | 21 |
| Turkey and Christendom | 22 |
| Turner's Anglo-Saxons | 23 |
| " Middle Ages | 23 |
| " Sacred Hist. of the World | 23 |
| Vehse's Austrian Court | 23 |
| Wade's England's Greatness | 24 |
| Whitlocke's Swedish Embassy | 24 |
| Young's Christ of History | 24 |

Geography and Atlases.

| | |
|-------------------------------------|----|
| Brewer's Historical Atlas | 4 |
| Butter's Geography and Atlases | 6 |
| Cabinet Gazetteer | 6 |
| Corwall, the Mines, &c. | 22 |
| Darrius's Morocco | 22 |
| Hugbie's Australian Colonies | 22 |
| Johnston's General Gazetteer | 11 |
| McCulloch's Geographical Dictionary | 14 |
| " Russia and Turkey | 22 |
| Maunder's Treasury of Geography | 14 |
| Mayne's Arctic Discoveries | 22 |
| Murray's Encyclo. of Geography | 16 |
| Sharp's British Gazetteer | 19 |

Juvenile Books.

| | |
|-----------------------------|----|
| Amey Herbert | 18 |
| Cleve Hall | 19 |
| Earl's Danger (Tae) | 19 |
| Experience of Life | 19 |
| Gertrude | 19 |
| Howitt's Boy's Country Book | 10 |
| " (Mary) Children's Year | 10 |
| Ivora | 19 |
| Katharine Ashton | 19 |
| Lanston Parsonage | 19 |
| Margaret Percival | 19 |

Medicine and Surgery.

| | |
|----------------------------------|----|
| Brodie's Psychological Inquiries | 4 |
| Bull's Hints to Mothers | 4 |
| " Menstruation of Children | 4 |
| Copland's Dictionary of Medicine | 6 |
| Curt's Invalid's Own Book | 7 |
| Holland's Mental Physiology | 9 |
| " Medical Notes and Reflect. | 9 |
| How to Nurse Sick Children | 10 |
| Kesteven's Domestic Medicine | 11 |
| L'pereira's Materia Medica | 17 |
| Reece's Medical Guide | 18 |
| Richardson's Cold-Water Cure | 19 |
| West on Diseases of Infancy | 24 |

Miscellaneous and General Literature.

| | |
|--|----|
| Bacon's (Lord) Works | 3 |
| " Lectures and Addresses | 22 |
| Defence of Eclipse of Faith | 7 |
| Eclipse of Faith | 7 |
| Greg's Political and Social Essays | 8 |
| Greyson's Select Correspondence | 8 |
| Gray's Evening Recitations | 8 |
| Hassall's Advertisements Detected, &c. | 8 |
| Haydn's Book of Dignities | 9 |
| Holland's Mental Physiology | 9 |
| Hooker's New Guides | 9 |
| Howitt's Rural Life of England | 10 |
| " Visiting Remarkable Places | 10 |
| Hutton's 100 Years Ago | 10 |
| Jameson's Commonplace-Book | 11 |
| Jeffrey's (Lord) Contributions | 11 |

| | |
|---|----|
| John's Lands of Silence and of Darkness | 11 |
| Last of the Old Squires | 16 |
| Macaulay's Crit. and Hist. Essays | 13 |
| " Speeches | 13 |
| Macintosh's Miscellaneous Works | 13 |
| Memoirs of a Maître d'Armes | 22 |
| Maitland's Church in the Catacombs | 14 |
| Martineau's Miscellanies | 14 |
| Moore's Church Cases | 16 |
| Printing: Its Origin &c. | 22 |
| Pycroft's English Reading | 17 |
| Rich's Comp. to Latin Dictionary | 18 |
| Riddle's Latin Dictionaries | 16 |
| Rowton's Debater | 18 |
| Seaward's Narrative of his Shipwreck | 19 |
| Sir Roger De Coverley | 19 |
| Smith's (Rev. Sydney) Works | 20 |
| " Common-place Books | 20 |
| " The Doctor &c. | 20 |
| Soerestre's Art of the Philosopher | 22 |
| " Confessions of a Working Man | 22 |
| Stephen's Essays | 20 |
| Stow's Training System | 21 |
| Thomson's Lives of the Thought | 21 |
| Townsend's State Trials | 21 |
| Willich's Popular Tables | 24 |
| Yonge's English-Greek Lexicon | 24 |
| " Latin Gradus | 24 |
| Zumpt's Latin Grammar | 24 |

Natural History general.

| | |
|--|----|
| Catlow's Popular Conchology | 6 |
| Ephemera and Young on the Salmon | 7 |
| Garra's Marvels of Insect | 8 |
| Gosse's Natural History of Jamaica | 6 |
| Kemp's Natural History of Creation | 22 |
| Kirby and Spence's Entomology | 11 |
| Lee's Elements of Natural History | 11 |
| Maudslayi's Natural History | 14 |
| Turton's Shells of the British Islands | 23 |
| Van der Hoeven's Zoology | 22 |
| Von Tschudi's Sketches in the Alps | 22 |
| Waterston's Essays on Natural Hist. | 24 |
| Yonath's The Dog | 24 |
| " The Horse | 24 |

1-Volume Encyclopedias and Dictionaries.

| | |
|---------------------------------------|----|
| Blaise's Rural Sports | 4 |
| Brande's Science, Literature, and Art | 4 |
| Copland's Dictionary of Medicine | 6 |
| Cressy's Civil Engineering | 8 |
| Gwilt's Architecture | 8 |
| Johston's Geographical Dictionary | 11 |
| Loudon's Agriculture | 12 |
| " Rural Architecture | 13 |
| " Gardening | 13 |
| " Plants | 13 |
| " Trees and Shrubs | 12 |
| M'Culloch's Geographical Dictionary | 14 |
| " Dictionary of Commerce | 14 |
| Murray's Encyclo. of Geography | 16 |
| Sharp's British Gazetteer | 16 |
| Ure's Dictionary of Arts, &c. | 23 |
| Webster's Domestic Economy | 24 |

Religious & Moral Works.

| | |
|--------------------------------------|----|
| Amy Herbert | 19 |
| Bloomfield's Greek Testament | 4 |
| Calvert's Wife's Manual | 6 |
| Cleve Hall | 19 |
| Conybeare's Essays | 6 |
| Conybeare and Hewson St. Paul | 6 |
| Cotton's Instruction in Christianity | 6 |
| Dale's Domestic Liturgy | 7 |
| Defence of Eclipse of Faith | 7 |
| Discipline | 7 |
| Earl's Daughters (The) | 19 |
| Eclipse of Faith | 7 |
| Englishman's Greek Concordance | 7 |
| " Heb. & Chald. Concord. | 7 |
| Experience (The) of Life | 19 |
| Gettrude | 19 |
| Harrison's Light of the Forge | 8 |
| Hook's Lectures on Passion Week | 9 |
| Horne's Introduction to Scriptures | 9 |
| " Abolishment of ditto | 10 |
| Hue's Christianity in China | 10 |
| Humphreys's Parables Illuminated | 10 |
| Ivors | 19 |
| Jameson's Sacred Legends | 11 |
| " Monastic Legends | 11 |
| " Legends of the Madonna | 11 |
| " Lectures on Female Employment | 11 |
| Jeremy Taylor's Works | 11 |
| Katherine Ashton | 11 |
| Knox's Fictional Life of Luther | 9 |
| Laneton Parsonage | 19 |
| Letters to my Unknown Friends | 11 |
| " on Happiness | 11 |
| Lynn Germanicus | 11 |
| Neosacht on Inspiration | 14 |
| Nagure's Rome | 14 |
| Maitland's Church in Catacombs | 14 |
| Margaret Percival | 19 |

| | |
|---|----|
| Martineau's Christian Life | 14 |
| " Hymns | 14 |
| Merivale's Christian Records | 16 |
| Milner's Church of Christ | 16 |
| Moore on the Use of the Body | 16 |
| " Soul and Body | 16 |
| " 'A Men and his Motives | 16 |
| Morisonism | 22 |
| Morning Clouds | 16 |
| Neale's Closing Scene | 16 |
| Rankin's Ferdinand & Maximilian | 18 |
| Readings for Lent | 18 |
| " Confirmation | 19 |
| Riddle's Household Prayers | 18 |
| Robinson's Lexicon to the Greek Testament | 18 |
| Saints our Example | 18 |
| Sermon in the Month | 19 |
| Sinclair's Journey of Life | 19 |
| Smith's (Sydney) Moral Philosophy | 20 |
| " G.V. Assyrian Prophecy | 20 |
| " (G.) Wesleyan Methodism | 19 |
| " (J.) St. Paul's Shipwreck | 20 |
| Southey's Life of Wesley | 20 |
| Stephen's Ecclesiastical Biography | 21 |
| Taylor's Loyals | 21 |
| " Wesley | 21 |
| Theologia Germanica | 5 |
| Thamb Bible (The) | 21 |
| Tomlin's Introduction to the Bible | 21 |
| Turner's Sacred History | 23 |
| Young's Christ of History | 24 |
| " Mystery | 24 |

Poetry and the Drama.

| | |
|-----------------------------------|----|
| Aikin's (Dr.) British Poets | 3 |
| Arnold's Poems | 3 |
| Baillie's (Joanna) Poetical Works | 3 |
| Calvert's Wife's Manual | 6 |
| De Vere's May Carols | 7 |
| Estcourt's Music of Creation | 7 |
| Fairy Family (The) | 7 |
| Goldsmith's Poems, illustrated | 11 |
| L. E. L.'s Poetical Works | 11 |
| Linwood's Anthologia Oxoniensis | 12 |
| Lyra Germanica | 5 |
| Macaulay's Lays of Ancient Rome | 13 |
| Mac Donald's Within and Without | 13 |
| " Poems | 13 |
| Montgomery's Poetical Works | 15 |
| Noore's Poetical Works | 15 |
| " Selections (illustrated) | 15 |
| " Lalla Rookh | 16 |
| " Irish Melodies | 16 |
| " Songs and Ballads | 15 |
| Reade's Poetical Works | 17 |
| Shakespeare, by Bowdler | 19 |
| Smith's Poetical Works | 20 |
| British Poets | 20 |
| Thomson's Seasons, illustrated | 21 |

Political Economy and Statistics.

| | |
|--------------------------------------|----|
| Dodd's Food of London | 7 |
| Greg's Political and Social Essays | 8 |
| Lane's Notes of a Traveller | 22 |
| M'Culloch's Geog. Statist. &c. Dict. | 14 |
| " Dictionary of Commerce | 14 |
| " London | 22 |
| Willich's Popular Tables | 24 |

The Sciences in general and Mathematics.

| | |
|--|----|
| Arago's Meteorological Essays | 3 |
| " Popular Astronomy | 3 |
| Bourne on the Screw Propeller | 4 |
| " C. Calculation of Steam-Engines | 4 |
| Boyd's Naval Cadet's Manual | 4 |
| Brande's Dictionary of Science, &c. | 4 |
| " Lectures on Organic Chemistry | 4 |
| Cressy's Civil Engineering | 6 |
| De la Roche's Geology of Cornwall, &c. | 7 |
| De la Rive's Electricity | 7 |
| Grove's Correla. of Physical Forces | 8 |
| Herschel's Outlines of Astronomy | 9 |
| Holland's Mental Physiology | 10 |
| Humboldt's Aspects of Nature | 10 |
| " Cosmos | 10 |
| Hunt on Light | 10 |
| Lardner's Cabinet Cyclopaedia | 12 |
| Marek's (Mrs.) Conversations | 15 |
| Morrell's Elements of Psychology | 16 |
| Moseley's Engineering & Architecture | 16 |
| Our Coal Fields and our Coal-Pits | 22 |
| Owen's Lectures on Comp. Anatomy | 17 |
| Persin on Polarized Light | 17 |
| Pecheil's Elements of Physics | 17 |
| Pecheil's Fossils of Cornwall, &c. | 17 |
| " Mineralogy | 17 |
| " Guide to Geology | 17 |
| Portlock's Geology of Londonderry | 17 |
| Powell's Unity of Worlds | 17 |
| Smeat's Electro-Metallurgy | 19 |
| Steam-Engines (The) | 19 |
| Wilson's Electric Telegraph | 22 |

Rural Sports.

| | |
|-------------------------------------|----|
| Baker's Rifle and Hound in Ceylon | 3 |
| Berkeley's Forests of France | 4 |
| Blaine's Dictionary of Sports | 6 |
| Cecil's Stable Practice | 7 |
| " Stud Farm | 7 |
| The Cricket-Field | 8 |
| Davy's Fishing Excursions, 2 Series | 7 |
| Ephemera on Angling | 7 |
| " The Book of the Salmon | 7 |
| Hawker's Young Sportsman | 9 |
| The Hunting-Field | 8 |
| Idle's Hints on Shooting | 10 |
| Pocket and the Stud | 8 |
| Thacker's Courser's Guide | 21 |
| Richardson's Horsemanship | 18 |
| Ronald's Fly-Fisher's Entomology | 19 |
| Stable Talk and Table Talk | 8 |
| Stoehenge on the Greyhound | 20 |
| The Stud, for Practical Purpose | 8 |

Veterinary Medicine, &c.

| | |
|----------------------------|----|
| Cecil's Stable Practice | 6 |
| " Stud Farm | 6 |
| Huntin'-Field (The) | 8 |
| Miles's Horse-Shoeing | 15 |
| " on the Horse's Foot | 15 |
| Pocket and the Stud | 8 |
| Practical Horsemanship | 8 |
| Richardson's Horsemanship | 18 |
| Stable Talk and Table Talk | 8 |
| Stud (The) | 8 |
| Yonath's The Dog | 24 |
| " The Horse | 24 |

Voyages and Travels.

| | |
|---|----|
| Auldjo's Ascent of Mont Blanc | 22 |
| Baines's Vaudouin of Piedmont | 22 |
| Baker's Wanderings in Ceylon | 22 |
| Barrow's Coastland of Tonar | 22 |
| Barth's African Travels | 3 |
| Berkeley's Forests of France | 4 |
| Burton's East Africa | 5 |
| " Medina and Mecca | 5 |
| Carlisle's Turkey and Greece | 6 |
| De Custine's Russia | 22 |
| Zethou | 22 |
| Ferguson's Swiss Travels | 22 |
| Flemish Interiors | 22 |
| Forester's Rambles in Norway | 22 |
| " Sardinia and Corsica | 8 |
| Gironi's Philippines | 22 |
| Groenouius's Corsica | 22 |
| Hailoran's Japan | 8 |
| Hill's Travels in Siberia | 9 |
| Hinchliff's Travels in the Alps | 9 |
| Hope's Brittany and the Bible | 22 |
| " Chase in Brittany | 22 |
| Howitt's Art-Student in Menich | 10 |
| " (W.) Victoria | 10 |
| Hue's Chinese Empire | 10 |
| Hue and Gabet's Tartary & Tibet | 22 |
| Hugh and Kennedy's Mount Blanc | 22 |
| Hughes's Australian Colonies | 22 |
| Humboldt's Aspects of Nature | 10 |
| Hurtich's Pictures from Caba | 22 |
| Hutchinson's African Exploration | 22 |
| Jameson's Canada | 22 |
| Jermann's St. Petersburg | 22 |
| Laing's Norway | 22 |
| " Notes of a Traveller | 22 |
| MacClure's North-West Passage | 17 |
| MacDonald's Voyage of the <i>Keserius</i> | 13 |
| Mason's Zulus of Natal | 22 |
| Miles's Rambles in Iceland | 15 |
| Obb's Quebec | 22 |
| Pfeiffer's Voyage round the World | 22 |
| " Second ditto | 17 |
| Scherzer's Central America | 16 |
| Schwarz's Nautilus | 19 |
| Snow's Tierra del Fuego | 20 |
| Spottiswood's Eastern Russia | 20 |
| Von Tempeky's Mexico and Guatemala | 24 |
| Wid's Vacations in Ireland | 24 |
| " United States and Canada | 24 |
| Werne's African Wanderings | 22 |
| Wilberforce's Brazil & Slave-Trade | 22 |

Works of Fiction.

| | |
|------------------------------|----|
| Cruikshank's Falstaff | 6 |
| Howitt's Talangetta | 10 |
| Macdonald's Villa Verocchio | 13 |
| McNeill's Confidence-Men | 19 |
| Moore's Epistles | 19 |
| Sir Roger De Coverley | 19 |
| Sketches (The), Three Tales | 19 |
| Southey's The Doctor &c. | 20 |
| Trollope's Barchester Towers | 23 |
| " Waverley | 23 |

ALPHABETICAL CATALOGUE

of

NEW WORKS and NEW EDITIONS

PUBLISHED BY

Messrs. LONGMAN, BROWN, GREEN, LONGMANS, and ROBERTS,

PATERNOSTER ROW, LONDON.

Miss Acton's Modern Cookery for Private Families, reduced to a System of Easy Practice in a Series of carefully-tested Receipts, in which the Principles of Baron Liebig and other eminent Writers have been as much as possible applied and explained. Newly-revised and enlarged Edition; with 8 Plates, comprising 27 Figures, and 150 Woodcuts. Fcp. 8vo. 7s. 6d.

Acton's English Bread-Book for Domestic Use, adapted to Families of every grade: Containing the plainest and most minute Instructions to the Learner, and Practical Receipts for many varieties of Bread; with Notices of the present System of Adulteration and its Consequences, and of the Improved Baking Processes and Institutions established Abroad. Fcp. 8vo. price 4s. 6d. cloth.

Aikin.—Select Works of the British Poets, from Ben Jonson to Beattie. With Biographical and Critical Prefaces by Dr. AIKIN. New Edition, with Supplement by LUCY AIKIN; consisting of additional Selections from more recent Poets. 8vo. price 18s.

Arago (F.).—Biographies of Distinguished Scientific Men. Translated by Admiral W. H. SMYTH, D.C.L., F.R.S., &c.; the Rev. BADEN POWELL, M.A.; and ROBERT GRANT, M.A., F.R.A.S. 8vo. 18s.

Arago's Meteorological Essays. With an Introduction by BARON HUMBERT. Translated under the superintendence of Lieut.-Colonel E. SABINE, R.A., Treasurer and V.P.R.S. 8vo. 18s.

Arago's Popular Astronomy. Translated and edited by Admiral W. H. SMYTH, D.C.L., F.R.S.; and ROBERT GRANT, M.A., F.R.A.S. In Two Volumes. Vol. I. 8vo. with Plates and Woodcuts, 21s.

Arnold.—Poems. By Matthew Arnold. Third Edition of the *First Series*. Fcp. 8vo. price 5s. 6d.

Arnold.—Poems. By Matthew Arnold. Second Series, about one-third new; the rest finally selected from the Volumes of 1849 and 1852, now withdrawn. Fcp. 8vo. price 5s.

Lord Bacon's Works. A New Edition, revised and elucidated; and enlarged by the addition of many pieces not printed before Collected and Edited by ROBERT LESLIE ELLIS, M.A., Fellow of Trinity College, Cambridge; JAMES SPEDDING, M.A. of Trinity College, Cambridge; and DOUGLAS DENON HEATH, Esq., Barrister-at-Law, and late Fellow of Trinity College, Cambridge.—The publication has commenced with the Division of the *Philosophical Works*, to be completed in 5 vols., of which Vols. I. to III. in 8vo., price 18s. each, are now ready. Vols. IV. and V. are in the press.

Joanna Baillie's Dramatic and Poetical Works: Comprising the Plays of the Passions, Miscellaneous Dramas, Metrical Legends, Fugitive Pieces, and *Ahalya Bae*. Second Edition, with a Life of Joanna Baillie, Portrait, and Vignette. Square crown 8vo. 21s. cloth; or 42s. bound in morocco by Hayday.

Baker.—The Rifle and the Hound in Ceylon. By S. W. BAKER, Esq. New Edition, with 13 Illustrations engraved on Wood. Fcp. 8vo. 4s. 6d.

Baker.—Eight Years' Wanderings in Ceylon. By S. W. BAKER, Esq. With 6 coloured Plates. 8vo. price 15s.

Barth.—Travels and Discoveries in North and Central Africa: Being the Journal of an Expedition undertaken under the auspices of Her Britannic Majesty's Government in the Years 1849—1855. By HENRY BARTH, Ph.D., D.C.L., Fellow of the Royal Geographical and Asiatic Societies, &c. Vols. I. to III., with 11 Maps, 100 Engravings on Wood, and 36 Illustrations in tinted Lithography, price 63s.—Vols. IV. and V., completing the work, are in the press.

Bayldon's Art of Valuing Rents and Tillages, and Claims of Tenants upon Quitting Farms, at both Michaelmas and Lady-Day; as revised by Mr. DONALDSON. *Seventh Edition*, enlarged and adapted to the Present Time: With the Principles and Mode of Valuing Land and other Property for Parochial Assessment and Enfranchisement of Copyholds, under the recent Acts of Parliament. By ROBERT BAKER, Land-Agent and Valuer. 8vo. 10s. 6d.

A Month in the Forests of France. By the Hon. GRANTLEY F. BERKELEY, Author of *Reminiscences of a Huntsman*. 8vo. with 2 Etchings by John Leech (1 coloured). [Nearly ready.]

Black's Practical Treatise on Brewing, based on Chemical and Economical Principles: With Formulæ for Public Brewers, and Instructions for Private Families. New Edition, with Additions. 8vo. 10s. 6d.

Blaine's Encyclopædia of Rural Sports; or, a complete Account, Historical, Practical, and Descriptive, of Hunting, Shooting, Fishing, Racing, and other Field Sports and Athletic Amusements of the present day. New Edition, revised by HARRY HIEROVER, EPHEMERA, and Mr. A. GRAHAM. With upwards of 600 Woodcuts. 8vo. 50s.

Blair's Chronological and Historical Tables, from the Creation to the Present Time: With Additions and Corrections from the most authentic Writers; including the Computation of St. Paul, as connecting the Period from the Exode to the Temple. Under the revision of SIR HENRY ELLIS, K.H. Imperial 8vo. 31s. 6d. half-morocco.

Bloomfield. — The Greek Testament, with copious English Notes, Critical, Philological, and Explanatory. Especially adapted to the use of Theological Students and Ministers. By the Rev. S. T. BLOOMFIELD, D.D., F.S.A. Ninth Edition, revised. 2 vols. 8vo. with Map, price £2. 8s.

Dr. Bloomfield's College and School Edition of the Greek Testament: With brief English Notes, chiefly Philological and Explanatory, especially formed for use in Colleges and the Public Schools. Seventh Edition, improved; with Map and Index. Fcp. 8vo. 7s. 6d.

Dr. Bloomfield's College and School Lexicon to the Greek Testament. New Edition, carefully revised. Fcp. 8vo. price 10s. 6d. cloth.

Bourne. — A Treatise on the Steam-Engine, in its Application to Mines, Mills, Steam-Navigation, and Railways. By the Artisan Club. Edited by JOHN BOURNE, C.E. New Edition; with 33 Steel Plates and 349 Wood Engravings. 4to. price 27s.

Bourne's Catechism of the Steam-Engine in its various Applications to Mines, Mills, Steam-Navigation, Railways, and Agriculture: With Practical Instructions for the Manufacture and Management of Engines of every class. Fourth Edition, enlarged; with 89 Woodcuts. Fcp. 8vo. 6s.

Bourne. — A Treatise on the Screw Propeller: With various Suggestions of Improvement. By JOHN BOURNE, C.E. New Edition, thoroughly revised and corrected. With 20 large Plates and numerous Woodcuts. 4to. price 38s.

Boyd. — A Manual for Naval Cadets. Published with the sanction and approval of the Lords Commissioners of the Admiralty. By JOHN McNEILL BOYD, Captain, R.N. With Compass-Signals in Colours, and 236 Woodcuts. Fcp. 8vo. 10s. 6d.

Brande. — A Dictionary of Science, Literature, and Art: Comprising the History, Description, and Scientific Principles of every Branch of Human Knowledge; with the Derivation and Definition of all the Terms in general use. Edited by W. T. BRANDE, F.R.S.L. and E.; assisted by DR. J. CAUVIN. Third Edition, revised and corrected; with numerous Woodcuts. 8vo. 60s.

Professor Brande's Lectures on Organic Chemistry, as applied to Manufactures; including Dyeing, Bleaching, Calico-Printing, Sugar-Manufacture, the Preservation of Wood, Tanning, &c.; delivered before the Members of the Royal Institution. Arranged by permission from the Lecturer's Notes by J. SCOFFERN, M.B. Fcp. 8vo. with Woodcuts, price 7s. 6d.

Brewer. — An Atlas of History and Geography, from the Commencement of the Christian Era to the Present Time: Comprising a Series of Sixteen coloured Maps, arranged in Chronological Order, with Illustrative Memoirs. By the Rev J. S. BREWER, M.A., Professor of English History and Literature in King's College, London. Second Edition, revised and corrected. Royal 8vo. 12s. 6d. half-bound.

Brodie. — Psychological Inquiries, in a Series of Essays intended to illustrate the Influence of the Physical Organisation on the Mental Faculties. By SIR BENJAMIN C. BRODIE, Bart. Third Edition. Fcp. 8vo. 5s.

Bull. — The Maternal Management of Children in Health and Disease. By T. BULL, M.D., Member of the Royal College of Physicians; formerly Physician-Accoucheur to the Finsbury Midwifery Institution. New Edition. Fcp. 8vo. 5s.

Dr. T. Bull's Hints to Mothers on the Management of their Health during the Period of Pregnancy and in the Lying-in Room: With an Exposure of Popular Errors in connexion with those subjects, &c.; and Hints upon Nursing. New Edition. Fcp. 8vo. 5s.

Buckingham.—Autobiography of James Silk Buckingham: Including his Voyages, Travels, Adventures, Speculations, Successes and Failures, frankly and faithfully narrated; with Characteristic Sketches of Public Men. Vols. I. and II. post 8vo. 21s.

Bunsen.—Christianity and Mankind, their Beginnings and Prospects. By CHRISTIAN CHARLES JOSIAS BUNSEN, D.D., D.C.L., D.Ph. Being a New Edition, corrected, remodelled, and extended, of *Hippolytus and his Age*. 7 vols. 8vo. £5. 5s.

. This Second Edition of the *Hippolytus* is composed of three distinct works, which may be had separately, as follows:—

1. *Hippolytus and his Age; or, the Beginnings and Prospects of Christianity*. 2 vols. 8vo. price £1. 10s.
2. *Outline of the Philosophy of Universal History applied to Language and Religion: Containing an Account of the Alphabetical Conferences*. 2 vols. 8vo. price £1. 13s.
3. *Analecta Ante-Nicæna*. 3 vols. 8vo. price £2. 2s.

Bunsen.—Lyra Germanica: Hymns for the Sundays and chief Festivals of the Christian Year. Translated from the German by CATHERINE WINKWORTH. Third Edition. Fcp. 8vo. 5s.

. This selection of German Hymns has been made from a collection published in Germany by the Chevalier BUNSEN; and forms a companion volume to

Theologia Germanica: Which setteth forth many fair lineaments of Divine Truth, and saith very lofty and lovely things touching a Perfect Life. Translated by SUSANNA WINKWORTH. With a Preface by the Rev. CHARLES KINGSLEY; and a Letter by Chevalier BUNSEN. Third Edition. Fcp. 8vo. 5s.

Bunsen.—Egypt's Place in Universal History: An Historical Investigation, in Five Books. By C. C. J. BUNSEN, D.D., D.C.L., D.Ph. Translated from the German by C. H. COTTELL, Esq., M.A. With many Illustrations. Vol. I. 8vo. 28s.; Vol. II. 8vo. 30s.

Burton (J. H.).—The History of Scotland from the Revolution to the Extinction of the Last Jacobite Insurrection (1689–1748). By JOHN HILL BURTON. 2 vols. 8vo. 26s.

Bishop S. Butler's General Atlas of Modern and Ancient Geography; comprising Fifty-two full-coloured Maps; with complete Indices. New Edition, nearly all re-engraved, enlarged, and greatly improved. Edited by the Author's Son. Royal 4to. 24s. half-bound.

Separately { The Modern Atlas of 28 full-coloured Maps. Royal 8vo. price 13s.
The Ancient Atlas of 24 full-coloured Maps. Royal 8vo. price 12s.

Bishop S. Butler's Sketch of Modern and Ancient Geography. New Edition, thoroughly revised, with such Alterations introduced as continually progressive Discoveries and the latest Information have rendered necessary. Post 8vo. price 7s. 6d.

Burton.—First Footsteps in East Africa; or, an Exploration of Harar. By RICHARD F. BURTON, Captain, Bombay Army. With Maps and coloured Plates. 8vo. 18s.

Burton.—Personal Narrative of a Pilgrimage to El Medinah and Meccah. By RICHARD F. BURTON, Captain, Bombay Army. Second Edition, revised; with coloured Plates and Woodcuts. 2 vols. crown 8vo. price 24s.

The Cabinet Lawyer: A Popular Digest of the Laws of England, Civil and Criminal; with a Dictionary of Law Terms, Maxims, Statutes, and Judicial Antiquities; Correct Tables of Assessed Taxes, Stamp Duties, Excise Licences, and Post-Horae Duties; Post-Office Regulations; and Prison Discipline. 17th Edition, comprising the Public Acts of the Session 1857. Fcp. 8vo. 10s. 6d.

The Cabinet Gazetteer: A Popular Exposition of All the Countries of the World: their Government, Population, Revenues, Commerce, and Industries; Agricultural, Manufactured, and Mineral Products; Religion, Laws, Manners, and Social State; With brief Notices of their History and Antiquities. By the Author of *The Cabinet Lawyer*. Fcp. 8vo. 10s. 6d. cloth; or 13s. bound in calf.

"The author has neglected no modern sources of information, and all his short, succinct, and neat descriptions of the different places are quite conformable to present knowledge. Sarawak, for example, in Borneo, is not omitted, and of San Francisco there is quite a detailed description. The work is compiled with considerable care, and in the 912 pages that it contains there is a vast amount of geographical and topographical information pleasantly condensed. The *Cabinet Gazetteer*, though not intended to supersede more elaborate works, will, to some extent, have that effect; but it will be sure to find a large and permanent circulation of its own."

ECONOMIST.

Calendar of English State Papers, Domestic Series, of the Reigns of Edward VI., Mary, Elizabeth, 1547–1580, preserved in the State Paper Department of Her Majesty's Public Record Office. Edited by ROBERT LEMON, Esq., F.S.A., under the direction of the Master of the Rolls, and with the sanction of Her Majesty's Secretary of State for the Home Department. Imperial 8vo. 15s.

Calendar of English State Papers, Domestic Series, of the Reign of James I., 1603–1610 (comprising the Papers relating to the Gunpowder Plot), preserved in the State Paper Department of H.M. Public Record Office. Edited by MARY ANNE EVERETT GREEN, Author of *The Lives of the Princesses of England*, &c., under the direction of the Master of the Rolls, and with the sanction of H.M. Secretary of State for the Home Department. Imperial 8vo. 15s.

Calvert.—*The Wife's Manual; or, Prayers, Thoughts, and Songs on Several Occasions of a Matron's Life.* By the Rev. W. CALVERT, M.A. Ornamented from Designs by the Author in the style of *Queen Elizabeth's Prayer-Book.* Second Edition. Crown 8vo. 10s. 6d.

Carlisle (Lord).—*A Diary in Turkish and Greek Waters.* By the Right Hon. the EARL OF CARLISLE. Fifth Edition. Post 8vo. price 10s. 6d.

Catlow.—*Popular Conchology; or, the Shell Cabinet arranged according to the Modern System: With a detailed Account of the Animals, and a complete Descriptive List of the Families and Genera of Recent and Fossil Shells.* By AGNES CATLOW. Second Edition, much improved; with 405 Woodcut Illustrations. Post 8vo. price 14s.

Cecil.—*The Stud Farm; or, Hints on Breeding Horses for the Turf, the Chase, and the Road.* Addressed to Breeders of Race-Horses and Hunters, Landed Proprietors, and especially to Tenant Farmers. By CECIL. Fcp. 8vo. with Frontispiece, 5s.

Cecil's Stable Practice; or, Hints on Training for the Turf, the Chase, and the Road; with Observations on Racing and Hunting, Wasting, Race-Riding, and Handicapping: Addressed to Owners of Racers, Hunters, and other Horses, and to all who are concerned in Racing, Steeple-Chasing, and Fox-Hunting. Fcp. 8vo. with Plate, price 5s. half-bound.

Chapman.—*History of Gustavus Adolphus, and of the Thirty Years' War up to the King's Death: With some Account of its Conclusion by the Peace of Westphalia, in 1648.* By B. CHAPMAN, M.A., Vicar of Letherhead. 8vo. with Plans, 12s. 6d.

Chevreul On the Harmony and Contrast of Colours, and their Applications to the Arts: Including Painting, Interior Decoration, Tapestries, Carpets, Mosaics, Coloured Glazing, Paper-Staining, Calico-Printing, Letterpress-Printing, Map-Colouring, Dress, Landscape and Flower-Gardening, &c. &c. Translated by CHARLES MARTEL. Second Edition; with 4 Plates. Crown 8vo. price 10s. 6d.

Connolly.—*History of the Royal Sappers and Miners: Including the Services of the Corps in the Crimea and at the Siege of Sebastopol.* By T. W. J. CONNOLLY, Quartermaster of the Royal Engineers. Second Edition, revised and enlarged; with 17 coloured plates. 2 vols. 8vo. price 30s.

Conybeare and Howson.—*The Life and Epistles of Saint Paul: Comprising a complete Biography of the Apostle, and a Translation of his Epistles inserted in Chronological Order.* By the Rev. W. J. CONYBEARE, M.A.; and the Rev. J. S. HOWSON, M.A. Second Edition, revised and corrected; with several Maps and Woodcuts, and 4 Plates. 2 vols. square crown 8vo. 31s. 6d. cloth.

. The Original Edition, with more numerous Illustrations, in 2 vols. 4to. price 48s.—may also be had.

Conybeare.—*Essays, Ecclesiastical and Social: Reprinted, with Additions, from the Edinburgh Review.* By the Rev. W. J. CONYBEARE, M.A., late Fellow of Trinity College, Cambridge. 8vo. 12s.

Dr. Copland's Dictionary of Practical Medicine: Comprising General Pathology, the Nature and Treatment of Diseases, Morbid Structures, and the Disorders especially incidental to Climates, to Sex, and to the different Epochs of Life; with numerous approved Formulae of the Medicines recommended. Vols. I. and II. 8vo. price £3; and Parts X. to XVIII. 4s. 6d. each.

. Part XIX., completing the work, is nearly ready.

Cotton.—*Instructions in the Doctrine and Practice of Christianity.* Intended chiefly as an Introduction to Confirmation. By G. E. L. COTTON, M.A. 18mo. 2s. 6d.

Cresy's Encyclopædia of Civil Engineering, Historical, Theoretical, and Practical. Illustrated by upwards of 3,000 Woodcuts. Second Edition, revised and brought down to the Present Time in a Supplement, comprising Metropolitan Water-Supply, Drainage of Towns, Railways, Cubical Proportion, Brick and Iron Construction, Iron Screw Piles, Tubular Bridges, &c. 8vo. 63s. cloth.—THE SUPPLEMENT separately, price 10s. 6d. cloth.

The Cricket-Field; or, the Science and History of the Game of Cricket. By the Author of *Principles of Scientific Batting.* Second Edition, greatly improved; with Plates and Woodcuts. Fcp. 8vo. price 5s.

Crosse.—*Memorials, Scientific and Literary, of Andrew Crosse, the Electrician.* Edited by Mrs. CROSSE. Post 8vo. 9s. 6d.

Cruikshank.—*The Life of Sir John Falstaff, illustrated by George Cruikshank.* With a Biography of the Knight, from authentic sources, by ROBERT B. BROUGH, Esq. Royal 8vo.—In course of publication monthly, and to be completed in 10 Numbers, each containing 2 Plates, price 1s. The first 6 Numbers are now ready.

Lady Cust's Invalid's Book.—The Invalid's Own Book: A Collection of Recipes from various Books and various Countries. By the Honourable LADY CUST. *Second Edition.* Fcp. 8vo. price 2s. 6d.

Dale.—The Domestic Liturgy and Family Chaplain, in Two Parts: PART I. Church Services adapted for Domestic Use, with Prayers for Every Day of the Week, selected from the Book of Common Prayer; PART II. an appropriate Sermon for Every Sunday in the Year. By the Rev. THOMAS DALE, M.A., Canon Residentiary of St. Paul's. *Second Edition.* Post 4to. 21s. cloth; 31s. 6d. calf; or £2. 10s. morocco.

Separately { THE FAMILY CHAPLAIN, 12s.
THE DOMESTIC LITURGY, 10s. 6d.

Davy (Dr. J.)—The Angler and his Friend; or, Piscatory Colloquies and Fishing Excursions. By JOHN DAVY, M.D., F.R.S., &c. Fcp. 8vo. price 6s.

The Angler in the Lake District: Or, Piscatory Colloquies and Fishing Excursions in Westmoreland and Cumberland. By JOHN DAVY, M.D., F.R.S. Fcp. 8vo. 6s. 6d.

Delabèche.—Report on the Geology of Cornwall, Devon, and West Somerset. By SIR H. T. DELABÈCHE, F.R.S. With Maps, Plates, and Woodcuts. 8vo. price 14s.

De la Rive.—A Treatise on Electricity in Theory and Practice. By A. DELA RIVE, Professor in the Academy of Geneva. Translated for the Author by C. V. WALKER, F.R.S. *In Three Volumes*; with numerous Woodcuts. Vol. I. 8vo. price 18s.; Vol. II. price 28s.—Vol. III. is in the press.

De Vere.—May Carols. By ANBREY DE VERE, Author of *The Search after Prosperine*, &c. Fcp. 8vo. 5s.

Discipline. By the Author of "Letters to my Unknown Friends," &c. *Second Edition, enlarged.* 18mo. price 2s. 6d.

Dodd.—The Food of London: A Sketch of the chief Varieties, Sources of Supply, probable Quantities, Modes of Arrival, Processes of Manufacture, suspected Adulteration, and Machinery of Distribution of the Food for a Community of Two Millions and a Half. By GEORGE DODD, Author of *British Manufactures*, &c. Post 8vo. 10s. 6d.

Estcourt.—Music the Voice of Harmony in Creation. Selected and arranged by Mary Jane Estcourt. Fcp. 8vo. 7s. 6d.

The Eclipse of Faith; or, a Visit to a Religious Sceptic. *8th Edition.* Fcp. 8vo. 5s.

Defence of The Eclipse of Faith, by its Author: Being a Rejoinder to Professor Newman's *Reply*: Including a full Examination of that Writer's Criticism on the Character of Christ; and a Chapter on the Aspects and Pretensions of Modern Deism. *Second Edition, revised.* Post 8vo. 5s. 6d.

The Englishman's Greek Concordance of the New Testament: Being an Attempt at a Verbal Connexion between the Greek and the English Texts; including a Concordance to the Proper Names, with Indexes, Greek-English and English-Greek. *New Edition,* with a new Index. Royal 8vo. price 42s.

The Englishman's Hebrew and Chaldean Concordance of the Old Testament: Being an Attempt at a Verbal Connexion between the Original and the English Translations; with Indexes, a List of the Proper Names and their Occurrences, &c. 2 vols. royal 8vo. £3. 13s. 6d.; large paper, £4. 14s. 6d.

Ephemera's Handbook of Angling; teaching Fly-Fishing, Trolling, Bottom-Fishing, Salmon-Fishing: With the Natural History of River-Fish, and the best Modes of Catching them. *Third Edition, corrected and improved;* with Woodcuts. Fcp. 8vo. 5s.

Ephemera.—The Book of the Salmon: Comprising the Theory, Principles, and Practice of Fly-Fishing for Salmon; Lists of good Salmon Flies for every good River in the Empire; the Natural History of the Salmon, its Habits described, and the best way of artificially Breeding it. By EPHEMERA; assisted by ANDREW YOUNG. Fcp. 8vo. with coloured Plates, price 14s.

Fairbairn.—Useful Information for Engineers: Being a Series of Lectures delivered to the Working Engineers of Yorkshire and Lancashire. With Appendices, containing the Results of Experimental Inquiries into the Strength of Materials, the Causes of Boiler Explosions, &c. By WILLIAM FAIRBAIRN, F.R.S., F.G.S. *Second Edition*; with numerous Plates and Woodcuts. Crown 8vo. price 10s. 6d.

The Fairy Family: A Series of Ballads and Metrical Tales illustrating the Fairy Mythology of Europe. With Frontispiece and Pictorial Title. Crown 8vo. 10s. 6d.

Flemish Interiors. By the Writer of *A Glance behind the Grilles of Religious Houses in France.* Fcp. 8vo. 7s. 6d.

Forester.—*Travels in the Islands of Corsica and Sardinia.* By THOMAS FORESTER, Author of *Rambles in Norway*. With numerous coloured Illustrations and Woodcuts, from Sketches made during the Tour by Lieutenant-Colonel M. A. BIDDULPH, R.A. Imperial 8vo. [In the press.]

Garratt.—*Marvels and Mysteries of Instinct; or, Curiosities of Animal Life.* By GEORGE GARRATT. *Second Edition*, revised and improved; with a Frontispiece. Fcp. 8vo. 4s. 6d.

Gilbart.—*A Practical Treatise on Banking.* By JAMES WILLIAM GILBART, F.R.S., General Manager of the London and Westminster Bank. *Sixth Edition*, revised and enlarged. 2 vols. 12mo. Portrait, 16s.

Gilbart.—*Logic for the Million: A Familiar Exposition of the Art of Reasoning.* By J. W. GILBART, F.R.S. 5th Edition; with Portrait of the Author. 12mo. 3s. 6d.

The Poetical Works of Oliver Goldsmith. Edited by BOLTON CORNEY, Esq. Illustrated by Wood Engravings, from Designs by Members of the Etching Club. Square crown 8vo. cloth, 21s.; morocco, £1. 16s.

Gosse.—*A Naturalist's Sojourn in Jamaica.* By P. H. GOSSE, Esq. With Plates. Post 8vo. price 14s.

Green.—*Lives of the Princesses of England.* By MRS. MARY ANNE EVERETT GREEN, Editor of the *Letters of Royal and Illustrious Ladies*. With numerous Portraits. Complete in 6 vols. post 8vo. price 10s. 6d. each.—Any Volume may be had separately to complete sets.

Mr. W. R. Greg's Essays on Political and Social Science, contributed chiefly to the *Edinburgh Review*. 2 vols. 8vo. price 24s.

Greyson.—*Selections from the Correspondence of R. E. H. GREYSON, Esq.* Edited by the Author of *The Eclipse of Faith*. 2 vols. fcp. 8vo. price 12s.

Grove.—*The Correlation of Physical Forces.* By W. R. GROVE, Q.C., M.A., F.R.S., &c. *Third Edition*. 8vo. price 7s.

Gurney.—*St. Louis and Henri IV.: Being a Second Series of Historical Sketches.* By the Rev. JOHN H. GURNEY, M.A., Rector of St. Mary's, Marylebone. Fcp. 8vo. 6s.

Evening Recreations; or, Samples from the Lecture-Room. Edited by the Rev. J. H. GURNEY, M.A. Crown 8vo. 5s.

Gwilt's Encyclopædia of Architecture, Historical, Theoretical, and Practical. By JOSEPH GWILT. With more than 1,000 Wood Engravings, from Designs by J. S. GWILT. *Third Edition*. 8vo. 42s.

Halloran.—*Eight Months' Journal kept on board one of H.M. Sloops of War, during Visits to Loochoo, Japan, and Pootoo.* By ALFRED L. HALLORAN, Master, R.N. With Etchings and Woodcuts. Post 8vo. 7s. 6d.

Hare (Archdeacon).—*The Life of Luther*, in Forty-eight Historical Engravings. By GUSTAV KÖNIG. With Explanations by ARCHDEACON HARE and SUSANNA WINKWORTH. Fcp. 4to. price 28s.

Harford.—*Life of Michael Angelo Buonarroti: With Translations of many of his Poems and Letters; also Memoirs of Savonarola, Raphael, and Vittoria Colonna.* By JOHN S. HARFORD, Esq., D.C.L., F.R.S., Member of the Academy of Painting of St. Luke, at Rome, and of the Roman Archaeological Society. With Portrait and Plates. 2 vols. 8vo. 25s.

Illustrations, Architectural and Pictorial, of the Genius of Michael Angelo Buonarroti. With Descriptions of the Plates, by the Commendatore CANINA; C. R. COCKEBELL, Esq., R.A.; and J. S. HARFORD, Esq., D.C.L., F.R.S. Folio, 73s. 6d. half-bound.

Harrison.—*The Light of the Forge; or, Counsels drawn from the Sick-Bed of E. M.* By the Rev. W. HARRISON, M.A., Domestic Chaplain to H.R.H. the Duchess of Cambridge. Fcp. 8vo. price 5s.

Harry Hieover.—*Stable Talk and Table Talk; or, Spectacles for Young Sportsmen.* By HARRY HIEOVER. *New Edition*, 2 vols. 8vo. with Portrait, price 24s.

Harry Hieover.—*The Hunting-Field.* By HARRY HIEOVER. With Two Plates. Fcp. 8vo. 5s. half-bound.

Harry Hieover.—*Practical Horsemanship.* By HARRY HIEOVER. *Second Edition*; with 2 Plates. Fcp. 8vo. 5s. half-bound.

Harry Hieover.—*The Pocket and the Stud; or, Practical Hints on the Management of the Stable.* By HARRY HIEOVER. *Second Edition*; with Portrait of the Author. Fcp. 8vo. price 5s. half-bound.

Harry Hieover.—*The Stud, for Practical Purposes and Practical Men: Being a Guide to the Choice of a Horse for use more than for show.* By HARRY HIEOVER. With 2 Plates. Fcp. 8vo. price 5s. half-bound.

Hassall.—Adulterations Detected; or, Plain Instructions for the Discovery of Frauds in Food and Medicine. By ARTHUR HILL HASSALL, M.D. Lond., Analyst of *The Lancet* Sanitary Commission, and Author of the Reports of that Commission published under the title of *Food and its Adulterations* (which may also be had, in 8vo. price 28s.) With 225 Illustrations, engraved on Wood. Crown 8vo. 17s. 6d.

Hassall.—A History of the British Fresh Water Algae: Including Descriptions of the Desmidiæ and Diatomaceæ. With upwards of One Hundred Plates of Figures, illustrating the various Species. By ARTHUR HILL HASSALL, M.D., Author of *Microscopic Anatomy of the Human Body*, &c. 2 vols. 8vo. with 103 Plates, price £1. 15s.

Col. Hawker's Instructions to Young Sportsmen in all that relates to Guns and Shooting. 10th Edition, revised and brought down to the Present Time, by the Author's Son, Major P. W. L. HAWKER. With a Portrait of the Author, and numerous Plates and Woodcuts. 8vo. 21s.

Haydn's Book of Dignities: Containing Rolls of the Official Personages of the British Empire, Civil, Ecclesiastical, Judicial, Military, Naval, and Municipal, from the Earliest Periods to the Present Time. Together with the Sovereigns of Europe, from the Foundation of their respective States; the Peerage and Nobility of Great Britain; &c. Being a New Edition, improved and continued, of Beaton's Political Index. 8vo. 25s. half-bound.

Sir John Herschel.—Essays from the Edinburgh and Quarterly Reviews, with Addresses and other Pieces. By SIR JOHN F. W. HERSCHEL, Bart., K.H., M.A. 8vo. price 18s.

Sir John Herschel.—Outlines of Astronomy. By SIR JOHN F. W. HERSCHEL, Bart., K.H., M.A. New Edition; with Plates and Woodcuts. 8vo. price 18s.

Hill.—Travels in Siberia. By S. S. Hill, Esq., Author of *Travels on the Shores of the Baltic*. With a large Map of European and Asiatic Russia. 2 vols. post 8vo. 24s.

Hinchliff.—Summer Months among the Alps: With the Ascent of Monte Rosa. By THOMAS W. HINCHLIFF, of Lincoln's Inn, Barrister-at-Law. With 4 tinted Views and 3 Maps. Post 8vo. price 10s. 6d.

Hints on Etiquette and the Usages of Society: With a Glance at Bad Habits. New Edition, revised (with Additions) by a Lady of Rank. Fcp. 8vo. price Half-a-Crown.

Holland.—Medical Notes and Reflections. By SIR HENRY HOLLAND, Bart., M.D., F.R.S., &c., Physician in Ordinary to the Queen and Prince Albert. Third Edition. 8vo. 18s.

Holland.—Chapters on Mental Physiology. By SIR HENRY HOLLAND, Bart., F.R.S., &c. Founded chiefly on Chapters contained in the First and Second Editions of *Medical Notes and Reflections* by the same Author. 8vo. price 10s. 6d.

Hook.—The Last Days of Our Lord's Ministry: A Course of Lectures on the principal Events of Passion Week. By the Rev. W. F. HOOK, D.D. New Edition. Fcp. 8vo. price 6s.

Hooker.—Kew Gardens; or, a Popular Guide to the Royal Botanic Gardens of Kew. By SIR WILLIAM JACKSON HOOKER, K.H., &c., Director. New Edition; with many Woodcuts. 16mo. price Sixpence.

Hooker.—Museum of Economic Botany; or, a Popular Guide to the Useful and Remarkable Vegetable Products of the Museum in the Royal Gardens of Kew. By SIR W. J. HOOKER, K.H., &c., Director. With 29 Woodcuts. 16mo. price 1s.

Hooker and Arnott.—The British Flora: comprising the Phænogamous or Flowering Plants, and the Ferns. Seventh Edition, with Additions and Corrections; and numerous Figures illustrative of the Umbelliferous Plants, the Composite Plants, the Grasses, and the Ferns. By SIR W. J. HOOKER, F.R.A. and L.S., &c.; and G. A. WALKER-ARNOTT, LL.D., F.L.S. 12mo. with 12 Plates, price 14s.; with the Plates coloured, price 21s.

Horne's Introduction to the Critical Study and Knowledge of the Holy Scriptures. Tenth Edition, revised, corrected, and brought down to the present time. Edited by the Rev. T. HARTWELL HORNE, B.D. (the Author); the Rev. SAMUEL DAVIDSON, D.D. of the University of Halle, and LL.D.; and S. PRIDEAUX TREGELLES, LL.D. With 4 Maps and 22 Vignettes and Facsimiles. 4 vols. 8vo. £3. 13s. 6d.

* * * The Four Volumes may also be had *separately* as follows:—

VOL. I.—A Summary of the Evidence for the Genuineness, Authenticity, Uncontested Preservation, and Inspiration of the Holy Scriptures. By the Rev. T. H. Horne, B.D. 8vo. 15s.

VOL. II.—The Text of the Old Testament considered: With a Treatise on Sacred Interpretation; and a brief Introduction to the Old Testament Books and the Apocrypha. By S. Davidson, D.D. (Halle) and LL.D. 8vo. 25s.

VOL. III.—A Summary of Biblical Geography and Antiquities. By the Rev. T. H. Horne, B.D. 8vo. 18s.

VOL. IV.—An Introduction to the Textual Criticism of the New Testament. By the Rev. T. H. Horne, B.D. The Critical Part re-written, and the remainder revised and edited by S. P. Tregelles, LL.D. 8vo. 18s.

Horne.—*A Compendious Introduction to the Study of the Bible.* By the Rev. T. HARTWELL HORNE, B.D. New Edition, with Maps and Illustrations. 12mo. 9s.

Hoskyns.—*Talpa; or, the Chronicles of a Clay Farm: An Agricultural Fragment.* By CHANDOS WREN HOSKYNs, Esq. Fourth Edition. With 24 Woodcuts from the original Designs by GEORGE CRUIKSHANK. 16mo. price 5s. 6d.

How to Nurse Sick Children: Intended especially as a Help to the Nurses in the Hospital for Sick Children; but containing Directions of service to all who have the charge of the Young. Fcp. 8vo. 1s. 6d.

Howitt (A. M.)—*An Art-Student in Munich.* By ANNA MARY HOWITT. 2 vols. post 8vo. price 14s.

Howitt.—*The Children's Year.* By Mary HOWITT. With Four Illustrations, from Designs by A. M. HOWITT. Square 16mo. 5s.

Howitt.—*Tallangetta, the Squatter's Home: A Story of Australian Life.* By WILLIAM HOWITT, Author of *Two Years in Victoria*, &c. 2 vols. post 8vo. price 18s.

Howitt.—*Land, Labour, and Gold; or, Two Years in Victoria: With Visit to Sydney and Van Diemen's Land.* By WILLIAM HOWITT. 2 vols. post 8vo. 21s.

Howitt.—*Visits to Remarkable Places: Old Halls, Battle-Fields, and Scenes illustrative of Striking Passages in English History and Poetry.* By WILLIAM HOWITT. With about 80 Wood Engravings. *New Edition.* 2 vols. square crown 8vo. price 25s.

William Howitt's Boy's Country Book: Being the Real Life of a Country Boy, written by himself; exhibiting all the Amusements, Pleasures, and Pursuits of Children in the Country. *New Edition;* with 40 Woodcuts. Fcp. 8vo. price 6s.

Howitt.—*The Rural Life of England.* By WILLIAM HOWITT. *New Edition,* corrected and revised; with Woodcuts by Bewick and Williams. Medium 8vo. 21s.

Huc.—*Christianity in China, Tartary, and Thibet.* By M. l'Abbé HUC, formerly Missionary Apostolic in China; Author of *The Chinese Empire*, &c. 2 vols. 8vo. 21s.

Huc.—*The Chinese Empire: A Sequel to Huc and Gabet's Journey through Tartary and Thibet.* By the Abbé HUC, formerly Missionary Apostolic in China. *Second Edition;* with Map. 2 vols. 8vo. 24s.

Hudson's Plain Directions for Making Wills in conformity with the Law: With a clear Exposition of the Law relating to the distribution of Personal Estate in the case of Intestacy, two Forms of Wills, and much useful information. *New and enlarged Edition;* including the Provisions of the Wills Act Amendment Act. Fcp. 8vo. 2s. 6d.

Hudson's Executor's Guide. *New and improved Edition;* with the Statutes enacted, and the Judicial Decisions pronounced since the last Edition incorporated, comprising the Probate and Administration Acts for England and Ireland, passed in the first Session of the New Parliament. Fcp. 8vo. [Just ready.]

Hudson and Kennedy.—*Where there's a Will there's a Way: An Ascent of Mont Blanc by a New Route and Without Guides.* By the Rev. C. HUDSON, M.A., St. John's College, Cambridge; and E. S. KENNEDY, B.A., Caius College, Cambridge. *Second Edition,* with Two Ascents of Monte Rosa; a Plate, and a coloured Map. Post 8vo. 5s. 6d.

Humboldt's Cosmos. Translated, with the Author's authority, by MRS. SABINE. Vols. I. and II. 16mo. Half-a-Crown each, sewed; 3s. 6d. each, cloth; or in post 8vo. 12s. each, cloth. Vol. III. post 8vo. 12s. 6d. cloth; or in 16mo. Part I. 2s. 6d. sewed, 3s. 6d. cloth; and Part II. 3s. sewed, 4s. cloth.

Humboldt's Aspects of Nature. Translated, with the Author's authority, by MRS. SABINE. 16mo. price 6s.: or in 2 vols. 3s. 6d. each, cloth; 2s. 6d. each, sewed.

Humphreys.—*Parables of Our Lord,* illuminated and ornamented in the style of the Missals of the Renaissance by HENRY NOEL HUMPHREYS. Square fcp. 8vo. 21s. in massive carved covers; or 30s. bound in morocco by Hayday.

Hunt.—*Researches on Light in its Chemical Relations;* embracing a Consideration of all the Photographic Processes. By ROBERT HUNT, F.R.S. *Second Edition,* with Plate and Woodcuts. 8vo. 10s. 6d.

Hutton.—*A Hundred Years Ago: An Historical Sketch, 1755 to 1756.* By JAMES HUTTON. Post 8vo.

Idle.—*Hints on Shooting, Fishing, &c., both on Sea and Land, and in the Fresh-Water Lochs of Scotland: Being the Experiences of C. IDLE, Esq.* Fcp. 8vo. 5s.

Mrs. Jameson's Legends of the Saints and Martyrs, as represented in Christian Art: Forming the First Series of *Sacred and Legendary Art*. Third Edition, revised and improved; with 17 Etchings and upwards of 180 Woodcuts, many of which are new in this Edition. 2 vols. square crown 8vo. 31s. 6d.

Mrs. Jameson's Legends of the Monastic Orders, as represented in Christian Art. Forming the Second Series of *Sacred and Legendary Art*. Second Edition, enlarged; with 11 Etchings by the Author, and 88 Woodcuts. Square crown 8vo. price 28s.

Mrs. Jameson's Legends of the Madonna, as represented in Christian Art: Forming the Third Series of *Sacred and Legendary Art*. Second Edition, revised and improved: with numerous Etchings from Drawings by the Author, and upwards of 150 Woodcuts. Square crown 8vo. [Nearly ready.]

Mrs. Jameson's Commonplace-Book of Thoughts, Memories, and Fancies, Original and Selected. Part I. Ethics and Character; Part II. Literature and Art. *Second Edit.* revised and corrected; with Etchings and Woodcuts. Crown 8vo. 18s.

Mrs. Jameson's Two Lectures on the Employment of Women.

1. *STORIES of CHARITY*, Catholic and Protestant, Abroad and at Home. *Second Edition*, with new Preface. Fcp. 8vo. 4s.
2. *THE COMMUNION of LABOUR: A Second Lecture on the Social Employments of Women.* Fcp. 8vo. 3s.

Jaquemet's Compendium of Chronology: Containing the most important Dates of General History, Political, Ecclesiastical, and Literary, from the Creation of the World to the end of the Year 1854. Edited by the Rev. J. ALCOCK, M.A. *Second Edition.* Post 8vo. price 7s. 6d.

Lord Jeffrey's Contributions to The Edinburgh Review. A New Edition, complete in One Volume, with a Portrait engraved by Henry Robinson, and a Vignette. Square crown 8vo. 21s. cloth; or 30s. calf. —Or in 3 vols. 8vo. price 42s.

Bishop Jeremy Taylor's Entire Works: With Life by BISHOP HEBER. Revised and corrected by the Rev. CHARLES PAGE EDEN, Fellow of Oriel College, Oxford. Now complete in 10 vols. 8vo. 10s. 6d. each.

Johns.—The Land of Silence and the Land of Darkness. Being Two Essays on the Blind and on the Deaf and Dumb. By the Rev. B. G. JOHNS, Chaplain of the Blind School, St. George's Fields, Southwark. Fcp. 8vo. price 4s. 6d.

Johnston.—A Dictionary of Geography, Descriptive, Physical, Statistical, and Historical: Forming a complete General Gazetteer of the World. By A. KEITH JOHNSTON, F.R.S.E., F.R.G.S., F.G.S., Geographer at Edinburgh in Ordinary to Her Majesty. Second Edition, thoroughly revised. In 1 vol. of 1,360 pages, comprising about 50,000 Names of Places. 8vo. 36s. cloth; or half-bound in Russia, 41s.

Kemble.—The Saxons in England: A History of the English Commonwealth till the Norman Conquest. By JOHN M. KEMBLE, M.A., &c. 2 vols. 8vo. 28s.

Kesteven.—A Manual of the Domestic Practice of Medicine. By W. B. KESTIVEN, Fellow of the Royal College of Surgeons of England, &c. Square post 8vo. 7s. 6d.

Kirby and Spence's Introduction to Entomology; or, Elements of the Natural History of Insects: Comprising an Account of Noxious and Useful Insects, of their Metamorphoses, Food, Stratagems, Habitations, Societies, Motions, Noises, Hybernation, Instinct, &c. *Seventh Edition*, with an Appendix relative to the Origin and Progress of the work. Crown 8vo. 5s.

Mrs. R. Lee's Elements of Natural History; or, First Principles of Zoology: Comprising the Principles of Classification, interspersed with amusing and instructive Accounts of the most remarkable Animals. New Edition; Woodcuts. Fcp. 8vo. 7s. 6d.

Letters to my Unknown Friends. By a LADY, Author of *Letters on Happiness*. Fourth Edition. Fcp. 8vo. 5s.

Letters on Happiness, addressed to a Friend. By a LADY, Author of *Letters to my Unknown Friends*. Fcp. 8vo. 6s.

L.E.L.—The Poetical Works of Letitia Elizabeth Landon; comprising the *Improvvisatrice*, the *Venetian Bracelet*, the *Golden Violet*, the *Troubadour*, and Poetical Remains. New Edition; with 2 Vignettes by R. Doyle. 2 vols. 16mo. 10s. cloth; morocco, 21s.

Dr. John Lindley's Theory and Practice of Horticulture; or, an Attempt to explain the principal Operations of Gardening upon Physiologcal Grounds: Being the Second Edition of the *Theory of Horticulture*, much enlarged; with 98 Woodcuts. 8vo. 21s.

Dr. John Lindley's Introduction to Botany. New Edition, with Corrections and copious Additions. 2 vols. 8vo. with Six Plates and numerous Woodcuts, price 24s.

LARDNER'S CABINET CYCLOPÆDIA

Of History, Biography, Literature, the Arts and Sciences, Natural History, and Manufactures.
A Series of Original Works by

SIR JOHN HERSCHEL,
SIR JAMES MACKINTOSH,
ROBERT SOUTHY,
SIR DAVID BREWSTER,

THOMAS KEIGHTLEY,
JOHN FORSTER,
SIR WALTER SCOTT,
THOMAS MOORE,

BISHOP THIRLWALL,
THE REV. G. R. GLEIG,
J. C. L. DE SISMONDI,
JOHN PHILLIPS, F.R.S., G.S.

AND OTHER EMINENT WRITERS.

Complete in 132 vols. fcp. 8vo. with Vignette Titles, price, in cloth, Nineteen Guineas.
The Works *separately*, in Sets or Series, price Three Shillings and Sixpence each Volume.

A List of the WORKS composing the CABINET CYCLOPÆDIA:—

- | | | | |
|--|------------------|---|------------------|
| 1. Bell's History of Russia | 3 vols. 10s. 6d. | 31. Lardner on Heat | 1 vol. 3s. 6d. |
| 2. Bell's Lives of British Poets | 2 vols. 7s. | 33. Lardner's Hydrostatics and Pneumatics | 1 vol. 3s. 6d. |
| 3. Brewster's Optics | 1 vol. 3s. 6d. | 36. Lardner and Walker's Electricity and | |
| 4. Cooley's Maritima and Inland Discovery | 3 vols. 10s. 6d. | Magnetism | 2 vols. 7s. |
| 5. Crowe's History of France | 3 vols. 10s. 6d. | 37. Mackintosh, Forster, and Courtenay's | |
| 6. De Morgan on Probabilities | 1 vol. 3s. 6d. | Lives of British Statesmen | 7 vols. 24s. 6d. |
| 7. De Sismondi's History of the Italian | | 38. Mackintosh, Wallace, and Bell's History | |
| Republics | 1 vol. 3s. 6d. | of England | 10 vols. 35s. |
| 8. De Sismondi's Fall of the Roman Empire | 2 vols. 7s. | 39. Montgomery and Shelley's eminent Ita- | |
| 9. Donovan's Chemistry | 1 vol. 3s. 6d. | lian, Spanish, and Portuguese Authors | 3 vols. 10s. 6d. |
| 10. Donovan's Domestic Economy | 2 vols. 7s. | 40. Moore's History of Ireland | 4 vols. 14s. |
| 11. Dunham's Spain and Portugal | 5 vols. 17s. 6d. | 41. Nicolas's Chronology of History | 1 vol. 3s. 6d. |
| 12. Dunham's History of Denmark, Sweden, | | 42. Phillips's Treatise on Geology | 2 vols. 7s. |
| and Norway | 3 vols. 10s. 6d. | 43. Powell's History of Natural Philosophy | 1 vol. 3s. 6d. |
| 13. Dunham's History of Poland | 1 vol. 3s. 6d. | 44. Porter's Treatise on the Manufacture of | |
| 14. Dunham's Germanic Empire | 3 vols. 10s. 6d. | Silk | 1 vol. 3s. 6d. |
| 15. Dunham's Europe during the Middle | | 45. Porter's Manufactures of Porcelain and | |
| Ages | 3 vols. 14s. | Glass | 1 vol. 3s. 6d. |
| 16. Dunham's British Dramatists | 2 vols. 7s. | 46. Roscoe's British Lawyers | 1 vol. 3s. 6d. |
| 17. Dunham's Lives of Early Writers of | | 47. Scott's History of Scotland | 2 vols. 7s. |
| Great Britain | 1 vol. 3s. 6d. | 48. Shelley's Lives of eminent French | |
| 18. Ferguson's History of the United States | 2 vols. 7s. | Authors | 2 vols. 7s. |
| 19. Foshrooke's Grecian & Roman Antiquities | 2 vols. 7s. | 49. Shackard and Swainson's Insects | 1 vol. 3s. 6d. |
| 20. Forster's Lives of the Statesmen of the | | 50. Southey's Lives of British Admirals | 5 vols. 17s. 6d. |
| Commonwealth | 5 vols. 17s. 6d. | 51. Stebbing's Church History | 2 vols. 7s. |
| 21. Gleig's Lives of British Military Com- | | 52. Stebbing's History of the Reformation .. | 2 vols. 7s. |
| manders | 3 vols. 10s. 6d. | 53. Swainson's Discourse on Natural History | 1 vol. 3s. 6d. |
| 22. Grattan's History of the Netherlands | 1 vol. 3s. 6d. | 54. Swainson's Natural History and Classi- | |
| 23. Henslow's Botany | 1 vol. 3s. 6d. | fication of Animals | 1 vol. 3s. 6d. |
| 24. Herschel's Astronomy | 1 vol. 3s. 6d. | 55. Swainson's Habits and Instincts of | |
| 25. Herschel's Discourse on Natural Philo- | | Animals | 1 vol. 3s. 6d. |
| sophy | 1 vol. 3s. 6d. | 56. Swainson's Birds | 2 vols. 7s. |
| 26. History of Rome | 2 vols. 7s. | 57. Swainson's Fish, Reptiles, &c. | 2 vols. 7s. |
| 27. History of Switzerland | 1 vol. 3s. 6d. | 58. Swainson's Quadrupeds | 1 vol. 3s. 6d. |
| 28. Holland's Manufactures in Metal | 3 vols. 10s. 6d. | 59. Swainson's Shells and Shell-Fish | 1 vol. 3s. 6d. |
| 29. James's Lives of Foreign Statesmen | 5 vols. 17s. 6d. | 60. Swainson's Animals in Menageries | 1 vol. 3s. 6d. |
| 30. Kater and Lardner's Mechanics | 1 vol. 3s. 6d. | 61. Swainson's Taxidermy and Biography of | |
| 31. Keightley's Outlines of History | 1 vol. 3s. 6d. | Zoologists | 1 vol. 3s. 6d. |
| 32. Lardner's Arithmetic | 1 vol. 3s. 6d. | 62. Thirlwall's History of Greece | 8 vols. 28s. |
| 33. Lardner's Geometry | 1 vol. 3s. 6d. | | |

Linwood.—*Anthologia Oxoniensis*, sive Florilegium c Lusibus poeticis diversorum Oxoniensium Græcis et Latinis decerptum. Curante GUILIELMO LINWOOD, M.A., Ædis Christi Alumno. 8vo. price 14s.

Lorimer's (C.) Letters to a Young Master Mariner on some Subjects connected with his Calling. New Edition. Fcp. 8vo. 5s. 6d.

Loudon's Encyclopædia of Gardening: Comprising the Theory and Practice of Horticulture, Floriculture, Arboriculture, and Landscape-Gardening. With many hundred Woodcuts. New Edition, corrected and improved by MRS. LOUDON. 8vo. 50s.

Loudon's Encyclopædia of Trees and Shrubs, or *Arboretum et Fruticetum Britannicum* abridged: Containing the Hardy Trees and Shrubs of Great Britain, Native and Foreign, Scientifically and Popularly Described. With about 2,000 Woodcuts. 8vo. 50s.

Loudon's Encyclopædia of Agriculture: Comprising the Theory and Practice of the Valuation, Transfer, Laying-out, Improvement, and Management of Landed Property, and of the Cultivation and Economy of the Animal and Vegetable Productions of Agriculture. New and cheaper Edition; with 1,100 Woodcuts. 8vo. 31s. 6d.

Loudon's Encyclopædia of Plants: Comprising the Specific Character, Description, Culture, History, Application in the Arts, and every other desirable Particular respecting all the Plants found in Great Britain. New Edition, corrected by MRS. LOUDON. With upwards of 12,000 Woodcuts. 8vo. £3. 13s. 6d.—Second Supplement, 21s.

Loudon's Encyclopædia of Cottage, Farm, and Villa Architecture and Furniture. New Edition, edited by MRS. LOUDON; with more than 2,000 Woodcuts. 8vo. 63s.

Loudon's Self-Instruction for Young Gardeners, Foresters, Bailiffs, Land Stewards, and Farmers; in Arithmetic, Book-keeping, Geometry, Mensuration, Practical Trigonometry, Mechanics, Land-Surveying, Levelling, Planning and Mapping, Architectural Drawing, and Isometrical Projection and Perspective. 8vo. Portrait, 7s. 6d.

Loudon's Hortus Britannicus; or, Catalogue of all the Plants found in Great Britain. New Edition, corrected by MRS. LOUDON. 8vo. 31s. 6d.

Mrs. Loudon's Lady's Country Companion; or, How to Enjoy a Country Life Rationally. Fourth Edition, with Plates and Woodcuts. Fcp. 8vo. 5s.

Mrs. Loudon's Amateur Gardener's Calendar, or Monthly Guide to what should be avoided and done in a Garden. 16mo. with Woodcuts, 7s. 6d.

Low's Elements of Practical Agriculture; comprehending the Cultivation of Plants, the Husbandry of the Domestic Animals, and the Economy of the Farm. New Edition; with 200 Woodcuts. 8vo. 21s.

Macaulay.—Speeches of the Right Hon. Lord Macaulay. Corrected by HIMSELF. 8vo. price 12s.

Macaulay. — The History of England from the Accession of James II. By the Right Hon. LORD MACAULAY. New Edition. Vols. I. and II. 8vo. price 32s.; Vols. III. and IV. price 36s.

Lord Macaulay's Critical and Historical Essays contributed to The Edinburgh Review. Four Editions, as follows:—

1. A LIBRARY EDITION (the Eighth), in 3 vols. 8vo. price 36s.
2. Complete in ONE VOLUME, with Portrait and Vignette. Square crown 8vo. price 21s. cloth; or 30s. calf.
3. Another NEW EDITION, in 3 vols. fcp. 8vo. price 21s. cloth.
4. THE PEOPLE'S EDITION, in 2 vols. crown 8vo. price 8s. cloth.

Macaulay.—Lays of Ancient Rome, with Iery and the Armada. By the Right Hon. LORD MACAULAY. New Edition. 16mo. price 4s. 6d. cloth; or 10s. 6d. bound in morocco.

Lord Macaulay's Lays of Ancient Rome. With numerous Illustrations, Original and from the Antique, drawn on Wood by George Scharf, jun., and engraved by Samuel Williams. New Edition. Fcp. 4to. price 21s. boards; or 42s. bound in morocco.

Mac Donald. — Poems. By George MAC DONALD, Author of *Within and Without*. Fcp. 8vo. 7s.

Mac Donald.—Within and Without: A Dramatic Poem. By GEORGE MAC DONALD. Second Edition, revised; fcp. 8vo. 4s. 6d.

Macdonald.—Villa Verocchio; or, the Youth of Leonardo da Vinci: A Tale. By the late Miss D. L. MACDONALD. Fcp. 8vo. price 6s.

MacDougall.—The Theory of War illustrated by numerous Examples from History. By Lieutenant-Colonel MACDOUGALL, Superintendent of Studies in the Royal Military College, Sandhurst. Post 8vo. with 10 Plans of Battles, price 10s. 6d.

M'Dougall.—The Eventful Voyage of H.M. Discovery Ship Resolute to the Arctic Regions in Search of Sir John Franklin and the Missing Crews of H.M. Discovery Ships Erebus and Terror, 1852, 1853, 1854. To which is added an Account of her being fallen in with by an American Whaler, after her abandonment in Barrow Straite, and of her presentation to Queen Victoria by the Government of the United States. By GEORGE F. M'DOUGALL, Master. With a coloured Chart; 8 Illustrations in tinted Lithography; and 22 Woodcuts. 8vo. price 21s. cloth.

Sir James Mackintosh's Miscellaneous Works: Including his Contributions to The Edinburgh Review. Complete in One Volume; with Portrait and Vignette. Square crown 8vo. 21s. cloth; or 30s. bound in calf: or in 3 vols. fcp. 8vo. 21s.

Sir James Mackintosh's History of England from the Earliest Times to the final Establishment of the Reformation. Library Edition, revised. 2 vols. 8vo. 21s.

Macleod.—The Theory and Practice of Banking: With the Elementary Principles of Currency, Prices, Credit, and Exchanges. By HENRY DUNNING MACLEOD, of the Inner Temple, Esq., Barrister-at-Law. 2 vols. royal 8vo. price 30s.

Macnaught.—The Doctrine of Inspiration: Being an Inquiry concerning the Infallibility, Inspiration, and Authority of Holy Writ. By the Rev. JOHN MACNAUGHT, M.A. *Second Edition*, revised. Crown 8vo. price 4s. 6d.

McCulloch's Dictionary, Practical, Theoretical, and Historical, of Commerce and Commercial Navigation. Illustrated with Maps and Plans. New Edition, corrected to the Present Time; with a Supplement. 8vo. price 50s. cloth; half-russia, 55s.

McCulloch's Dictionary, Geographical, Statistical, and Historical, of the various Countries, Places, and principal Natural Objects in the World. Illustrated with Six large Maps. New Edition, revised; with a Supplement. 2 vols. 8vo. price 63s.

Maguire.—Rome; its Ruler and its Institutions. By JOHN FRANCIS MAGUIRE, M.P. With a Portrait of Pope Pius IX. Post 8vo. price 10s. 6d.

Maitland.—The Church in the Catacombs: A Description of the Primitive Church of Rome. Illustrated by its Sepulchral Remains. By the Rev. CHARLES MAITLAND. New Edition; with several Woodcuts. 8vo. price 14s.

Out-of-Doors Drawing.—Aphorisms on Drawing. By the Rev. S. C. MALAN, M.A. of Balliol College, Oxford; Vicar of Broadwindsor, Dorset. Post 8vo. 3s. 6d.

Mrs. Marcet's Conversations on Chemistry, in which the Elements of that Science are familiarly explained and illustrated by Experiments. New Edition, enlarged and improved. 2 vols. fcp. 8vo. price 14s.

Mrs. Marcet's Conversations on Natural Philosophy, in which the Elements of that Science are familiarly explained. New Edition, enlarged and corrected; with 23 Plates. Fcp. 8vo. price 10s. 6d.

Martineau.—Endeavours after the Christian Life: Discourses. By JAMES MARTINEAU. 2 vols. post 8vo. 7s. 6d. each.

Martineau.—Hymns for the Christian Church and Home. Collected and edited by JAMES MARTINEAU. *Eleventh Edition*, 32mo. 3s. 6d. cloth, or 5s. calf; *Fifth Edition*, 32mo. 1s. 4d. cloth, or 1s. 8d. roan.

Martineau.—Miscellanies: Comprising Essays on Dr. Priestley, Arnold's Life and Correspondence, Church and State, Theodore Parker's Discourse of Religion, "Phases of Faith," the Church of England, and the Battle of the Churchæa. By JAMES MARTINEAU. Post 8vo. 9s.

Maunder's Scientific and Literary Treasury: A new and popular Encyclopædia of Science and the Belles-Lettres; including all branches of Science, and every subject connected with Literature and Art. New Edition. Fcp. 8vo. price 10s. cloth; bound in roan, 12s.; calf, 12s. 6d.

Maunder's Biographical Treasury; consisting of Memoirs, Sketches, and brief Notices of above 12,000 Eminent Persons of All Ages and Nations, from the Earliest Period of History: Forming a new and complete Dictionary of Universal Biography. Ninth Edition, revised throughout. Fcp. 8vo. 10s. cloth; bound in roan, 12s.; calf, 12s. 6d.

Maunder's Treasury of Knowledge, and Library of Reference. Comprising an English Dictionary and Grammar, a Universal Gazetteer, a Classical Dictionary, a Chronology, a Law Dictionary, a Synopsis of the Peerage, numerous useful Tables, &c. New Edition, carefully revised and corrected throughout: With Additions. Fcp. 8vo. 10s. cloth; bound in roan, 12s.; calf, 12s. 6d.

Maunder's Treasury of Natural History; or, a Popular Dictionary of Animated Nature: In which the Zoological Characteristics that distinguish the different Classes, Genera, and Species, are combined with a variety of interesting Information illustrative of the Habits, Instincts, and General Economy of the Animal Kingdom. With 900 Woodcuts. New Edition. Fcp. 8vo. price 10s. cloth; roan, 12s.; calf, 12s. 6d.

Maunder's Historical Treasury; comprising a General Introductory Outline of Universal History, Ancient and Modern, and a Series of separate Histories of every principal Nation that exists; their Rise, Progress, and Present Condition, the Moral and Social Character of their respective Inhabitants, their Religion, Manners and Customs, &c. New Edition; revised throughout, with a new GENERAL INDEX. Fcp. 8vo. 10s. cloth; roan, 12s.; calf, 12s. 6d.

Maunder's Geographical Treasury.—The Treasury of Geography, Physical, Historical, Descriptive, and Political; containing a succinct Account of Every Country in the World: Preceded by an Introductory Outline of the History of Geography; a Familiar Inquiry into the Varieties of Race and Language exhibited by different Nations; and a View of the Relations of Geography to Astronomy and the Physical Sciences. Commenced by the late SAMUEL MAUNDER; completed by WILLIAM HUGHES, F.R.G.S., late Professor of Geography in the College for Civil Engineers. *New Edition*; with 7 Maps and 16 Steel Plates. Fcp. 8vo. 10s. cloth; roan, 12s.; calf, 12s. 6d.

Melville.—The Confidence-Man: His Masquerade. By HERMAN MELVILLE, Author of *Typee*, *Omoo*, &c. Fcp. 8vo. 5s.

Merivale.—A History of the Romans under the Empire. By the Rev. CHARLES MERIVALE, B.D., late Fellow of St. John's College, Cambridge. 8vo. with Maps.

Vols. I. and II. comprising the History to the Fall of *Julius Cæsar*. Second Edition. 28s.
Vol. III. to the establishment of the Monarchy by *Augustus*. Second Edition. 14s.
Vols. IV. and V. from *Augustus* to *Claudius*, A.C. 27 to A.D. 54. 32s.

Merivale.—The Fall of the Roman Republic: A Short History of the Last Century of the Commonwealth. By the Rev. C. MERIVALE, B.D., late Fellow of St. John's College, Cambridge. New Edition. 12mo. 7s. 6d.

Merivale.—An Account of the Life and Letters of Cicero. Translated from the German of ABKEN; and Edited by the Rev. CHARLES MERIVALE, B.D. 12mo. 9s. 6d.

Merivale (L. A.).—Christian Records: A Short History of Apostolic Age. By L. A. MERIVALE. Fcp. 8vo. 7s. 6d.

Miles.—The Horse's Foot, and How to Keep it Sound. *Eighth Edition*; with an Appendix on Shoeing in general, and Hunters in particular, 12 Plates and 12 Woodcuts. By W. MILES, Esq. Imperial 8vo. 12s. 6d.

* * Two Casts or Models of Off Fore Feet, No. 1, *Shod for All Purposes*, No. 2, *Shod with Leather*, on Mr. Miles's plan, may be had, price 3s. each.

Miles.—A Plain Treatise on Horse-Shoeing. By WILLIAM MILES, Esq. With Plates and Woodcuts. Small 4to. price 5s.

Milner's History of the Church of Christ. With Additions by the late Rev. ISAAC MILNER, D.D., F.R.S. A New Edition, revised, with additional Notes by the Rev. T. GRANTHAM, B.D. 4 vols. 8vo. price 52s.

Montgomery.—Memoirs of the Life and Writings of James Montgomery: Including Selections from his Correspondence, Remains in Prose and Verse, and Conversations. By JOHN HOLLAND and JAMES EVERETT. With Portraits and Vignettes. 7 vols. post 8vo. price £3. 13s. 6d.

James Montgomery's Poetical Works: Collective Edition; with the Author's Autobiographical Prefaces, complete in One Volume; with Portrait and Vignette. Square crown 8vo. price 10s. 6d. cloth; morocco, 21s.—Or, in 4 vols. fcp. 8vo. with Portrait, d 7 other Plates, price 14s.

Moore.—The Power of the Soul over the Body, considered in relation to Health and Morals. By GEORGE MOORE, M.D. *Fifth Edition*. Fcp. 8vo. 6s.

"It shows that unless the inward principle be disciplined, purified, and enlightened, vainly must we look for that harmony between mind and body so necessary to human enjoyment. . . . We would say, Read the book." ATHENÆUM.

Moore.—Man and his Motives. By George MOORE, M.D. *Third Edition*. Fcp. 8vo. 6s.

Moore.—The Use of the Body in relation to the Mind. By GEORGE MOORE, M.D. *Third Edition*. Fcp. 8vo. 6s.

Moore.—Memoirs, Journal, and Correspondence of Thomas Moore. Edited by the Right Hon. LORD JOHN RUSSELL, M.P. With Portraits and Vignette Illustrations. 8 vols. post 8vo. price 10s. 6d. each.

Thomas Moore's Poetical Works: Comprising the Author's recent Introductions and Notes. The Traveller's Edition, complete in One Volume, printed in Ruby Type; with a Portrait. Crown 8vo. 12s. 6d. cloth; morocco by Hayday, 21s.—Also the Library Edition complete in 1 vol. medium 8vo. with Portrait and Vignettes, 21s. cloth; morocco by Hayday, 42s.—And the First collected Edition, in 10 vols. fcp. 8vo. with Portrait and 19 Plates, price 35s.

Moore.—Poetry and Pictures from Thomas Moore: Being Selections of the most popular and admired of Moore's Poems, copiously illustrated with highly-finished Wood Engravings from original Designs by

| | |
|-----------------------|-------------------------|
| C. W. COPE, R.A. | F. R. PICKERSGILL, R.A. |
| E. C. CORBOULD, | S. READ, |
| J. CROFTSEY, | G. THOMAS, |
| E. DUNCAN, | F. TOPHAM, |
| BIRKET FOSTER, | H. WARREN, |
| J. C. HORSLEY, A.R.A. | HARRISON WEIR, and |
| E. LE JEUNE, | P. WYCHUD. |

Fcp. 4to., printed on toned paper, and elegantly bound. [Nearly ready.]

Moore's Epicurean. New Edition, with the Notes from the collective edition of Moore's Poetical Works; and a Vignette engraved on Wood from an original Design by D. MACLISE, R.A. 16mo. 5s. cloth; or 12s. 6d. morocco by Hayday.

Moore's Songs, Ballads, and Sacred Songs. New Edition, printed in Ruby Type; with the Notes from the collective edition of Moore's Poetical Works, and a Vignette from a Design by T. CRESWICK, R.A. 32mo. 2s. 6d.—An Edition in 16mo. with Vignette by R. Doyle, price 5s.; or 12s. 6d. morocco by Hayday.

Moore's Lalla Rookh: An Oriental Romance. With 13 highly-finished Steel Plates from Original Designs by Corbould, Meadows, and Stephanoff, engraved under the superintendence of the late Charles Heath. New Edition. Square crown 8vo. price 15s. cloth; morocco, 28s.

Moore's Lalla Rookh. New Edition, printed in Ruby Type; with the Preface and Notes from the collective edition of *Moore's Poetical Works*, and a Frontispiece from a Design by Kenny Meadows. 32mo. 2s. 6d. —An Edition in 16mo. with Vignette, 5s.; or 12s. 6d. morocco by Hayday.

Moore's Irish Melodies. A New Edition, with 13 highly-finished Steel Plates, from Original Designs by

| | |
|---------------------|-----------------------|
| C. W. COPE, R.A. | D. MACLISE, R.A. |
| T. CRESWICK, R.A. | J. E. MILLAIS, A.R.A. |
| A. L. EGG, A.R.A. | W. MULREATH, R.A. |
| W. P. FRITH, R.A. | J. SANT, R.A. |
| W. E. FROST, A.R.A. | F. STONE, A.R.A.; and |
| J. C. HORSLEY, | E. M. WARD, R.A. |

Square crown 8vo. price 21s. cloth; or 31s. 6d. handsomely bound in morocco.

Moore's Irish Melodies, printed in Ruby Type; with the Preface and Notes from the collective edition of *Moore's Poetical Works*, the Advertisements originally prefixed, and a Portrait of the Author. 32mo. 2s. 6d. —An Edition in 16mo. with Vignette, 5s.; or 12s. 6d. morocco by Hayday.

Moore's Irish Melodies. Illustrated by D. MacLise, R.A. New Edition; with 161 Designs, and the whole of the Letterpress engraved on Steel, by F. P. Becker. Super-royal 8vo. 31s. 6d. boards; £2. 12s. 6d. morocco by Hayday.

Moore's Irish Melodies, the Music with the Words; the Symphonies and Accompaniments by Sir John Stevenson, Mus. Doc. Complete in One Volume, small Music size, convenient and legible at the pianoforte, but more portable than the usual form of Musical publications. Imperial 8vo. 31s. 6d. cloth; or 42s. half-bound in morocco.

Moore.—The Crosses, Altar, and Ornaments in the Churches of St. Paul's, Knightsbridge, and St. Barnabas, Pimlico: A concise Report of the Proceedings and Judgments in the Cases of Westerton v. Liddell, Horne, and others, and Beal v. Liddell, Parke, and Evans; as heard and determined by the Consistory Court of London, the Arches Court of Canterbury, and the Judicial Committee of H.M. Most Hon. Privy Council. By EDMUND F. MOORE, Esq., M.A., Barrister-at-Law. Royal 8vo. price 12s. cloth.

Morell.—Elements of Psychology: Part I., containing the Analysis of the Intellectual Powers. By J. D. MORELL, M.A., One of Her Majesty's Inspectors of Schools. Post 8vo. 7s. 6d.

Morning Clouds. [A book of practical ethics, in form of letters of counsel, encouragement, and sympathy, specially addressed to young women on their entrance into life.] Post 8vo. price 7s.

Moseley.—The Mechanical Principles of Engineering and Architecture. By H. MOSELEY, M.A., F.R.S., Canon of Bristol, &c. Second Edition, enlarged; with numerous Corrections and Woodcuts. 8vo. 24s.

Memoirs and Letters of the late Colonel ARMINE S. H. MOUNTAIN, C.B., Aide-de-Camp to the Queen, and Adjutant-General of Her Majesty's Forces in India. Edited by Mrs. MOUNTAIN. With a Portrait drawn on Stone by R. J. LANE, A.E.R.A. Post 8vo. 8s. 6d.

Mure.—A Critical History of the Language and Literature of Ancient Greece. By WILLIAM MURE, M.P. of Caldwell. Second Edition. Vols. I. to III. 8vo. price 36s.; Vol. IV. price 15s.; Vol. V. price 18s.

Murray's Encyclopædia of Geography; comprising a complete Description of the Earth: Exhibiting its Relation to the Heavenly Bodies, its Physical Structure, the Natural History of each Country, and the Industry, Commerce, Political Institutions, and Civil and Social State of All Nations. Second Edition; with 82 Maps, and upwards of 1,000 other Woodcuts. 8vo. price 60s.

Neale.—The Closing Scene; or, Christianity and Infidelity contrasted in the Last Hours of Remarkable Persons. By the Rev. ERSKINE NEALE, M.A. New Editions. 2 vols. fcp. 8vo. price 6s. each.

Oldacre.—The Last of the Old Squires. A Sketch. By CEDRIC OLDACRE, Esq., of Sax-Normanbury, sometime of Christ Church, Oxon. Crown 8vo. price 9s. 6d.

Osborn.—Quedah; or, Stray Leaves from a Journal in Malayan Waters. By Captain SHEARD OSBORN, R.N., C.B., Author of *Stray Leaves from an Arctic Journal*, and of the *Narrative of the Discovery of the North-West Passage*. With a coloured Chart and tinted Illustrations. Post 8vo. price 10s. 6d.

Osborn.—The Discovery of the North-West Passage by H.M.S. *Investigator*, Captain R. M'CLURE, 1850–1854. Edited by Captain SHERARD OSBORN, C.B., from the Logs and Journals of Captain R. M'Clure. Second Edition, revised; with Additions to the Chapter on the Hybernation of Animals in the Arctic Regions, a Geological Paper by Sir RODERICK I. MURCHISON, a Portrait of Captain M'Clure, a coloured Chart and tinted Illustrations. 8vo. price 15s.

Owen.—Lectures on the Comparative Anatomy and Physiology of the Invertebrate Animals, delivered at the Royal College of Surgeons. By RICHARD OWEN, F.R.S., Hunterian Professor to the College. Second Edition, with 235 Woodcuts. 8vo. 21s.

Professor Owen's Lectures on the Comparative Anatomy and Physiology of the Vertebrate Animals, delivered at the Royal College of Surgeons in 1844 and 1846. With numerous Woodcuts. Vol. I. 8vo. price 14s.

Memoirs of Admiral Parry, the Arctic Navigator. By his Son, the Rev. E. PARRY, M.A. of Balliol College, Oxford; Domestic Chaplain to the Lord Bishop of London. Third Edition; with a Portrait and coloured Chart of the North-West Passage. Fcp. 8vo. price 5s.

Dr. Pereira's Elements of Materia Medica and Therapeutics. *Third Edition*, enlarged and improved from the Author's Materials, by A. S. TAYLOR, M.D., and G. O. REES, M.D.: With numerous Woodcuts. Vol. I. 8vo. 28s.; Vol. II. Part I. 21s.; Vol. II. Part II. 24s.

Dr. Pereira's Lectures on Polarised Light, together with a Lecture on the Microscope. 2d Edition, enlarged from Materials left by the Author, by the Rev. B. POWELL, M.A., &c. Fcp. 8vo. with Woodcuts, 7s.

Perry.—The Franks, from their First Appearance in History to the Death of King Pepin. By WALTER C. PERRY, Barrister-at-Law, Doctor in Philosophy and Master of Arts in the University of Göttingen. 8vo. price 12s. 6d.

Peschel's Elements of Physics. Translated from the German, with Notes, by E. WEST. With Diagrams and Woodcuts. 3 vols. fcp. 8vo. 21s.

Ida Pfeiffer's Lady's Second Journey round the World: From London to the Cape of Good Hope, Borneo, Java, Sumatra, Celebes, Ceram, the Moluccas &c., California, Panama, Peru, Ecuador, and the United States. 2 vols. post 8vo. 21s.

Phillips's Elementary Introduction to Mineralogy. A New Edition, with extensive Alterations and Additions, by H. J. BROOKE, F.R.S., F.G.S.; and W. H. MILLER, M.A., F.G.S. With numerous Wood Engravings. Post 8vo. 18s.

Phillips.—A Guide to Geology. By JOHN PHILLIPS, M.A., F.R.S., F.G.S., &c. Fourth Edition, corrected to the Present Time; with 4 Plates. Fcp. 8vo. 5s.

Phillips.—Figures and Descriptions of the Palæozoic Fossils of Cornwall, Devon, and West Somerset; observed in the course of the Ordnance Geological Survey of that District. By JOHN PHILLIPS, F.R.S., F.G.S., &c. 8vo. with 60 Plates, price 9s.

Piesse's Art of Perfumery, and Methods of Obtaining the Odours of Plants: With Instructions for the Manufacture of Perfumes for the Handkerchief, Scented Powders, Odorous Vinegars, Dentifrices, Pomatums, Cosmétiques, Perfumed Soap, &c.; and an Appendix on the Colours of Flowers, Artificial Fruit Essences, &c. *Second Edition*, revised and improved; with 46 Woodcuts. Crown 8vo. 8s. 6d.

Captain Portlock's Report on the Geology of the County of Londonderry, and of Parts of Tyrone and Fermanagh, examined and described under the Authority of the Master-General and Board of Ordnance. 8vo. with 48 Plates, price 24s.

Powell.—Essays on the Spirit of the Inductive Philosophy, the Unity of Worlds, and the Philosophy of Creation. By the Rev. BADEN POWELL, M.A., F.R.S., F.R.A.S., F.G.S., Savilian Professor of Geometry in the University of Oxford. Second Edition, revised. Crown 8vo. with Woodcuts, 12s. 6d.

Pycroft's Course of English Reading, adapted to every taste and capacity: With Literary Anecdotes. New and cheaper Edition. Fcp. 8vo. price 5s.

Raikes.—A Portion of the Journal kept by THOMAS RAIKES, Esq., from 1831 to 1847: Comprising Reminiscences of Social and Political Life in London and Paris during that period. Vols. I. and II. (*Second Edition*), post 8vo. 21s.; Vols. III. and IV. with *Index*, completing the work, price 21s.

Reade.—The Poetical Works of John Edmund Reade. New Edition, revised and corrected; with Additional Poems. 4 vols. fcp. 8vo. price 20s.

Dr. Reece's Medical Guide: Comprising a complete Modern Dispensatory, and a Practical Treatise on the distinguishing Symptoms, Causes, Prevention, Cure, and Palliation of the Diseases incident to the Human Frame. Seventeenth Edition, corrected and enlarged by the Author's Son, Dr. H. REECE, M.R.C.S., &c. 8vo. 12s.

Rich's Illustrated Companion to the Latin Dictionary and Greek Lexicon: Forming a Glossary of all the Words representing Visible Objects connected with the Arts, Manufactures, and Every-Day Life of the Ancients. With about 2,000 Woodcuts from the Antique. Post 8vo. 21s.

Richardson. — Fourteen Years' Experience of Cold Water: Its Uses and Abuses. By Captain M. RICHARDSON, late of the 4th Light Dragoons. Post 8vo. with Woodcuts, price 6s.

"The first object of Captain Richardson's book is to extend the use of the cold-water cure to the humbler classes, by a simpler mode of treatment. This simplicity principally consists in the substitution of wet bandages covered by dry bandages for the wet sheet and other processes of established hydropathy. Captain Richardson considers the bandage not only more beneficial medically than the sheet, but much more easily applied, while it does not interrupt a man's avocations, but can be worn even at work. The general expositions are followed by directions for the treatment of diseases under the Captain's system." SPECTATOR.

Horsemanship; or, the Art of Riding and Managing a Horse, adapted to the Guidance of Ladies and Gentlemen on the Road and in the Field: With Instructions for Breaking-in Colts and Young Horses. By CAPTAIN RICHARDSON, late of the 4th Light Dragoons. With 5 Plates. Square crown 8vo. 14s.

Household Prayers for Four Weeks; With additional Prayers for Special Occasions. To which is added a Course of Scripture Reading for Every Day in the Year. By the Rev. J. E. RIDDLE, M.A., Incumbent of St. Philip's, Leckhampton. Crown 8vo. price 3s. 6d.

Riddle's Complete Latin-English and English-Latin Dictionary, for the use of Colleges and Schools. *New and cheaper Edition, revised and corrected.* 8vo. 21s.

Separately {The English-Latin Dictionary, 7s.
{The Latin-English Dictionary, 15s.

Riddle's Diamond Latin-English Dictionary. A Guide to the Meaning, Quality, and right Accentuation of Latin Classical Words. Royal 32mo. price 4s.

Riddle's Copious and Critical Latin-English Lexicon, founded on the German-Latin Dictionaries of Dr. William Freund. *New and cheaper Edition.* Post 4to. 31s. 6d.

Rivers's Rose-Amateur's Guide; containing ample Descriptions of all the finest leading varieties of Roses, regularly classed in their respective Families; their History and Mode of Culture. Fifth Edition, corrected and improved. Fcp. 8vo. 3s. 6d.

Dr. E. Robinson's Greek and English Lexicon to the Greek Testament. A New Edition, revised and in great part re-written. 8vo. price 18s.

Mr. Henry Rogers's Essays selected from Contributions to the *Edinburgh Review*. Second and cheaper Edition, with Additions. 3 vols. fcp. 8vo. 21s.

Dr. Roget's Thesaurus of English Words and Phrases classified and arranged so as to facilitate the Expression of Ideas and assist in Literary Composition. Sixth Edition, revised and improved. Crown 8vo. 10s. 6d.

Ronalds's Fly-Fisher's Entomology: With coloured Representations of the Natural and Artificial Insect, and a few Observations and Instructions on Trout and Grayling Fishing. Fifth Edition, thoroughly revised by an Experienced Fly-Fisher; with 20 new coloured Plates. 8vo. 14s.

Rowton's Debater: A Series of complete Debates, Outlines of Debates, and Questions for Discussion; with ample References to the best Sources of Information. New Edition. Fcp. 8vo. 6s.

The Saints our Example. By the Author of *Letters to my Unknown Friends*, &c. Fcp. 8vo. price 7s.

Scherzer. — Travels in the Free States of Central America: Nicaragua, Honduras, and San Salvador. By Dr. CARL SCHERZER. With a coloured Map. 2 vols. post 8vo. 16s.

"Central America is not an inviting place for the lounging traveller. The roads are bad; there are no inns; food is scarce; the people are dishonest; scoundrels swarm; neither life nor property is safe. Dr. Scherzer travelled with guides of doubtful fidelity, was forced to keep his hand on his gun and revolver, sometimes compelled to eat a few black beans or starve; now groping at night through a forest, now escorted by barefooted soldiers with helmets of pasteboard and tin; now swinging in a hammock in a filthy hovel; anon receiving the President of a State by the light of a candle stuck in a bottle. Altogether having a hard and hazardous life of it. But he does not complain. A cheasier and braver traveller seldom has made his way in outlandish tracks far beyond the limits of the civilised world. . . . The Central American question will probably endow Dr. Scherzer's book with an additional attraction." GLEANER.

Dr. L. Schmitz's History of Greece, from the Earliest Times to the Taking of Corinth by the Romans, B.C. 146, mainly based upon Bishop Thirlwall's History. Fourth Edition, with Supplementary Chapters on the Literature and the Arts of Ancient Greece; and illustrated with a Map of Athens and 137 Woodcuts, designed from the Antique by G. Scharf, jun., F.S.A. 12mo. 7s. 6d.

Scrivenor's History of the Iron Trade, from the Earliest Records to the Present Period. New Edition, corrected. 8vo. price 10s. 6d.

Sir Edward Seaward's Narrative of his Shipwreck, and consequent Discovery of certain Islands in the Caribbean Sea. Third Edition. 2 vols. post 8vo. 21s.—An ABRIDGMENT, in 16mo. price 2s. 6d.

The Sermon in the Mount. Printed by C. Whittingham, uniformly with the *Thames Bible*; bound and clasped. 64mo. 1s. 6d.

Sewell.—Amy Herbert. By a Lady. Edited by the Rev. WILLIAM SEWELL, B.D., Fellow and Tutor of Exeter College, Oxford. New Edition. Fcp. 8vo. price 6s.

Sewell.—The Earl's Daughter. By the Author of *Amy Herbert*. Edited by the Rev. W. SEWELL, B.D. 2 vols. fcp. 8vo. 9s.

Sewell.—Gertrude: A Tale. By the Author of *Amy Herbert*. Edited by the Rev. W. SEWELL, B.D. Fcp. 8vo. price 6s.

Sewell.—Margaret Percival. By the Author of *Amy Herbert*. Edited by the Rev. W. SEWELL, B.D. 2 vols. fcp. 8vo. price 12s.

By the same Author, New Editions,

Ivors. 2 vols. fcp. 8vo. price 12s.

Cleve Hall. 2 vols. fcp. 8vo. price 12s.

Katharine Ashton. 2 vols. fcp. 8vo. 12s.

The Experience of Life. Fcp. 8vo. price 7s. 6d.

Laneton Parsonage: A Tale for Children, on the Practical Use of a portion of the Church Catechism. 2 vols. fcp. 8vo. price 12s.

Readings for Every Day in Lent: Compiled from the Writings of BISHOP JEREMY TAYLOR. Fcp. 8vo. price 5s.

Readings for a Month preparatory to Confirmation: Compiled from the Works of Writers of the Early and of the English Church. New and cheaper Edition. Fcp. 8vo. 4s.

Bowdler's Family Shakspeare: In which nothing is added to the Original Text; but those words and expressions are omitted which cannot with propriety be read aloud. Illustrated with Thirty-six Vignettes engraved on Wood from original Designs by

G. COOKE, R.A.
E. COOKE,
H. HOWARD, R.A.
H. SINGLETON,
E. SMIRKE, R.A.

T. SPOTTHARD, R.A.
H. THOMSON, R.A.
E. WESSELL, R.A.
E. WORDSWORTH, R.A.

New Edition, printed in a more convenient form. 6 vols. fcp. 8vo. price 30s. cloth; separately, 5s. each.

*. The LIBRARY EDITION, with the same Illustrations, in One Volume, medium 8vo, price 21s. cloth.

Sharp's New British Gazetteer, or Topographical Dictionary of the British Islands and Narrow Seas: Comprising concise Descriptions of about Sixty Thousand Places, Seats, Natural Features, and Objects of Note, founded on the best authorities. 2 vols. 8vo. price £2. 16s.

"We have already had occasion to mention this book, and a careful examination of its contents has convinced us of its great value. The remarkable clearness with which its condensations and abbreviations are made appears to us its most admirable feature. We have no book of similar bulk in the language containing anything like the amount of information of various kinds so well arranged and so easily accessible as in this new gazetteer. Every article bears the mark of studied, careful, and exact compilation. It comprehends both the topography and the hydrography of the United Kingdom, and is constructed on the plan of facilitating reference by bringing together as many articles as possible under distinct heads. . . . All the positions have been retaken from the maps; and not only the county but the quarter of the county given in which a name might be looked for. We must, in short, repeat with a liberal acquiescence what Mr. Sharp himself remarks of his five years' diligent labour, that it will be found to comprise, in a clear and legible type, more substantial information, collected from original sources, and put into a convenient form, than the bulkiest of its class." EXAMINER.

Short Whist; its Rise, Progress, and Laws: With Observations to make any one a Whist-Player. Containing also the Laws of Piquet, Cassino, Ecarté, Cribbage, Backgammon. By Major A. New Edition; to which are added, Precepts for Tyros, by Mrs. B. Fcp. 8vo. 3s.

Sinclair.—The Journey of Life. By CATHERINE SINCLAIR, Author of *The Business of Life*. New Edition. Fcp. 8vo. 5s.

Sir Roger De Coverley. From the Spectator. With Notes and Illustrations, by W. HENRY WILLS; and 12 Wood Engravings from Designs by F. TAYLER. Second and cheaper Edition. Crown 8vo. 10s. 6d.; or 21s. in morocco by Hayday.—An Edition without Woodcuts, in 16mo. price 1s.

The Sketches: Three Tales. By the Authors of *Amy Herbert*, *The Old Man's Home*, and *Hawkstone*. The Third Edition; with 6 Illustrations in Aquatint. Fcp. 8vo. price 4s. 6d. boards.

Smee's Elements of Electro-Metallurgy. Third Edition, revised, corrected, and considerably enlarged; with Electrotypes and numerous Woodcuts. Post 8vo. 10s. 6d.

Smith (G.)—History of Wesleyan Methodism: Vol. I. Wesley and his Times. By GEORGE SMITH, F.A.S., Member of the Royal Asiatic Society, &c.; Author of *Sacred Annals, or Researches into the History and Religion of Mankind*, &c. Crown 8vo. with 8 Facsimiles of Methodist Society Tickets, price 10s. 6d. cloth.

Smith (G. V.).—The Prophecies relating to Nineveh and the Assyrians. Translated from the Hebrew, with Historical Introductions and Notes, exhibiting the principal Results of the recent Discoveries. By **GEORGE VANCE SMITH, B.A.** Post 8vo. with a Map, price 10s. 6d. cloth.

Smith (J.).—The Voyage and Shipwreck of St. Paul : With Dissertations on the Life and Writings of St. Luke, and the Ships and Navigation of the Ancients. By **JAMES SMITH, of Jordanhill, Esq., F.R.S.** *Second Edition ;* with Charts, Views, and Woodcuts. Crown 8vo. 8s. 6d.

A Memoir of the Rev. Sydney Smith. By his Daughter, **LADY HOLLAND.** With a Selection from his Letters, edited by **MRS. AUSTIN.** *New Edition.* 2 vols. 8vo. 28s.

The Rev. Sydney Smith's Miscellaneous Works : Including his Contributions to The Edinburgh Review. Three Editions :—

1. A LIBRARY EDITION (the *Fourth*), in 3 vols. 8vo. with Portrait, 36s.
2. Complete in ONE VOLUME, with Portrait and Vignette. Square crown 8vo. price 21s. cloth ; or 30s. bound in calf.
3. Another NEW EDITION, in 3 vols. fcp. 8vo. price 21s.

The Rev. Sydney Smith's Elementary Sketches of Moral Philosophy, delivered at the Royal Institution in the Years 1804, 1805, and 1806. Third Edition. Fcp. 8vo. 7s.

Snow.—A Two-Years' Cruise off Tierra del Fuego, the Falkland Islands, Patagonia, and the River Plate. A Narrative of Life in the Southern Seas. By **W. PARKER SNOW,** late Commander of the Mission Yacht *Allen Gardiner* ; Author of "Voyage of the *Prince Albert* in Search of Sir John Franklin." With 3 coloured Charts and 6 tinted Illustrations. 2 vols. post 8vo. 24s.

"A Robinson-Crusoe style of narration, and a kind of rough and picturesque treatment, sustain the interest of the nautical descriptions more than might be supposed ; the wild and violent weather of the Falkland Islands, with the dangers of their navigation and the peculiar character of the River Plate, have a novelty beyond the common run of voyaging. The adventures in Tierra del Fuego are very interesting." **SPECTATOR.**

Robert Southey's Complete Poetical Works ; containing all the Author's last Introductions and Notes. *The Library Edition,* complete in One Volume, with Portrait and Vignette. Medium 8vo. price 21s. cloth ; 42s. bound in morocco. — Also, the *First collected Edition,* in 10 vols. fcp. 8vo. with Portrait and 19 Vignettes, price 35s.

Select Works of the British Poets ; from Chaucer to Lovelace inclusive. With Biographical Sketches by the late **ROBERT SOUTHEY.** Medium 8vo. price 30s.

Southey's Correspondence. — Selections from the Letters of Robert Southey, &c. Edited by his Son-in-Law, the Rev. **JOHN WOOD WARTER, B.D.,** Vicar of West Tarring, Sussex. 4 vols. post 8vo. price 42s.

The Life and Correspondence of the late Robert Southey. Edited by his Son, the Rev. **C. C. SOUTHEY, M.A.,** Vicar of Arleigh. With Portraits and Landscape Illustrations. 6 vols. post 8vo. price 63s.

Southey's Doctor, complete in One Volume. Edited by the Rev. **J. W. WARTER, B.D.** With Portrait, Vignette, Bust, and coloured Plate. Square crown 8vo. 21s.

Southey's Commonplace-Books, complete in Four Volumes. Edited by the Rev. **J. W. WARTER, B.D.** 4 vols. square crown 8vo. price £3. 18s.

Each *Commonplace-Book*, complete in itself, may be had separately, as follows :—

FIRST SERIES—CHOICE PASSAGES, &c. 18s.

SECOND SERIES—SPECIAL COLLECTIONS. 18s.

THIRD SERIES—ANALYTICAL READINGS. 21s.

FOURTH SERIES—ORIGINAL MEMORANDA, &c. 21s.

Southey's Life of Wesley ; and Rise and Progress of Methodism. *New Edition,* with Notes and Additions. Edited by the Rev. **C. C. SOUTHEY, M.A.** 2 vols. 8vo. with 2 Portraits, price 28s.

Spottiswoode.—A Tarantasse Journey through Eastern Russia, in the Autumn of 1856. By **WILLIAM SPOTTISWOODE, M.A., F.R.S.** With a Map of Russia, several Wood Engravings, and Seven Illustrations in tinted Lithography from Sketches by the Author. Post 8vo. price 10s. 6d.

Stephen.—Lectures on the History of France. By the Right Hon. **SIR JAMES STEPHEN, K.C.B., LL.D.,** Professor of Modern History in the University of Cambridge. Third Edition. 2 vols. 8vo. price 24s.

Stephen.—Essays in Ecclesiastical Biography ; from *The Edinburgh Review.* By the Right Hon. **SIR JAMES STEPHEN, K.C.B., LL.D.,** Professor of Modern History in the University of Cambridge. Third Edition. 2 vols. 8vo. 24s.

Stonehenge.—The Greyhound : Being a Treatise on the Art of Breeding, Rearing, and Training Greyhounds for Public Running ; their Diseases and Treatment : Containing also Rules for the Management of Coursing Meetings, and for the Decision of Courses. By **STONEHENGE.** With Frontispiece and Woodcuts. Square crown 8vo. price 21s. half-bound.

Stow.—The Training System, Moral Training School, and Normal Seminary for preparing Schoolmasters and Governesses. By DAVID STOW, Esq., Honorary Secretary to the Glasgow Normal Free Seminary. Tenth Edition; with Plates and Woodcuts. Post 8vo. price 6s.

Strickland.—Lives of the Queens of England. By AGNES STRICKLAND. Dedicated, by express permission, to Her Majesty. Embellished with Portraits of every Queen, engraved from the most authentic sources. Complete in 8 vols. post 8vo. price 7s. 6d. each.—Any Volume may be had separately to complete Sets.

Memoirs of Rear-Admiral Sir William Symonds, Knt., C.B., F.R.S., late Surveyor of the Navy. Published with the sanction of his Executors, as directed by his Will; and edited by J. A. SHARP. 8vo. with Plates and Wood Engravings.

[In the press.]

Taylor.—Loyola: and Jesuitism in its Rudiments. By ISAAC TAYLOR. Post 8vo. price 10s. 6d.

Taylor.—Wesley and Methodism. By ISAAC TAYLOR. Post 8vo. Portrait, 10s. 6d.

Thacker's Courser's Annual Remembrancer and Stud-Book: Being an Alphabetical Return of the Running at all the Public Coursing Clubs in England, Ireland, and Scotland, for the Season 1856-57; with the *Pedigrees* (as far as received) of the Dogs. By ROBERT ABRAM WELSH, Liverpool. 8vo. 21s.

. Published annually in October.

Thirlwall.—The History of Greece. By the Right Rev. the LORD BISHOP of ST. DAVID'S (the Rev. Connop Thirlwall). An improved Library Edition; with Maps. 8 vols. 8vo. price £3.—An Edition in 8 vols. fcp. 8vo. with Vignette Titles, price 28s.

Thomas.—Historical Notes relative to the History of England; embracing the Period from the Accession of King Henry VIII. to the Death of Queen Anne inclusive (1509 to 1714): Designed as a Book of instant Reference for the purpose of ascertaining the Dates of Events mentioned in History and in Manuscripts. The Names of Persons and Events mentioned in History within the above period placed in Alphabetical and Chronological Order, with Dates; and the Authority from whence taken given in each case, whether from Printed History or from Manuscripts. By F. S. THOMAS, Secretary of the Public Record Department. 3 vols. royal 8vo. price £2.

Thomson's Seasons. Edited by BOLTON CORNEY, Esq. Illustrated with 77 fine Wood Engravings from Designs by Members of the Etching Club. Square crown 8vo. 21s. cloth; or 36s. bound in morocco.

Thomson (the Rev. Dr.)—An Outline of the necessary Laws of Thought: A Treatise on Pure and Applied Logic. By WILLIAM THOMSON, D.D., Provost of Queen's College, Oxford. Fourth Edition, carefully revised. Fcp. 8vo. price 7s. 6d.

Thomson's Tables of Interest, at Three, Four, Four-and-a-Half, and Five per Cent., from One Pound to Ten Thousand, and from 1 to 365 Days, in a regular progression of single Days; with Interest at all the above Rates, from One to Twelve Months, and from One to Ten Years. Also, numerous other Tables of Exchanges, Time, and Discounts. New Edition. 12mo. price 8s.

Thornbury.—Shakspeare's England; or, Sketches of Social History during the Reign of Elizabeth. By G. W. THORNBURY, Author of *History of the Buccaneers, &c.* 2 vols. crown 8vo. 21s.

"A work which stands unrivalled for the variety and entertaining character of its contents, and which well deserves a place on the library-shelf, by the side either of the historians of England or the prince of dramatists."
JOHN BULL.

The Thumb Bible; or, Verbum Semper. By J. TAYLOR. Being an Epitome of the Old and New Testaments in English Verse. Reprinted from the Edition of 1693; bound and clasped. 64mo. 1s. 6d.

Bishop Tomline's Introduction to the Study of the Bible: Containing Proofs of the Authenticity and Inspiration of the Scriptures; a Summary of the History of the Jews; an Account of the Jewish Sects; and a brief Statement of Contents of several Books of the *Old Testament*. New Edition. Fcp. 8vo. 5s. 6d.

Tooke.—History of Prices, and of the State of the Circulation, during the Nine Years from 1848 to 1856 inclusive. Forming Vols. V. and VI. of Tooke's *History of Prices from 1792 to the Present Time*; and comprising a copious Index to the whole of the Six Volumes. By THOMAS TOOKE, F.R.S. and WILLIAM NEWMARCH. 2 vols. 8vo. price 52s. 6d.

Townsend.—Modern State Trials revised and illustrated with Essays and Notes. By W. C. TOWNSEND, Esq., M.A., Q.C. 2 vols. 8vo. price 30s.

COMPLETION OF THE TRAVELLER'S LIBRARY.

Summary of the Contents of the TRAVELLER'S LIBRARY, now complete in 102 Parts, price One Shilling each, or in 50 Volumes, price 2s. 6d. each in cloth.— To be had also, in complete Sets only, at Five Guineas per Set, bound in cloth, lettered, in 25 Volumes, classified as follows:—

VOYAGES AND TRAVELS.

IN EUROPE.

A CONTINENTAL TOUR BY J. BARROW.
ARCTIC VOYAGES AND } BY F. MAYNE.
DISCOVERIES }
BRITANY AND THE BIBLE BY I. HOPE.
BRITANY AND THE CHASE BY I. HOPE.
COESICA BY F. GREGOROVIVS.
GERMANY, &c. NOTES OF } BY S. LAING.
A TRAVELLER }
ICELAND BY P. MILES.
NORWAY, A RESIDENCE IN BY S. LAING.
NORWAY, RAMBLES IN BY T. FORESTER.
RUSSIA BY THE MARQUIS DE CUSTINE.
RUSSIA AND TURKEY .. BY J. R. MCCULLOCH.
ST. PETERSBURG BY M. JERRMANN.
THE RUSSIANS OF THE SOUTH, BY S. BROOKES.
SWISS MEN AND SWISS } BY R. FERGUSON.
MOUNTAINS }
MONT BLANC, ASCENT OF BY J. AULDJO.
SKETCHES OF NATURE } BY F. VON TSCHUDI.
IN THE ALPS }
VISIT TO THE VAUDOIS } BY E. BAINES.
OF PIEDMONT }

IN ASIA.

CHINA AND THIBET BY THE ABBE HUC.
SYRIA AND PALESTINE "EÖTHEN."
THE PHILIPPINE ISLANDS, BY P. GIRONIÈRE.

IN AFRICA.

AFRICAN WANDERINGS BY M. WERNE.
MOROCCO BY X. DURRIEU.
NIGER EXPLORATION, BY T. J. HUTCHINSON.
THE ZULUS OF NATAL BY G. H. MASON.

IN AMERICA.

BRAZIL BY E. WILBERFORCE.
CANADA BY A. M. JAMESON.
CUBA BY W. H. HURLBUT.
NORTH AMERICAN WILDS BY C. LANMAN.

IN AUSTRALIA.

AUSTRALIAN COLONIES BY W. HUGHES.

ROUND THE WORLD.

A LADY'S VOYAGE BY IDA PFEIFFER.

HISTORY AND BIOGRAPHY.

MEMOIR OF THE DUKE OF WELLINGTON.
THE LIFE OF MARSHAL } BY THE REV. T. O.
TURENNE COCKAYNE.
SCHAMYL BY BODENSTEDT AND WAGNER.
FERDINAND I. AND MAXIMILIAN II. } BY RANKE.
FRANCIS ARAGO'S AUTOBIOGRAPHY.
THOMAS HOLCROFT'S MEMOIRS.

CHESTERFIELD & SELWYN, BY A. HAYWARD.
SWIFT AND RICHARDSON, BY LORD JEFFREY.
DEFOE AND CHURCHILL BY J. FORSTER.
ANECDOTES OF DR. JOHNSON, BY MRS. PIOZZI.
TURKEY AND CHRISTENDOM.
LEIPSIĆ CAMPAIGN, BY THE REV. G. R. GLEIG.
AN ESSAY ON THE LIFE AND } BY HENRY
GENIUS OF THOMAS FULLER } ROGERS.

ESSAYS BY LORD MACAULAY.

WARREN HASTINGS.
LORD CLIVE.
WILLIAM PITT.
THE EARL OF CHATHAM.
RANKE'S HISTORY OF THE POPES.
GLADSTONE ON CHURCH AND STATE.
ADDISON'S LIFE AND WRITINGS.
HORACE WALPOLE.
LORD BACON.

LORD BYRON.
COMIC DRAMATISTS OF THE RESTORATION.
FREDERIC THE GREAT.
HALLAM'S CONSTITUTIONAL HISTORY.
CROKER'S EDITION OF BOSWELL'S LIFE OF JOHNSON.

LORD MACAULAY'S SPEECHES ON PARLIAMENTARY REFORM.

WORKS OF FICTION.

THE LOVE STORY, FROM SOUTHEY'S DOCTOR.
SIR ROGER DE COVERLEY, FROM THE }
SPECTATOR. }
MEMOIRS OF A MAITRE-D'ARMES, BY DUMAS.
CONFESSIONS OF A } BY E. SOUVESTRE.
WORKING MAN }

AN ATTIC PHILOSOPHER IN PARIS BY E. SOUVESTRE.
SIR EDWARD SEAWARD'S NARRATIVE OF HIS SHIPWRECK.

NATURAL HISTORY, &c.

NATURAL HISTORY OF } BY DR. L. KEMP.
CREATION }
INDICATIONS OF INSTINCT, BY DR. L. KEMP.

ELECTRIC TELEGRAPH, &c. BY DR. G. WILSON.
OUR COAL-FIELDS AND OUR COAL-PITS.
CORNWALL, ITS MINES, MINERS, &c.

MISCELLANEOUS WORKS.

LECTURES AND ADDRESSES { BY THE EARL OF
CARLISLE.
SELECTIONS FROM SYDNEY SMITH'S
WRITINGS.
PRINTING BY A. STARK.

RAILWAY MORALS AND } BY H. SPENCER.
RAILWAY POLICY }
MORMONISM .. BY THE REV. W. J. CONYBEARE.
LONDON BY J. R. MCCULLOCH.

The Traveller's Library being now complete, the Publishers call attention to this collection as well adapted for *Travellers* and *Emigrants*, for *School-room Libraries*, the *Libraries of Mechanics' Institutions*, *Young Men's Libraries*, the *Libraries of Ships*, and similar purposes. The separate volumes are suited for *School Prizes*, *Presents to Young People*, and for general instruction and entertainment. The Series comprises fourteen of the most popular of Lord Macaulay's *Essays*, and his *Speeches* on Parliamentary Reform. The department of *Travels* contains some account of eight of the principal countries of Europe, as well as travels in four districts of Africa, in four of America, and in three of Asia. *Madame Pfeiffer's First Journey round the World* is included; and a general account of the *Australian Colonies*. In *Biography and History* will be found Lord Macaulay's *Biographical Sketches of Warren Hastings, Clive, Pitt, Walpole, Bacon*, and others; besides *Memoirs of Wellington, Turenne, F. Arago, &c.*; an *Essay on the Life and Genius of Thomas Fuller*, with *Selections from his Writings*, by Mr. Henry Rogers; and a history of the *Leipsc Campaign*, by Mr. Gleig, — which is the only separate account of this remarkable campaign. Works of Fiction did not come within the plan of the *TRAVELLER'S LIBRARY*; but the *Confessions of a Working Man*, by Souvestre, which is indeed a fiction founded on fact, has been included, and has been read with unusual interest by many of the working classes, for whose use it is especially recommended. Dumas's story of the *Maitre-d'Armes*, though in form a work of fiction, gives a striking picture of an episode in the history of Russia. Amongst the works on Science and Natural Philosophy, a general view of Creation is embodied in Dr. Kemp's *Natural History of Creation*; and in his *Indications of Instinct* remarkable facts in natural history are collected. Dr. Wilson has contributed a popular account of the *Electric Telegraph*. In the volumes on the *Coal-Fields*, and on the *Tin* and other Mining Districts of *Cornwall*, is given an account of the mineral wealth of England, the habits and manners of the miners, and the scenery of the surrounding country. It only remains to add, that among the *Miscellaneous Works* are a Selection of the best Writings of the Rev. Sydney Smith; Lord Carlisle's *Lectures and Addresses*; an account of *Mormonism*, by the Rev. W. J. Conybeare; an exposition of *Railway management and mismanagement*, by Mr. Herbert Spencer; an account of the *Origin and Practice of Printing*, by Mr. Stark; and an account of *London*, by Mr. McCulloch.

"If we were called upon to lay the first stone of a Mechanics' Institute or Book-Society's Collection, it should be composed of the hundred and two parts of the *Traveller's Library*. It is the best Shilling Series extant. Here are Mr. Macaulay's best writings, the anthology of Sydney Smith, some admirable literary essays by different authors, several excellent volumes of science, narratives of travel in eight European, four American, four African, and three Asi-

atic countries, and examples from the works of Souvestre and Dumas. Bound together, they form twenty-five convenient volumes, which any society of a hundred and five members may possess, upon payment of one shilling each. An association of this kind, formed in every small town, would thus create sufficient basis for a free library upon a modest scale. Good books are not beyond the reach of working men, if working men will combine to obtain them."

LEADER.

☛ The *Traveller's Library* may also be had as originally issued in 102 parts, 1s. each, forming 50 vols. 2s. 6d. each; or any separate parts or volumes.

Trollope.—Barchester Towers. By ANTHONY TROLLOPE. 3 vols. post 8vo. price 31s. 6d.

"*Barchester Towers* (a kind of sequel in continuation of Mr. Trollope's former novel *The Warden*) does not depend only on story for its interest; the careful writing, the good humour with a tendency often to be Shandean in its expression, and the sense and right feeling with which the way is threaded among questions of high church and low church, are very noticeable, and secure for it unquestionable rank among the few really well-written tales that every season produces."

EXAMINER.

Trollope.—The Warden. By Anthony Trollope. Post 8vo. 10s. 6d.

Sharon Turner's Sacred History of the World, attempted to be Philosophically considered, in a Series of Letters to a Son. New Edition, edited by the Rev. S. TURNER. 3 vols. post 8vo. price 31s. 6d.

Sharon Turner's History of England during the Middle Ages: Comprising the Reigns from the Norman Conquest to the Accession of Henry VIII. Fifth Edition, revised by the Rev. S. TURNER. 4 vols. 8vo. price 50s.

Sharon Turner's History of the Anglo-Saxons, from the Earliest Period to the Norman Conquest. Seventh Edition, revised by the Rev. S. TURNER. 3 vols. 8vo. 36s.

Dr. Turton's Manual of the Land and Fresh-Water Shells of Great Britain. New Edition, thoroughly revised and brought up to the Present Time. Edited by Dr. J. E. GRAY, F.R.S., &c., Keeper of the Zoological Department in the British Museum. Crown 8vo. with Coloured Plates. [*In the press.*]

Dr. Ure's Dictionary of Arts, Manufactures, and Mines: Containing a clear Exposition of their Principles and Practice. Fourth Edition, much enlarged; most of the Articles being entirely re-written, and many new Articles added. With nearly 1,600 Woodcuts. 2 vols. 8vo. price 60s.

Van Der Hoeven's Handbook of Zoology. Translated from the Second Dutch Edition by the Rev. WILLIAM CLARK, M.D., F.R.S., &c., late Fellow of Trinity College, and Professor of Anatomy in the University of Cambridge; with additional References furnished by the Author. In Two Volumes. Vol. I. *Invertebrate Animals*; with 15 Plates, comprising numerous Figures. 8vo. 30s.

Vehse.—Memoirs of the Court, Aristocracy, and Diplomacy of Austria. By Dr. E. VEHSE. Translated from the German by FRANZ DEMMLER. 2 vols. post 8vo. 21s.

Von Tempsky. — Mitla: A Narrative of Incidents and Personal Adventures on a Journey in Mexico and Guatemala in the Years 1853 and 1854: With Observations on the Modes of Life in those Countries. By G. F. VON TEMPSKY. Edited by J. S. BELL, Author of *Journal of a Residence in Circassia in the Years 1836 to 1839*. With Illustrations in Chromolithography and Engravings on Wood. 8vo. [In the press.]

Wade. — England's Greatness: Its Rise and Progress in Government, Laws, Religion, and Social Life; Agriculture, Commerce, and Manufactures; Science, Literature, and the Arts, from the Earliest Period to the Peace of Paris. By JOHN WADE, Author of the *Cabinet Lawyer*, &c. Post 8vo. 10s. 6d.

Waterton. — Essays on Natural History, chiefly Ornithology. By C. WATERTON, Esq. With an Autobiography of the Author, and Views of Walton Hall. New and cheaper Edition. 2 vols. fcp. 8vo. price 10s.

Waterton's Essays on Natural History. Third Series; with a Continuation of the Autobiography, and a Portrait of the Author. Fcp. 8vo. price 6s.

Webster and Parkes's Encyclopædia of Domestic Economy; comprising such subjects as are most immediately connected with Housekeeping: As, The Construction of Domestic Edifices, with the Modes of Warming, Ventilating, and Lighting them—A description of the various articles of Furniture, with the nature of their Materials—Duties of Servants—&c. New Edition; with nearly 1,000 Woodcuts. 8vo. price 50s.

Weld. — Vacations in Ireland. By CHARLES RICHARD WELD, Barrister-at-Law. Post 8vo. with a tinted View of Birr Castle, price 10s. 6d.

Weld. — A Vacation Tour in the United States and Canada. By C. R. WELD, Barrister-at-Law. Post 8vo. with Map, 10s. 6d.

West. — Lectures on the Diseases of Infancy and Childhood. By CHARLES WEST, M.D., Physician to the Hospital for Sick Children; Physician-Accoucheur to, and Lecturer on Midwifery at, St. Bartholomew's Hospital. Third Edition. 8vo. 14s.

Willich's Popular Tables for ascertaining the Value of Lifehold, Leasehold, and Church Property, Renewal Fines, &c. With numerous additional Tables—Chemical, Astronomical, Trigonometrical, Common and Hyperbolic Logarithms; Constants, Squares, Cubes, Roots, Reciprocals, &c. Fourth Edition, enlarged. Post 8vo. price 10s.

Whitelocke's Journal of the English Embassy to the Court of Sweden in the Years 1653 and 1654. A New Edition, revised by HENRY REEVE, Esq., F.S.A. 2 vols. 8vo. 24s.

Wilmot's Abridgment of Blackstone's Commentaries on the Laws of England, intended for the use of Young Persons, and comprised in a series of Letters from a Father to his Daughter. 12mo. price 6s. 6d.

Wilson (W.). — Bryologia Britannica: Containing the Mosses of Great Britain and Ireland systematically arranged and described according to the Method of Bruch and Schimper; with 61 illustrative Plates. Being a New Edition, enlarged and altered, of the *Muscologia Britannica* of Messrs. Hooker and Taylor. By WILLIAM WILSON, President of the Warrington Natural History Society. 8vo. 42s.; or, with the Plates coloured, price £4. 4s. cloth.

Yonge. — A New English-Greek Lexicon: Containing all the Greek Words used by Writers of good authority. By C. D. YONGE, B.A. Second Edition, revised and corrected. Post 4to. price 21s.

Yonge's New Latin Gradus: Containing Every Word used by the Poets of good authority. For the use of Eton, Westminster, Winchester, Harrow, Charterhouse, and Rugby Schools; King's College, London; and Marlborough College. Fifth Edition. Post 8vo. price 9s.; or with APPENDIX of Epithets classified according to their English Meaning, 12s.

Youatt. — The Horse. By William Youatt. With a Treatise of Draught. New Edition, with numerous Wood Engravings, from Designs by William Harvey. (Messrs. LONGMAN and Co.'s Edition should be ordered.) 8vo. price 10s.

Youatt. — The Dog. By William Youatt. A New Edition; with numerous Engravings, from Designs by W. Harvey. 8vo. 6s.

Young. — The Christ of History: An Argument grounded in the Facts of His Life on Earth. By JOHN YOUNG, LL.D. Second Edition. Post 8vo. 7s. 6d.

Young. — The Mystery; or, Evil and God. By JOHN YOUNG, LL.D. Post 8vo. 7s. 6d.

Zumpt's Grammar of the Latin Language. Translated and adapted for the use of English Students by DR. L. SCHMITZ, F.R.S.E.: With numerous Additions and Corrections by the Author and Translator. 4th Edition, thoroughly revised. 8vo. 14s.

[October 1857.]

CORNELL UNIVERSITY LIBRARY

JUL 18 1991

MATHEMATICS LIBRARY

